

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Should Postanoxic Status Epilepticus be Treated Aggressively?-
No!
Authors: Rossetti AO
Journal: Journal of clinical neurophysiology : official publication of the
American Electroencephalographic Society
Year: 2015 Dec
Volume: 32
Issue: 6
Pages: 447-8
DOI: 10.1097/WNP.0000000000000208

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Should postanoxic Status epilepticus be treated aggressively?

- No!

Andrea O. Rossetti, MD¹

¹ Department of Clinical Neurosciences, Centre Hospitalier Universitaire Vaudois (CHUV) and University of Lausanne, Lausanne, Switzerland.

Acknowledgment: The author received unrestricted research contributions from Sage Therapeutics, Eisai, UCB, and is supported by the Swiss National Science Foundation (grant nr. CR32I3_143780).

Contents:

Title: 66 characters

Abstract: 100 words

Text: 1180 words

28 References

Keywords: General anesthetics, outcome, complications, myoclonus, EEG, reactivity.

Address correspondence to:

Dr Andrea O. Rossetti

Service de Neurologie

CHUV-BH07

CH-1011 Lausanne, Switzerland

Phone: +41 21 314 1220

Fax: +41 21 314 1290

andrea.rossetti@chuv.ch

Abstract

Electrographic Status epilepticus (SE) and myoclonus represent frequent findings in patients surviving cardiac arrest; both features have been related to poor clinical outcome. Recent data have outlined that SE appearing during therapeutic hypothermia and sedation is practically invariably related to a fatal issue, as opposed to some patients presenting SE and/or myoclonus after return to normothermic conditions. While it seems reasonable to give a chance of awakening to the latter patients by administering consequent antiepileptic treatment, especially if other favorable prognostic markers are observed, an aggressive treatment of SE arising during hypothermia seems futile in view of the existing evidence.

Background

Despite recent advances in resuscitation and intensive care in the domain of hypoxic-ischemic encephalopathy after cardiac arrest [1], only about 5% of patients survive to hospital discharge [2], and several patients are left with long-term cognitive sequelae [3]. Outcome prediction in this clinical setting represents a daunting task for neurologists and intensivists; therefore, guidelines have been proposed, in which clinical examination and EEG represent paramount tools. As researchers have been particularly successful in refining prognostication for dismal prognosis [4], false-positive rates (FPR) for poor outcome prediction (Cerebral Performance Category of 3 or higher on a scale of 5 [5]) should aim towards 0, with narrow confidence intervals.

In 2006, the American Academy of Neurology published evidence-based practice parameters [6]. Early myoclonus within 24h of cardiac arrest (FPR=0), and an EEG with generalized suppression, burst-suppression, or periodic discharges on a suppressed background (FPR<5%), were reported as reliable markers of unfavorable prognosis (Level C); however, this analysis included almost exclusively subjects treated before the hypothermia era.

Current evidence

More recently, several studies conducted in different cohorts in Europe and North America have addressed the prognostic role of electrographic epileptiform abnormalities and clinical myoclonus in patients treated with controlled temperature management. For the purposes of this controversy, status epilepticus (SE) will be defined broadly as electroencephalographic evidence of repetitive (rhythmic, periodic, evolving) epileptiform transients over more than 5 minutes, and “aggressive” treatment as the use of prolonged pharmacological sedation to induce electrographic burst-suppression or complete suppression.

After therapeutic hypothermia, electrographic seizures or repetitive epileptiform patterns are found in about 1/3 of adult patients [7-9], while early myoclonus (arising after weaning of sedation and hypothermia) has been reported in about 1/6 of patients [10, 11], and thus represent relatively frequent observations in this setting. Several independent investigators have found that the presence of EEG epileptiform alterations suggesting ongoing electrical seizures is incompatible with survival (FPR=0), even if patients were treated with antiepileptic drugs (AED) [9, 12-14].

However, several patients displaying postanoxic SE have in fact been reported to recover with at times a reasonable functional outcome [15, 16]. How is this to understand in view of the previous paragraph? To address this, at first glance, discrepancy, one should target her or his attention towards the early phase of intensive care, in which patients under hypothermia may undergo

EEG. Indeed, the extent of hypothermic EEG abnormalities (particularly, lack of background reactivity and discontinuity) has been shown to correlate with the magnitude of neuronal damage assessed with the neuron-specific enolase (NSE), a marker of neuronal death [17, 18]. A neurophysiologic, non-invasive tool may therefore prove highly informative, especially if recorded just after the first 9-12 hours following the initial insult [19].

During therapeutic hypothermia, recent evidence on 134 consecutive patients highlighted that the presence of epileptiform EEG activity is indeed not compatible with survival, even under continuous infusion of moderate doses of midazolam or propofol, and after additional AED treatment (FPR 0%); this contrasts with a FPR of 7% in patients displaying epileptiform activity only after return to normothermia, or the FPR of 13% for overt myoclonus within the first 48-72 hours [10]. Of note, the lack of EEG background reactivity during hypothermia was also clearly related to a poor prognosis (FPR 2%) [10]. These findings are similar to those reported by another group on 56 subjects investigated with continuous EEG during hypo- and normothermia: epileptiform transients were found in 10% of the 22 patients with good neurological outcome, at 48 hours (thus after hypothermia), but not under hypothermia, and lasted less than 2 hours under AED treatment [20]. A recent study focusing on postanoxic myoclonus highlighted that survival is clearly more frequent in patients without concomitant EEG epileptiform activity [11]. Moreover, an earlier Scandinavian study conducted with amplitude-integrated EEG on 95 patients demonstrated that among the 26 subjects with electrographic SE, the 2 who survived presented this condition only after return of normothermia, and out of a continuous background activity, but not burst-suppression [21]. Finally, occurrence of rhythmic, periodic, or ictal epileptiform features induced by stimulations, called “SIRPIDs” was recently assessed in a relatively large patient cohort: these were found in 13% of patients, and their occurrence during hypothermia was not compatible with survival [22].

Conclusions

The aforementioned data seem to converge to the following three points. First, while electrographic epileptiform features, which nearly always present in a rhythmic or repetitive fashion at least for several hours [23], thus fulfilling the SE definition, are general markers of poor prognosis after cardiac arrest, their occurrence during the early resuscitation period, particularly under mild therapeutic hypothermia and sedation, has not been so far reported to be compatible with survival; this is in contrast with some patients who may benefit from AED treatment if their SE declares itself after return to normothermic conditions. It may be speculated that SE arising in hypothermia despite sedation is the hallmark of an agonal cortical activity resulting from a devastating brain insult [10], whereas SE appearing in normothermia may be sometimes amenable to treatment and a reasonable outcome. Second, EEG background reactivity represents a practical additional prognostic marker: its absence during hypothermia seems to

have a smaller FPR for poor outcome, as compared to normothermic conditions (FPR=7-9%) While EEG reactivity in this setting has recently been shown to have reasonable inter-rater agreements among EEG readers, and even with automated algorithms [24], another study found lower agreements [25]. Third, myoclonus, which is also consistently related to poor outcome, is nevertheless not an invariable marker of mortality; of note, sedation and muscle relaxants often prevent its observation during controlled temperature management.

On top of these considerations, risks inherent to aggressive SE treatment, mostly related to potential complications of prolonged ICU treatment, sedation, and mechanical ventilation, have been outlined in recent literature, albeit outside of the setting of post-anoxic encephalopathy [26-28].

It appears therefore reasonable to suggest the following approach in patients with hypoxic ischemic encephalopathy. If EEG features compatible with SE arise only after hypothermia and upon sedation weaning, on a reactive EEG in an otherwise potentially promising clinical context (e.g., return of brainstem reflexes, preserved cortical somatosensory evoked responses), a treatment with non-sedating and at times sedating AED seems warranted, at least for some days. If, conversely, SE appears during sedation and hypothermia on a non-reactive EEG background, aggressive antiepileptic treatment with long-term sedative agents appears futile, as can worsen comorbidities, may divert resources from patients who may better take advantage from intensive treatment, and might unnecessarily induce false expectations in relatives and caregivers.

References

1. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, Horn J, Hovdenes J, Kjaergaard J, Kuiper M *et al*: **Targeted temperature management at 33 degrees C versus 36 degrees C after cardiac arrest.** *The New England journal of medicine* 2013, **369**(23):2197-2206.
2. Stiell IG, Wells GA, Field B, Spaite DW, Nesbitt LP, De Maio VJ, Nichol G, Cousineau D, Blackburn J, Munkley D *et al*: **Advanced cardiac life support in out-of-hospital cardiac arrest.** *The New England journal of medicine* 2004, **351**(7):647-656.
3. Moulaert VR, Verbunt JA, van Heugten CM, Wade DT: **Cognitive impairments in survivors of out-of-hospital cardiac arrest: a systematic review.** *Resuscitation* 2009, **80**(3):297-305.
4. Sandroni C, Cariou A, Cavallaro F, Cronberg T, Friberg H, Hoedemaekers C, Horn J, Nolan JP, Rossetti AO, Soar J: **Prognostication in comatose survivors of cardiac arrest: an advisory statement from the European Resuscitation Council and the European Society of Intensive Care Medicine.** *Intensive care medicine* 2014, **40**(12):1816-1831.
5. Booth CM, Boone RH, Tomlinson G, Detsky AS: **Is this patient dead, vegetative, or severely neurologically impaired? Assessing outcome for comatose survivors of cardiac arrest.** *Jama* 2004, **291**(7):870-879.
6. Wijdicks EF, Hijdra A, Young GB, Bassetti CL, Wiebe S: **Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology.** *Neurology* 2006, **67**(2):203-210.
7. Rossetti AO, Oddo M, Logroscino G, Kaplan PW: **Prognostication after cardiac arrest and hypothermia: a prospective study.** *Annals of neurology* 2010, **67**(3):301-307.
8. Legriel S, Hilly-Ginoux J, Resche-Rigon M, Merceron S, Pinoteau J, Henry-Lagarrigue M, Bruneel F, Nguyen A, Guezennec P, Troche G *et al*: **Prognostic value of electrographic postanoxic status epilepticus in comatose cardiac-arrest survivors in the therapeutic hypothermia era.** *Resuscitation* 2013, **84**(3):343-350.
9. Knight WA, Hart KW, Adeoye OM, Bonomo JB, Keegan SP, Ficker DM, Szaflarski JP, Privitera MD, Lindsell CJ: **The incidence of seizures in patients undergoing therapeutic hypothermia after resuscitation from cardiac arrest.** *Epilepsy research* 2013, **106**(3):396-402.
10. Oddo M, Rossetti AO: **Early multimodal outcome prediction after cardiac arrest in patients treated with hypothermia.** *Crit Care Med* 2014, **42**(6):1340-1347.
11. Seder DB, Sunde K, Rubertsson S, Mooney M, Stammet P, Riker RR, Kern KB, Unger B, Cronberg T, Dziodzio J *et al*: **Neurologic outcomes and postresuscitation care of patients with myoclonus following cardiac arrest.** *Crit Care Med* 2015, **43**(5):965-972.
12. Fugate JE, Wijdicks EF, Mandrekar J, Claassen DO, Manno EM, White RD, Bell MR, Rabinstein AA: **Predictors of neurologic outcome in hypothermia after cardiac arrest.** *Annals of neurology* 2010, **68**(6):907-914.
13. Thomke F, Weilemann SL: **Poor prognosis despite successful treatment of postanoxic generalized myoclonus.** *Neurology* 2010, **74**(17):1392-1394.
14. Crepeau AZ, Rabinstein AA, Fugate JE, Mandrekar J, Wijdicks EF, White RD, Britton JW: **Continuous EEG in therapeutic hypothermia after cardiac arrest: prognostic and clinical value.** *Neurology* 2013, **80**(4):339-344.
15. Rossetti AO, Oddo M, Liaudet L, Kaplan PW: **Predictors of awakening from postanoxic status epilepticus after therapeutic hypothermia.** *Neurology* 2009, **72**(8):744-749.

16. Westhall E, Rundgren M, Lilja G, Friberg H, Cronberg T: **Postanoxic status epilepticus can be identified and treatment guided successfully by continuous electroencephalography.** *Therapeutic hypothermia and temperature management* 2013, **3**(2):84-87.
17. Cronberg T, Rundgren M, Westhall E, Englund E, Siemund R, Rosen I, Widner H, Friberg H: **Neuron-specific enolase correlates with other prognostic markers after cardiac arrest.** *Neurology* 2011, **77**(7):623-630.
18. Rossetti AO, Carrera E, Oddo M: **Early EEG correlates of neuronal injury after brain anoxia.** *Neurology* 2012, **78**(11):796-802.
19. Alvarez V, Sierra-Marcos A, Oddo M, Rossetti AO: **Yield of intermittent versus continuous EEG in comatose survivors of cardiac arrest treated with hypothermia.** *Crit Care* 2013, **17**(5):R190.
20. Cloostermans MC, van Meulen FB, Eertman CJ, Hom HW, van Putten MJ: **Continuous electroencephalography monitoring for early prediction of neurological outcome in postanoxic patients after cardiac arrest: a prospective cohort study.** *Crit Care Med* 2012, **40**(10):2867-2875.
21. Rundgren M, Westhall E, Cronberg T, Rosen I, Friberg H: **Continuous amplitude-integrated electroencephalogram predicts outcome in hypothermia-treated cardiac arrest patients.** *Crit Care Med* 2010, **38**(9):1838-1844.
22. Alvarez V, Oddo M, Rossetti AO: **Stimulus-induced rhythmic, periodic or ictal discharges (SIRPIDs) in comatose survivors of cardiac arrest: characteristics and prognostic value.** *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology* 2013, **124**(1):204-208.
23. Thomke F, Marx JJ, Sauer O, Hundsberger T, Hagele S, Wiechelt J, Weilemann SL: **Observations on comatose survivors of cardiopulmonary resuscitation with generalized myoclonus.** *BMC Neurol* 2005, **5**:14.
24. Noirhomme Q, Lehembre R, Lugo Zdel R, Lesenfants D, Luxen A, Laureys S, Oddo M, Rossetti AO: **Automated analysis of background EEG and reactivity during therapeutic hypothermia in comatose patients after cardiac arrest.** *Clinical EEG and neuroscience : official journal of the EEG and Clinical Neuroscience Society* 2014, **45**(1):6-13.
25. Westhall E, Rosén I, Rossetti AO, van Rootselaar AF, Kjaer TW, Friberg H, Horn J, Nielsen N, Ullén S, Cronberg T: **Interrater variability of EEG interpretation in comatose cardiac arrest patients.** *Clinical Neurophysiology* 2015.
26. Kowalski RG, Ziai WC, Rees RN, Werner JK, Jr., Kim G, Goodwin H, Geocadin RG: **Third-line antiepileptic therapy and outcome in status epilepticus: the impact of vasopressor use and prolonged mechanical ventilation.** *Crit Care Med* 2012, **40**(9):2677-2684.
27. Sutter R, Marsch S, Fuhr P, Kaplan PW, Ruegg S: **Anesthetic drugs in status epilepticus: Risk or rescue?: A 6-year cohort study.** *Neurology* 2014, **82**(8):656-664.
28. Marchi NA, Novy J, Faouzi M, Stahli C, Burnand B, Rossetti AO: **Status epilepticus: impact of therapeutic coma on outcome.** *Crit Care Med* 2015, **43**(5):1003-1009.