HISTORY OF AN OUTBREAK OF PSEUDO-DIPHTHERIA.*

By JOSEPH PRIESTLEY, B.A., M.D., D.P.H.,

Medical Officer of Health for the Metropolitan Borough of Lambeth.

My attention was drawn by the medical superintendent of a large boarding school in Lambeth to an outbreak of sore throats—slight throat troubles having been more or less prevalent for many months past amongst the scholars, but no case having been sufficiently marked in its clinical symptoms to call for interference previously. On examination, the clinical symptoms were found to be as follows:

Increase of temperature, headache, lassitude, pains over the body generally, and difficulty and pain in swallowing. The majority of the cases were mild, and the patients recovered, generally after short illnesses, and without any sequelæ or complications, with the exception of slight enlargements of the glands of the throat and neck (the tonsils and the submaxillary glands). The pharynx was slightly congested in each case, and the membrane, when present, as it was often, was in the form of isolated, irregular-shaped spots, varying in size from a pin-point to a pea. This membrane was readily detached, and left no bleeding surface. In no single instance was there any suspicion of subsequent paralysis of any of the throat or other muscles. No death occurred. Swabs were at once taken from several typical throats, and in all these cases cultivations of definite, characteristic bacilli were obtained, mostly in pure cultivation. These bacilli were short rods with round ends, 1μ to 1.5μ in length, and stained regularly and deeply with Löffler's methylene-blue, polar staining being conspicuous by its absence. The parallel grouping (so conspicuous in the case of Klebs-Löffler bacilli) was marked, but no involution forms were met with, the bacilli appearing to be all of the same uniform size, and slightly thicker at the centres than at the ends. The bacillus grew well on blood serum, forming in twenty-four to forty-eight hours large cream-coloured colonies or growths, indistinguishable by the naked eye from those consisting of the true Klebs-Löffler (diphtheria) bacilli. The reaction of the colonies or growths in glucose was alkaline. It was clear, therefore, that we had to deal with an outbreak of sore throats due to a bacillus, which in some points resembled the well-known Klebs-Löffler bacillus, but which

* Read at meeting of Metropolitan Branch, May 22nd, 1903.

in many other points offered marked differences therefrom. These bacilli were, in fact, the pseudo (diphtheria) bacilli of Hofmann, and totally distinct from the true (diphtheria) bacilli of Klebs and Löffler. This fact was confirmed by several independent bacteriologists. Systematic bacteriological examinations were made from time to time of swabs taken from the scholars' throats, with the result that, out of a total of 137 scholars, 38 throats showed the presence of the pseudo-bacilli of Hofmann, and 5 the presence of the true bacilli of Klebs and Löffler, *i.e.*, 27·7 per cent. and 3·6 per cent. respectively. Six throats showed staphylococci and streptococci only. It is interesting to note that only one adult (the porter) was affected out of a total of 28 adults (teachers and servants) employed in the school, *i.e.*, 3·6 per cent., and the bacillus found in this case was the pseudo-bacillus of Hofmann. The grouping of the cases showed that the disease was spread chiefly in the dormitories (which at the time were found to be somewhat overcrowded), and the means of the spreading was probably the transference of the bacilli from person to person by contact, directly through kissing, or indirectly through towels, glasses, etc. In this connection the girls were found to be proportionately most affected. The first introduction of the disease into the school was from without by some scholar or scholars on returning from their summer holidays on September 3rd, the first case being noticed on September 7th, followed by forty fresh cases discovered during the same month.

It was decided that the pseudo cases should be treated as infectious and isolated, and in this respect, fortunately, the school was provided with a sanatorium away from the main school buildings, and into this sanatorium all pupils, whose throats were found, on examination, to contain pseudo-bacilli, were at once removed, and kept isolated there until all traces of pseudo-bacilli were gone, as shown by negative results from repeated bacteriological examinations made. Antiseptic sprays were used regularly and systematically as part of the treatment (the most efficacious spray in the opinion of the medical superintendent being a 0·1 per cent. solution of formalin). The scholars were moved back to the main school buildings (1) when they were convalescent; (2) as already stated, when repeated bacteriological examinations showed that the pseudo-bacilli were no longer present in their throats, and (3) *after efficient disinfection of their clothes (and persons) had been carried out.*

The lesson to be learnt from the outbreak is sufficiently obvious, *viz.*, that pseudo-bacilli are communicable from person to person, and that throats containing pseudo-bacilli are (at times at least) infectious. The usual preventive measures, consequently, suggest

themselves, viz., notification, isolation, and disinfection. All patients suffering from pseudo-diphtheritic throats (with clinical symptoms) should be isolated, but I do not think that, in the present state of bacteriological knowledge, we should be justified in placing patients with throats infected with so-called pseudo-diphtheria bacilli in the same wards with patients suffering from true diphtheria bacilli. It is true that some bacteriologists regard the pseudo-bacillus of Hofmann as a modified (attenuated) form of the Klebs Löffler bacillus, and the five true diphtheria cases that were found along with the thirty-eight pseudo-diphtheria cases in the Lambeth outbreak (in one case the true bacilli being afterwards found grafted on a pseudo throat) would appear, *primâ facie*, to support such a view. I think, however, reviewing the Lambeth outbreak as a whole, and remembering that only one case developed true bacilli after pseudo-bacilli, that the two diseases are distinct, and that the one may be grafted upon the other in one and the same patient at one and the same time. The two diseases should not, therefore, I submit, be treated in the same ward, and, as a corollary, it follows that an infectious hospital should be provided with a separate ward for pseudo-diphtheria patients. If the medical superintendents of the Metropolitan Asylums Board share my view, something might be done in the way of providing such pseudo-diphtheria separate isolation accommodation. I merely offer the suggestion. In connection with the local outbreak with which I am now dealing, the five true diphtheria cases were removed to hospital, and the thirty-eight pseudo-diphtheria cases isolated in the separate sanatorium belonging to the school—a most useful and all-important additional building to boarding-school premises. The whole of the suspicious cases were, by bacteriological examination, discovered, separated out, and kept under observation for periods varying from one to five weeks. The end justified the means employed, as the outbreak was stamped out, and that, too, without it being necessary to close the school as a whole, and distribute the scholars over various parts of London (and elsewhere), thereby risking the spreading of the disease in many directions.

An interesting fact in connection with the outbreak may be mentioned before a society of medical officers of health. In examining some of the sanitary fittings of the school in the lavatories adjoining the dormitories, the waste-pipes of the lavatory basins, and the outlets of the open rectangular troughs underneath the basins, were found to be covered with a slimy deposit, consisting (as proved on examination) of organic and other débris (skin scales,

washings from mouths, hairs, cotton and woollen fluff, soap, etc.), together with pseudo (Hofmann) and true (Klebs-Löffler) bacilli. Cultivations of both were obtained by independent observers by simply inoculating serum tubes with portions of this slimy deposit. The lavatories, it may be mentioned, were used by the scholars, not only for the usual purposes of washing hands and faces, but also for washing teeth and mouths. The slimy deposit was a good nutrient medium, which was kept at the necessary suitable temperature by the hot water—used for washing at all hours of the day—the hot water trickling gently over the deposits in the waste-pipes and open troughs, which latter were made of both galvanized iron and wood. These open rectangular troughs were condemned and done away with, a course of action which I should advise for other schools that may be fitted with such appliances. At the same time, it is only fair to state that there was no reason for supposing that these infected waste-pipes and open rectangular troughs caused the disease to spread, as the germs found in the deposits were practically harmless at the time, being imprisoned on the moist surface, which was, by constant use, never allowed to dry. Had the deposit dried and been carried about as dust, the results might have been different.

This short history of an outbreak of pseudo-diphtheria in Lambeth is a good object-lesson, and as the medical officer of health who investigated the facts, I have no hesitation in asking my colleagues to endorse officially, in the name of the Metropolitan Branch, the following principle—viz.: "That, in the opinion of the branch, cases clinically resembling diphtheria, but in which pseudo-bacilli, and not Klebs-Löffler bacilli, are found, are infectious, and that, consequently, the usual precautionary measures should be taken in connection therewith—viz., notification, isolation, and disinfection."

DISCUSSION.

Dr. A. WELLESLEY HARRIS said he was of opinion that cases of pseudo-diphtheria were infectious, and should be isolated with similar precautions to true diphtheria. He was not prepared to go as far as Dr. Priestley and recommend the provision of special wards in hospitals for these cases. In December last year he attempted to ascertain the cause of a threatened outbreak of diphtheria in a large industrial school. With the assistance of the regular medical attendant some 250 boys were examined, and of these fourteen had congested throats, but no sign of diphtheria. Swabs from each throat were submitted to bacteriological examination. Two were reported as containing diphtheria, three streptococci, and nine pseudo-diphtheria. The diphtheria cases were removed to hospital, and a special isolation obtained in the school for the pseudo-diphtheria cases, with the result that the occurrence of fresh

cases ceased. He attributes great value to the isolation of pseudo-diphtheria cases from healthy individuals.

Dr. SIDNEY DAVIES stated that in November last sixteen cases of diphtheria were notified in a Board school. An examination of the children attending this school was made by Dr. Thomas, assistant medical officer London School Board, with the result that sixteen cases were picked out for bacteriological examination, having suspicious symptoms of either the throat or the nose. One of these was found to have obvious clinical diphtheria. Cultures were taken from the others, with the result that one was found to have diphtheria bacilli and four pseudo-diphtheria bacilli, the bacilli being found in all cases in the nasal discharge. With the exclusion of the above-mentioned children, the outbreak in the school came to an end. He has been unable to detect any difference in the symptoms of children suffering from nasal discharges between those having the diphtheria bacilli and those having the pseudo-diphtheria bacilli. If the pseudo-diphtheria bacilli were decided to be indicative of an infectious disease, distinct from diphtheria, and that for the disease the same steps were to be taken with regard to notification and isolation as for diphtheria, this was equivalent to adding another disease to the list of dangerous infectious diseases in the Public Health Act, and would require special steps taken in order to obtain notification, etc. His own opinion, from clinical observation, was that the pseudo-diphtheria bacilli should not be distinguished from the true diphtheria bacilli, but should be considered as indicating a mild type of true diphtheria.

Dr. BROWN, after expressing appreciation of Dr. Priestley's interesting account of the outbreak due to Hofmann's bacillus, and the great importance of the subject of the connection of Hofmann with true bacillus, expressed the opinion that the present evidence tends to show that this bacillus is really an attenuated form of the Klebs-Löffler bacillus. For differentiation of this bacillus he relied solely on morphology, and agreed that what he was accustomed to call Hofmann's bacillus was a short wedge-shaped bacillus of very uniform size, occasionally with the bases of the wedges approximated, that it stained deeply and uniformly with carbol methylene blue, and showed no polar staining with Neisser. In support of the view that it is an attenuated form of Klebs-Löffler, he mentioned the experiments of Salter and Richmond at the Jenner Institute, by which the Hofmann was converted into the Klebs-Löffler bacillus by passage through birds, and was not aware, till informed at that meeting, that an unsuccessful effort to repeat them had been undertaken at the Jenner Institute. The experiments of Hewlett and Knight also seemed to point to the possibility of converting one into the other by cultural methods. The very fact brought out by Dr. Priestley, that the bacillus in question possessed a decided though slight virulence, and was associated in three of the cases with true Klebs-Löffler bacillus, tended to prove the same thing. He stated that during the year 1902 he had visited families where 102 primary cases of diphtheria occurred, in which the throats of 259 contacts were examined bacteriologically for Klebs-Löffler bacillus. This number, added to swabs sent in by general practitioners, brought the total number of throats examined for Klebs-Löffler bacillus up to 412. In eighty cases Hofmann's bacillus was found either alone or with Klebs-Löffler bacillus. In twenty-one of this eighty, or 26 per cent., it was associated with Klebs-Löffler bacillus. In fifty-nine of this eighty, or 74 per cent., it occurred alone or with cocci of various sorts.

The majority of the throats containing Hofmann alone presented no clinical symptoms, but in 10 to 20 per cent. some clinical symptoms were seen, such as redness, on examining the throat. From this it would appear that, first, Hofmann is very frequently associated with Klebs-Löffler bacillus, and, secondly, that it is fairly prevalent among children. The following are a few examples of the kind of cases met during the examination of the 259 diphtheria contacts :

On February 19th, 1902, W. D., male, seven years, was notified as having diphtheria ; on following day took swabs from children in the family, and found that—

B. D., six months, female, no Klebs-Löffler bacillus, but cocci and Hofmann.

D. D., five years, male, ditto.

L. D., two years, female, Klebs-Löffler bacillus and Hofmann.

L. D., four years, female, no Klebs-Löffler bacillus, a few Hofmann.

On March 4th took second swab from L. D., two years, and found no Klebs-Löffler bacillus.

In several of these cases what looked like transitional forms between Hofmann and Klebs-Löffler bacillus occurred, and it was often difficult to distinguish which one had to deal with.

Baby M., a diphtheria contact, lived in Holcomb Buildings, Bermondsey, and had pure culture of Hofmann. On June 28th she was notified with genuine diphtheria from Nunhead, where she had been taken to friends just fourteen days after the pure Hofmann, and no sore throats were found. What happened in these fourteen days when Klebs-Löffler bacillus developed ? There had been no other recent case where she went to at Nunhead.

From these considerations, I always recommend the parents of children with Hofmann bacillus in their throats to call in their medical man and get an antiseptic gargle. I also endeavour to isolate them at home, though this is very difficult, and at any rate exclude them from school. At present the Metropolitan Asylums Board will not take such cases in, and until the whole relationship has been decided, I certainly think it would be best to isolate and treat them practically as mild cases of diphtheria.

Dr. GOODALL thought it would be well to drop the name "pseudo-diphtheria." He did not consider that the Hofmann had ever been converted into the Klebs-Löffler.

Dr. CALDWELL SMITH would like to see notified all cases of sore throat accompanied by a rise of temperature, as the early knowledge gained thereby would have prevented several serious outbreaks which had come under his observation.

INTERNATIONAL HOUSING CONGRESS.—This Congress is to be held from July to November at the Grand Palace, in the Champs Elysées. The object of the Congress is to show the world the great need for improvement in both rural and city homes for workmen and clerks, and to this end there will be a collection from many countries of models of improved houses, of materials and models of construction, and of furniture and decorations. There will also be models to show the sanitary, or, rather, unsanitary, conditions in factories and many other places in which men work. It is expected that the work of the Congress will exert a potent influence in the control of consumption and alcoholism.