felt at their site. There was also marked burning pain throughout the entire length of the alimentary canal, and erythematous patches were seen upon the gums. All these manifestations disappeared within three or four days.—*Bulletin Générale de Thérapeutique*, 1898, 13e liv. p. 492.

Bacteriuria.—DR. EDMUND BONN reports an instance of chronic deep urethritis and follicular prostatitis. He administered internally sodium salicylate, benzoic acid, and salol, with large quantities of water (three pints to two quarts), and finally irrigated the bladder with potassium permanganate (1 to 1000) and corrosive sublimate (1 to 10,000). These were unavailing. The best results were obtained by the use of diuretic infusions and fifteen to thirty grains of urotropin daily, although even this did not entirely clear the urine.—*Prager medicinische Wochenschrift*, 1898, No. 18, S. 208.

The Applications of the Gastric Juice of the Dog.—M. FRÉMONT has obtained benefit from this substance in cholera nostras, chronic enteritis, and old dyspepsias. Several excessively emaciated patients gained weight rapidly. This remedy is contraindicated in the presence of an excessive amount of hydrochloric acid and in cancer. All of the patients treated had been for some time upon a regulated diet without benefit. Suggestion would not explain the beneficial results because the majority of patients did not know what they were taking. The juice was administered during the meal either clear or in beer. The doses varied according to the acidity of the stomach-contents of the patients. The average dose was two ounces.—*Le Progrès Médical*, 1898, No. 18, p. 281.

MEDICINE.

UNDER THE CHARGE OF

WILLIAM OSLER, M.D.,

PROFESSOR OF MEDICINE IN THE JOHNS HOPKINS UNIVERSITY, BALTIMORE, MARYLAND,

AND

GEORGE DOCK, M.D.,

PROFESSOR OF MEDICINE IN THE UNIVERSITY OF MICHIGAN.

Cancer of the Œsophagus.—HUBER has reported a case with some unusual features. The patient, a restaurateur, aged fifty-five years, began to complain of weakness and of piercing pain on swallowing. For years he had had attacks of an anginose character, frequently associated with hystero-epileptic paroxysms provoked by acute alcoholism. The patient was weak and emaciated, the pupils were contracted, and reacted to accommodation, but not to light. The reflexes were normal. Solid food could not be swallowed. On attempting to swallow milk, a coughing paroxysm was excited and most of the milk vomited. Examination of the pharynx showed a dilatation of the lower end, containing food-remains and mucus. The right vocal cord was