TUBERCULOSIS OF RETINA.

By A. C. O'SULLIVAN, F.T.C.D.;
Lecturer on Pathology, Trinity College;

AND

J. B. STORY, F.R.C.S.;
Surgeon to St. Mark's Hospital;
Professor of Ophthalmic and Aural Surgery, Royal College of Surgeons.

[Read in the Section of Pathology, November 4, 1898.]

THE specimen was taken from a healthy-looking, well-developed girl of twenty-one, who was sent to me by my friend Dr. Powell, of Roscrea, in February, 1898. The only history to be obtained was that a cloud came over her right eye three months previously, and that it came on in one night. Her health was good, and had always been so, and there was no sign of disease or dyscrasia.

On examination the pupil of the right eye only acted consensually, and the vision of that eye was reduced to counting fingers, and that only in the nasal side of the field. The left eye was normal in all respects. Ophthalmoscopically the right eye presented the appearances of an intense papillitis, but differing from any such inflammation I have ever seen in the brilliant whiteness of the papilla, and in the extreme swelling, which was more marked than in any case in my past experience. The prominence and the whiteness combined made the appearances much resemble those seen in detachment of the retina. The blood vessels, however, in the distended whitened papilla were tortuous and dilated, and there were small white spots in the retina about the macular region. A definite diagnosis was not made at that time, but when I saw the case again in April we diagnosed tuberculosis, and persuaded the patient to allow the globe to be removed. In April she said the eye had been inflamed for a fortnight, and there was a pink circumcorneal vascularity, a discolouration of the iris, and an external strabismus. Tension normal, and vision just perception of light.

After hardening the globe the following conditions were found:—

At the posterior part of the retina, for some distance round the entrance of the optic nerve, there was a tumour one-third of an inch in diameter, and about one-fifth inch in depth. For some distance on both sides of this tumour the retina was detached, and a firm coagulum of a homogeneous appearance lay between it and the underlying chorioid coat. On examining the section with the microscope, the tumour was seen to be composed of typical tubercular tissue containing numerous small round cells, larger irregular cells and giant cells. It was traversed by strands of connective tissue containing blood vessels, and showed a slight caseation in parts.

This tissue passed over without a break into the inner layers of the retina, which contained much larger vessels than usual, and in which were scattered numerous small nodules containing lymphoid and endothelioid cells in a reticulum,

The illustration shows the microscopic appearance of the tumour on cutting open the globe.

which caused the retina to bulge in towards the vitreous, and were no doubt tubercular in character. The tubercular tumour was separated from the chorioid by the coagulum before mentioned, which under the microscope showed a homogeneous appearance, and stained like the colloid material of the thyroid gland.

Behind this material all round lay the outer or pigmentary layer of the retina, and then the chorioid; so that the so-called retinal detachment is really a separation of the retinal layers by the exudation.

The optic nerve behind the globe was, microscopically, perfectly normal. The disease evidently was spreading from the papilla to the anterior portion of the uveal tract, and not backwards.

Localised tuberculosis of the uveal tract is not a very uncommon occurrence in ophthalmic literature. Many cases have been recorded, and the first case of the present communication only adds another to an already extensive series.

Localised tuberculosis of the retina is, however, a very rare occurrence indeed, and it appears probable that the second case described is actually the first recorded of a purely ophthalmoscopic diagnosis of such a condition. Tuberculosis of the optic nerve, with or without implication of the retina, has been described in a small number of cases. In all cases of localised tuberculosis of the eyeball it is very difficult to determine whether the disease is a primary intra-ocular affection or merely secondary to tuberculosis elsewhere. In most of the recorded cases the eye affection was undoubtedly secondary, and in the remainder of the cases it is almost impossible to prove that it was not secondary. In the case here described the disease appears to be primary, but it would be rash to make a positive assertion on the subject.

A very interesting case is recorded by Sattler in *v. Graefe's Archives*, XXIV. 3, p. 127. It shows what would probably have been the result in our case if the eye had been left in the orbit for some months longer. Sattler's patient was a boy of five, who came to Arlt's Clinique with proptosis, distended eyelids, restricted ocular movements, and chemosis. Evidently there was an orbital tumour present, and that it had implicated the optic nerve itself was to be inferred from the appearance of the papilla and retina, which was almost identical with what was seen in the case which forms the subject of this communication. The papilla and surrounding retina were occupied by a white tumour projecting into the vitreous, and covered with enlarged and tortuous arteries and veins. There were no hæmorrhages, there was no pain, the globe was becoming phthisical, and vision was lost.

Enucleation was followed by local suppuration. (Case occurred in 1876). Death occurred five months afterwards from tubercular meningitis.

Tubercular disease was found in the whole tract of the optic nerve and its sheaths from the chiasma to its intra-ocular expansion, besides infiltration of the orbital tissues, and miliary deposits in the retina. Sattler is inclined to regard the case as secondary to tubercular disease which was present in the bronchial glands.

Cirincione records a case of tuberculosis of the orbital part of the optic nerve (referred to in Michel's *Jahresbericht of Ophthalmology*, 1890, p. 181). The patient, aged eleven, died of basilar meningitis, and there was found in the orbit a tumour of 14 by 17 mm. springing from the nerve, and of tubercular nature. The origin was thought to be a caries of the upper orbital border which had existed for two years previously.

A tubercular nodule in the optic nerve is described by Demours (*Traite des Maladies des Yeux*, 1818, T. 1, p. 75); and Cruveilhier records the presence of a similar condition in a case of tubercular basilar meningitis (*Traité d'anat. path. gèn.* T. IV., p. 793). Demours' case died of chronic lung disease, and Cruveilhier's had deposits in apices of lung.

Tuberculosis of the optic nerve complicating tuberculosis of the uveal tract is described by Wagenman, v. Graefe's *Archives* XXXIV. 4, s. 178, and Bongartz', *Inaug. Dissertation* (referred to in Michel's *Jahresbericht of Ophthalmology*, 1891, p. 165). In this case the retina and optic papilla were the parts affected, and there was disease of the apex of left lung. In Wagenman's case, a girl of sixteen enucleated in Professor Leber's Clinique in 1874, the patient was alive and well fifteen years after the operation.

Our patient is reported by Dr. Powell to be in good health now, some 18 months after the operation, only somewhat thinner than she used to be, and with a pain in the back of her head, which gives her some trouble.

One of her uncles died of meningitis, and another maternal relative of consumption.

a. sclerotic. b. choroid, with pigmentary layer of retina attached.
c. colloid material between choroid and detached retina.
d. detached retina. e. tubercular mass in the retina with caseation and giant cells.

a. b. c. d. as before. b'. pigmentary layer of retina.
e. tubercular nodule. f. giant cell.