This article was downloaded by: [York University Libraries]
On: 21 November 2014, At: 13:34
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Transactions of the Botanical Society of Edinburgh

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tped18>

XIII. Notice of a new species of Antrophyum

R. K. Greville F.R.S.E. F.L.S. c.

Published online: 01 Dec 2010.

To cite this article: R. K. Greville F.R.S.E. F.L.S. c. (1850) XIII. Notice of a new species of Antrophyum, Transactions of the Botanical Society of Edinburgh, 3:1-4, 63-64, DOI: [10.1080/03746605009467573](https://doi.org/10.1080/03746605009467573)

To link to this article: <http://dx.doi.org/10.1080/03746605009467573>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

XIII. *Notice of a new species of Antrophyum.* By R. K. GREVILLE, LL.D., F.R.S.E., F.L.S., &c.

READ 11TH MAY 1848.

IN addition to the two new species of ferns (*Oleandra Sibbaldii* and *Grammitis blechnoides*) recently communicated to Professor Balfour by Dr. Sibbald from the island of Tahiti, I have now to submit the description of a third to the Botanical Society. The discovery of this plant is an additional proof how much remains to be done in an island where numerous collections have been made, but which is evidently still rich in undescribed productions. It is to be hoped that Dr. Sibbald will have an opportunity of revisiting Tahiti under more favourable circumstances, and that he will add largely to his collections, especially of ferns and mosses.

The interesting fern which forms the subject of this short notice belongs to *Antrophyum*, a genus having undivided, more or less lanceolate fronds, in which the sori form continuous grooved lines on the simply reticulate venation. *Antrophyum* is thus nearly allied to *Hemionitis*, from which, it must be confessed, it scarcely differs, except in the simplicity of the frond, for the grooved sorus is a somewhat variable character.

There is however another genus, *Polytænium* of Desvaux, which has been separated on, as it appears to me, more slender grounds. In that genus the sori are not reticulated, but form uninterrupted parallel lines, connected however by non-soriferous veins.

The plant I am about to describe seems to do away with such a distinction; for in it the sori are parallel, uninterrupted, and might be described as remotely forked rather than reticulated, and so seldom does *any* division in the sorus take place, that it is sometimes simply continuous for two or three inches together.

Antrophyum Grevillii (Balfour in herb.); fronde sessili, late lineari-lanceolata, inferne præcipue attenuata, soris approximatis, parallelis, longissimis, villosis, remote furcatis.

I cannot find any described species of *Antrophyum* which at all corresponds with Dr. Sibbald's specimens. The fronds are tufted, ten to eighteen inches in length, fully an inch broad in

the widest part, from whence they become insensibly narrower towards the base, which however never passes into a true stipes, although there is for the space of two or three inches an obscure midrib. The sori, which constitute the most remarkable feature, are so approximated as to be not more than a line apart, forming twelve or more uninterrupted lines, which sometimes divide at very remote intervals, but scarcely ever anastomose. The capsules, which are similar to those of the other species of the genus, are almost quite concealed by the mass of ferruginous hairs which arise along with themselves from the soriferous vein.

EXPLANATION OF PLATE V.

Fig. 1. Antrophyum Grevillii, nat. size.

- 2. A portion of the frond, showing the groove and soriferous vein.
- 3. A capsule with some of the ferruginous hairs.
- 4. Seeds.

Antrophyum Grevillei.

D. Greville del.

J. De C. Sowerby sc.