This article was downloaded by: [University of Glasgow]
On: 19 December 2014, At: 01:40
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Annals and Magazine of Natural History: Series 2

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/tnah08>

On two new species of South American birds

Philip Lutley Sclater

Published online: 10 Dec 2009.

To cite this article: Philip Lutley Sclater (1854) On two new species of South American birds, *Annals and Magazine of Natural History: Series 2*, 14:80, 158-158, DOI: [10.1080/037454809494323](https://doi.org/10.1080/037454809494323)

To link to this article: <http://dx.doi.org/10.1080/037454809494323>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

On two new species of South American Birds.

By PHILIP LUTLEY SCLATER.

1. *CULICIVORA BOLIVIANA*, Sclater. *C. supra plumbea; infra alba plumbeo paululum tincta; ventre niveo; fronte regione oculari et genis nigris; alis nigricantibus, tectricibus et secundariis latè, primariis strictissimè albo limbatis; caudà nigrà lateralibus rectricibus albo terminatis 4; extimis ferè omninò albis—rostro et pedibus nigris.*

Long. tota $4\frac{3}{4}$; alæ $2\frac{1}{8}$; caudæ $2\frac{1}{4}$.

Hab. Bolivia (Bridges), D'Orbigny.

The present bird is the fifth of this interesting genus, of which the best known are the *C. cærulea* (Linn.) of the United States, and the *C. dumicola* (Vieill.) of Brazil and Paraguay. The Prince of Canino notices two other previously unrecognized species in his 'Conspectus Avium,' p. 316. These I have never seen; but his fifth species, the *C. budytoides*, De la Fresnaye, of which I have examined the type, belongs, I believe, more properly to another genus. There are specimens of the present species in the British and Paris Museums, and in the fine collection of the Baron de la Fresnaye at La Fresnaye near Falaise.

2. *PIPPA FLAVO-TINCTA*, Sclater. *P. alba, flavo pallidè tincta; pileo alis caudâque cum dorso inferiore nigris; uropygio cum ventre viridescente-cinereis; rostro nigro; pedibus flavis.*

Long. tota $3\frac{1}{2}$; alæ $1\frac{7}{8}$; caudæ $1\frac{1}{8}$.

Hab. Sta Fe de Bogota.

This species is very like the common *Pippa manacus*, but is smaller, and has the white parts of its plumage tinged with yellow and much less black on the back. There are examples in the Museum of the Jardin des Plantes at Paris and of the Baron de la Fresnaye.—*Proc. Zool. Soc.* Feb. 24, 1852.

Description of a new species of Hyrax from Fernando Po.

By LOUIS FRASER, H.M. Consul at Whidah.

HYRAX DORSALIS, Fraser.

Adult male. General colour grizzled brown, becoming darker towards the back, where the hairs are annulated with black; a line of yellowish white, about four inches long, commencing over the short ribs and running hindways; muzzle naked and of a brown colour; eyes light hazel.

Length of head $4\frac{1}{2}$ in.; neck and body about 18 in.; hind feet, from heel to toe, 3 in.

Hab. Island of Fernando Po.

Its native name is 'Naybar'; it is nocturnal in its habits, and is no doubt common, as its loud cry of ccurr-ccurr-ccurr may be heard every evening after dark, during the commencement of the rains.

The Boobies say, it sleeps in the trees all day, and feeds upon leaves at night, but is very difficult to find.