different, presenting characters as distinct histologically as those which clinically differentiate the two diseases. In neither case ought there to be the slightest difficulty in diagnosing between the two.

In conclusion, I hope this paper with the microscopical illustrations so well executed by Mr. Max Colhoun may serve to make the very interesting condition known as lichen more intelligible to those who may read it.

Cavendish-square, W.

INFANTILE SCURVY.

By EDMUND CAUTLEY, M.D. CANTAB.,

PHYSICIAN TO THE BELGRAVE HOSPITAL FOR CHILDREN; ASSISTANT
PHYSICIAN TO THE METROPOLITAN HOSPITAL.

IN view of the discussion raised in THE LANCET on the cooking of milk and its effects on nutrition the following cases of infantile scurvy which have recently come under my notice may prove of interest. Cases of this nature are far more common in private than in hospital practice. This is no doubt due to the fact that the same care is not devoted by the mother or nurse to the preparation of the child's food, and that even when fed on patent foods the child often gets in addition "tastes" of unboiled milk and other articles of diet. The children of the lower classes are rarely kept on a diet of sterilised milk or even on patent foods only for any very long time and, as these cases show, the development of scurvy is a very slow process.

CASE 1. Scurvy due to a prolonged diet of sterilised cream and milk.—The patient was a female, aged 10 months, living in Surrey. She was brought to me at the age of nine weeks, weighing eight pounds, just as much as when she was born. She had been fed on various kinds of diet. On a diet of cream, milk, water, sugar, and lime water, in suitable quantities and at proper intervals, she gained 73 ounces in 12 weeks and had not a single bad symptom. The food was sterilised for 20 minutes. I did not see her again for five months, during which time she had been fed on exactly the same diet, the same amount and at the same intervals, and the food had been continuously sterilised for 20 minutes. She had now general tenderness of the lower limbs, disinclination to move them, pain on movement, and some swelling of the middle of the right thigh. Two teeth were cut and the surrounding gums were swollen and rather dusky. Four more teeth were almost through. There was no evidence of rickets. The milk mixture was given boiled instead of sterilised, and in addition the child was given barley-water, fruit juice, and a little meat juice. On account of a family susceptibility to tuberculosis, two sisters having died from tuberculous meningitis, and not knowing the source of the milk-supply I did not care to run the risk of putting the child on unboiled milk. Recovery was rapid and uneventful.

CASE 2. Scurvy due to patent foods.—The patient was a male, aged 11 months, living at the seaside. For the first three months he was fed on a patent food advertised as suitable for an infant at that age. During the next three months he was fed on another patent food, provided by the same firm, for children at that age. At six months an attempt was made to put him on the third variety of the same firm's preparations, but his digestion was upset, and he was then fed on a malted milk. His first tooth was cut at five months. For three months he had suffered from variable pains, anorexia, and wasting. He was not anæmic. There was a moderate degree of rickets. He had seven teeth. The gums were dusky and spongy. There was general tenderness and pain on movement. On a suitable diet he recovered rapidly.

CASE 3. Scurvy due to patent foods.—The patient was a female, aged 11 months, living in the country. Since the age of five months she had been fed on patent foods only and seemed to thrive well, getting very fat. For a month she had been fretful, disinclined to move, and would cry on handling. For three weeks she had been treated for rheumatism. She was very fat, her limbs resembling Bologna sausages. She had seven teeth. The gums were spongy, purplish, and bled freely. The legs were kept partially flexed and she screamed when attempts were made to straighten them. There were general tenderness, pain, and inability to sleep. There was no evidence of rickets. She soon recovered.

CASE 4. Scurvy due to sterilised milk.—The patient was a male, aged 15 months, living in a London suburb. He had been fed on milk from a "milk laboratory" since a few months of age. Recently he had been at the seaside and had only returned the day before I saw him. He had 10 teeth, with deep purple congestion of the gums, was very anæmic, sweated freely, cried when moved, and presented general tenderness of the limbs and slight beading. There was marked swelling of the right leg and to a less extent of the right thigh and left leg. This was much the worst case of the series, but the patient recovered very rapidly.

The treatment in each case consisted in a diet of unboiled milk (except in Case 1), fruit juice, meat juice, and barley water. In the fourth case vegetable purées were also given and mashed sieved potato. All these cases occurred in infants of the upper middle class in the social scale and they had all been more or less under medical supervision. Climate does not appear to have exerted any influence, though it is possible that the healthy hygienic surroundings and a copious supply of pure air may have postponed the onset of the disease. In all cases the diet consisted of a cooked food, but the disease did not develop for many months. The first case was the mildest, although the food had been continuously sterilised for eight months and it was not necessary to stop cooking the milk in order to cure the disease. The milk was boiled for a few minutes instead of being sterilised for 20 minutes.

It strikes me as a remarkable fact that during the last seven years I have in hospital practice brought up a very large number of infants on boiled milk, with the addition of barley water after the age of two or three months, and yet during the whole of that time no one of them has developed scurvy. Many have had the milk boiled until they have reached the age of one year. The rate of gain in weight has always been satisfactory when the directions have been properly carried out. Consequently I am astonished to read in your valuable journal that a physician of one of the largest London hospitals actually states that numbers of infants "are made peevish, fretful, and even seriously ill by the pernicious effects of boiled milk on their nutrition." In large towns it is absolutely imperative that milk for infants should be boiled, though it is not so necessary that it should be sterilised. In 1895 I examined milk bought in London shops and found that it contained from 3,000,000 to 30,000,000 of micro-organisms in each teaspoonful. Surely no one can urge that such milk is better uncooked than cooked. Under the present methods of milking and distribution of milk it is quite impossible to prevent the contamination of milk and the rapid growth of micro-organisms in such a very nutrient fluid. The only means we can safely devise to minimise the risk is to cook the milk. No chemical treatment of the fluid is proved to be less harmful. It is an established fact that many epidemics have been traced to the milk-supply. Such epidemics are hardly possible from this cause if the milk is boiled, for the common disease germs are readily destroyed by the process. Many maintain that infected milk is a great cause of infantile tuberculosis, and if this is the case it still more strengthens the position of those who insist on the boiling of milk for infants.

Let us consider for a moment the arguments against cooking the milk. In the first place, it is said that it diminishes the nutrient value. Such an argument might be brought forward against many articles of food and yet would not induce us to eat the food uncooked. The experiments of Vassilieff on healthy young men showed that the nitrogenous ingredients and the fats of boiled milk are less readily assimilated than when the milk is unboiled. Koplik's observations on infants showed that the amount of unabsorbed nitrogenous constituents is practically the same whether the milk is boiled, pasteurised, or sterilised, and is greater in each case than when the milk is unboiled. Bendix came to the conclusion that, in the case of infants two years of age, the nutritive value of sterilised and unsterilised milk is practically the same. Weber's experiments on calves showed that the nutritive value of uncooked milk was greater than that of cooked milk. Though there is great liability to error in experimental work which involves very elaborate chemical processes it may fairly be concluded that the nutritive value of milk is somewhat lessened by the process of cooking. Its antiscorbutic properties are also diminished and, in my opinion, the longer the subjection of milk to a high temperature the greater the diminution in these latter properties. For this reason the proprietary foods made from milk are more likely to lead to scurvy than is a diet of boiled milk.

There is no doubt that sterilised milk as a continuous diet will in time produce scurvy. It is equally clear that such a diet must be continued for many months—eight months in Case 1 and then only a very mild form of the disease; more than a year in Case 4 and hence a more serious attack. In both these instances nothing else was given. But are we to condemn the cooking of milk for such a reason as this? I can point to a very large number of infants who have been fed on boiled milk for many months and have never developed scurvy and have gained weight as rapidly as the most anxious mother could wish. The other two cases illustrate still more strongly the disadvantages of cooked foods, especially the prolonged cooking to which the proprietary foods are subjected in order to preserve them. Even in these instances the scurvy took many months to develop and a little modification of the diet would have prevented it entirely.

In conclusion, I most strongly maintain that the boiling of milk for infants is essential, save under most exceptional surroundings; that the longer milk is cooked and the higher the temperature to which it is submitted the less will be its nutritive and antiscorbutic value; that such deterioration in its properties is of no serious importance, if it is not forgotten, for it can be very easily remedied; that any healthy infant can be brought up on a diet of boiled cow's milk, suitably diluted and mixed and given in proper quantities at due intervals; that failure to bring up infants on such a diet is rarely, if ever, due to the fact that the milk is cooked, but is due to some error in the mode of feeding or the amount given; that the very slight tendency to scurvy can be easily remedied by the addition of barley-water and a little fruit juice; and, finally, that healthy boys at a public school can drink uncooked milk with impunity as long as it is not contaminated with disease germs, but that to base on such a fact an argument that infants should be fed on unboiled milk is merely a delightful instance of illogical reasoning.

Upper Brook-street, W.

ON X RAYS IN THE TREATMENT OF LUPUS AND RODENT ULCER.

By JAMES STARTIN, M.R.C.S. ENG.,
SENIOR SURGEON TO THE LONDON SKIN HOSPITAL.

IN September, 1899, I found the beneficial effects of the x rays in lupus, rodent ulcer, and certain other affections of the skin. I have been working with an instrument provided by Mr. Isenthal, consisting of a 15-inch sectional coil worked at 16 volts with a mercuric interrupter. The average speed of interruptions is about 250 a minute, the average current about from three to five ampères. I used at first a five-inch tube, then a smaller one made by Cox. I place the tube generally at a distance of five inches between the anti-cathode and the part to be exposed, and the duration of the exposure lasts from 10 to 15 minutes, giving four or five exposures on consecutive days. In cases of lupus the skin becomes slightly red after the first four exposures, the nodules bright and shining, and the pus dries up, with no doubt destruction of the bacteria. After a few days scabs form, then drop off, leaving a red glazed surface, which may or may not scab over again. This process goes on until no more scabs are formed and the surface is smooth and normal. The surfaces acted upon become slightly cedematous and a healthy epidermis begins to form. A good deal of reaction takes place in the diseased areas, such as we used to see on the administration of tuberculin a few years ago, but without any detrimental constitutional effects. This is followed by a copious discharge of serous matter which ceases in a few days.

Whether the x rays operate injuriously on the micro-organisms or whether phagocytosis becomes more pronounced under the slight stimulus has not yet been definitely made out. What, then, is this curative action so destructive to bacteria and what is the agent that produces it? In what manner do the ethereal vibrations of the x rays correspond to the ultra-violet, photo-chemical rays of the solar spectrum?

As far as my experience leads me at present the most suitable cases for treatment by the x rays are lupus, both

true and erythematous, rodent ulcer, epithelioma, psoriasis, localised eczema, lichen, alopecia areata parasitica, and other parasitic diseases of the ringworm class. To these may be added superficial cancerous affections and dermal tuberculosis. When we can get the penetrative power to extend still further, if it be true that cultivated bacteria can be destroyed in from 15 to 20 minutes, the x rays show a future use in antiseptic surgery.

CASE 1.—The first case I wish to record is that of a man, aged about 60 years, who came under my notice in September, 1898, with a large rodent ulcer extending over the whole frontal region down to the supra-orbital and nasal region, with deep ulcerations, some extending to the periosteum, and having the distinct crateriform edge characteristic of rodent ulcer. I used various treatments by caustics, ethylate of sodium, chromic acid, salicylic collodion, powders and ointments with varied success. Nevertheless, the disease steadily advanced, so that in June, 1900, I decided to put him under the x-rays treatment and gave him his first exposure on the 18th of that month, repeating the exposure for seven consecutive days, for 15 minutes at six inches from the skin. After the first four exposures much disturbance of tissue was noticed. The skin became red and glazed, the edges red and inflamed, and a copious discharge took place. After a day or so all discharge ceased and the sore began to take on a more healthy appearance, granulation tissue beginning to form. He went home with a simple iodoform dressing and I saw him again in July. The appearance of the wound was more healthy, but still there were two large excavated ulcers of about the size of a two-shilling piece in the centre of the area. I decided to give him a few more exposures, seven, and again in October, with the result that the raised edges of the wound became flattened, granulation tissue and an apparently healthy skin forming on most of the old sites of disease.

CASE 2.—The patient was a strong man who came under my notice three years ago with a deep foul ulcer on the forehead, nose, and face near the inner angles of both eyes, covered with thick crusts and very tender and sore, caused by a wound which he had received from the branch of a tree. I put him under various treatments by caustics, ethylate of sodium and salicylic and iodoform dressings, as in the other case, with varied results, and on Oct. 22nd last I sent him into the hospital in Fitzroy-square and gave him 10 exposures to the x rays with very beneficial results. The ulcers on the forehead and nose have nearly healed and only the ulceration in the corner of the nose and right eye remains. It is difficult to direct the rays there on account of the near proximity of the eye. The sight is unaffected at present.

In both these cases there was some considerable dermatitis, which was to be expected, and we need not be afraid of it as it promotes that disintegrative resolution in the tissues which must be present in the formation of a healthy cicatrisation. The surrounding skin has also taken on a more healthy appearance. The scars that remain are more natural and resemble the surrounding skin.

CASE 3.—A girl, aged 17 years, came under my care at the hospital in May, 1890. It was one of the worst cases of true lupus that I have seen for some years, with perforation of the bridge of the nose. I commenced treating her with tonics, quinine, and arsenic and using an antiseptic dressing of salicylic acid, iodoform, and vaseline, and on June 20th I gave her seven exposures with much the same result as in the two previous cases. There was considerable disturbance in the ulcers with much discharge for a few days. Then there followed a complete drying up of the sores and the assumption of a much more healthy appearance.

CASE 4.—A young married woman, aged about 30 years, had well-marked butterfly lupus erythematosus on both cheeks and on the bridge of the nose. The patches, which were crimson red and inflamed, were of about the size of a two-shilling piece. This patient came under my care in October, 1897. I put her under treatment of various applications, salicylic collodion, &c., with some good results, but the case proved to be tedious, so I determined to put her under the x rays at intervals of three days for a few weeks on Nov. 5th last. She has now had six applications with capital results. Cicatrisation has taken place.

CASE 5.—This patient, a young woman, aged about 21 years, came under my care at the hospital in March, 1900, with psoriasis of the legs, the arms, and the body. After treatment with tonics, arsenic, and tarry preparations she got fairly well excepting one obstinate patch about six inches