surgery. The history was as follows. On Jan. 15th his mother had applied to me for medicine for the boy, who had shown symptoms of worms. The powders (santonin and calomel) and medicine given had apparently been followed by good effect, for on the 27th he was running about at play and appeared all right until 7 A.M. on the 28th. At this time he was suddenly seized with abdominal pain and great tenderness, and he was brought to my surgery, as stated, at 9 A.M., his death occurring half an hour later.

At the post-mortem examination it was discovered that the cæcum, the appendix, and part of the ileum lay behind the stomach, having passed through the foramen of Winslow. Reduction was impossible until the gastro-hepatic omentum containing the vessels, &c., passing to the liver was divided, when the mass was easily withdrawn. A curious part of the case was that the cæcum and appendix up to the tip of the latter were tightly packed with round worms, the appendix being about the diameter of the thumb and conical in shape. 53 large worms were removed from the cæcum and appendix. The strangulated portion was naturally deeply congested and swollen, but the rest of the small intestine was not dilated, although there was some injection of the mesenteric vessels generally.

Interesting points about this case are: (1) the age of the patient; (2) the accelerated collapse; and (3) the fact that the condition was apparently directly brought about by the presence of the worms.

Knysna, Cape Colony.

A CASE OF ACROMEGALY.

BY E. P. CARMODY, L.R.C.P. LOND., M.R.C.S. ENG.

THE patient, a married woman, aged 33 years, was seen by me first on Dec. 19th, 1908. She complained of constant headaches, most marked over the forehead, and general weakness, and stated that she had been in poor health for about seven years. She had noticed enlargement of her hands and face for the last two years and had become round-shouldered. Her voice had been getting deeper and harsher. Catamenia had ceased six years ago. There were profuse sweatings at night. There was a history of rheumatic fever. Her condition was as follows. She was intelligent. She was a sallow, unhealthy-looking woman. There was enlargement of the face, especially the nose, lower jaw, and lips. The frontal region was prominent. The skin and deeper tissues all over the body were thickened but could be pinched up without difficulty. The nasal and tarsal cartilages were increased in size. The ears were leathery. The lower jaw was large, but the lower teeth met the upper teeth when the mouth was shut. The teeth were good and regular. The lips were thick and pendulous. The tongue was slightly enlarged. There was marked kyphosis in the cervico-dorsal region. The clavicles and ribs were large and unduly prominent. The lungs and heart were normal. There was marked impairment to resonance under the manubrium sterni. The thyroid gland was not enlarged, but the patient had previously been under treatment for goitre. The skin was abnormally active. The hands were increased in breadth and the fingers much thickened; the natural creases of the hand were well marked. The feet were unaffected. The larynx was large. The vision was indistinct for near objects. There was slight dyspnoea. The urine had been examined on several occasions and contained no albumin nor sugar. Mentally the patient was usually melancholy. Although intelligent at times she was not clear as to her surroundings. Her memory for recent events was bad. She suffered intermittently with insomnia.

I have treated the patient with phenazone for the headaches and have given two-grain tabloids of pituitary extract three times a day for six weeks. Up to the present there is no improvement.

Tilehurst, Berks.

PROVIDENT NURSING.—A special meeting of the council of the Charity Organisation Society will be held on June 28th, at 4.30 P.M., at Denison House, Vauxhall Bridge-road, S.W., when Miss Amy Hughes of the Queen Victoria Jubilee Nurses Institute will read a paper on Provident Nursing. Sir Alfred Lyall, G.C.I.E., K.C.B., will take the chair, and the Right Rev. the Bishop of Islington will speak.

Medical Societies.

ROYAL SOCIETY OF MEDICINE.

MEDICAL SECTION.

Treatment of Typhoid Fever with an Anti-endotoxic Serum.

A MEETING of this section was held on May 25th, Dr. NORMAN MOORE being in the chair.

Professor R. T. HEWLETT read a paper on the Treatment of Typhoid Fever with an Anti-endotoxic Serum. He said that it was generally recognised that an anti-typhoid serum prepared by the injection of the horse with cultures of the typhoid bacillus was useless in the treatment of typhoid fever. An anti-endotoxic serum had been made by the injection of the horse with typhoid cell-juice. Such a serum had an active neutralising power against both the typhoid endotoxin and the typhoid bacillus. The results of the use of this serum in nine cases of typhoid fever were detailed. The impression which had been gained so far was that several cases were benefited and that in two at least the disease was cut short by its use. Typhoid fever was, however, such a variable disease that it was extremely difficult to estimate the value of any form of treatment unless a large number of cases was available. In no instance did the serum appear to do any harm. Unfortunately, cases of typhoid fever usually only came under observation when the disease had existed for a week or longer. If the disease had existed for some time no immediate effect could be hoped for from the use of a serum, however efficient that might be. Moreover, the existence of secondary infections, which often occurred in typhoid fever, must limit the value of any serum. As regards dosage of the serum, his impression gained from those few cases was that the serum should be given freely—namely, from 10 to 20 cubic centimetres daily—until the temperature had reached the normal, and perhaps in smaller doses for a week subsequently. Another point was that the preparation of an anti-endotoxic serum was at present in the experimental stage. They did not know for how long the horse should be immunised and he was inclined to think that it was possible to over-immunise, that was the serum did not increase in potency after a certain number of doses of the typhoid endotoxin; in fact, harm might result from continuing the injection of the horse. The horse seemed to be extremely susceptible to the typhoid endotoxin and a period of eight months elapsed before a dose of 30 cubic centimetres could be given. In a second horse which he was immunising the commencing dose of toxin was 0.1 cubic centimetre, and even that small dose caused so much disturbance that it had to be repeated five times at weekly intervals before the dose could be increased. He thought it would be of interest to test whether the typhoid endotoxin would give an ophthalmic reaction. For that purpose some typhoid endotoxin was carefully prepared, so as to eliminate as far as possible extraneous substances, and dried *in vacuo*. The substance gave a marked reaction in several cases of typhoid fever. It also gave a reaction in one or two cases of other diseases, but did not give a reaction with several cases not typhoid. The interest of the observation was that the substance producing the reaction was thus shown to be an intracellular constituent of the organism.

Dr. E. W. GOODALL, of the Eastern Hospital, Homerton, read a paper detailing the results of the treatment of 26 cases of typhoid fever with the same anti-endotoxic serum. In regard to the value of the treatment he pointed out one fact which, before he treated a single patient with serum, led him not to be too sanguine as to its results. He referred to the advanced period of the disease at which the patients were in most instances sent to hospital. That was a fact which militated against any form of treatment, and was, he believed, one of the chief reasons why in his hands the bath treatment, for instance, had not achieved the excellent results obtained by other physicians. Enteric fever was a disease which varied in severity to a considerable degree within short periods of time, and the number of cases he had recorded was very small, so that it was impossible for him to speak with certainty as to the efficacy of the treatment. He could only say that it appeared to have done good