Beobachtung der Sonnenfinsterniss 1884 März 26

auf der Sternwarte in Helsingfors.

	Eintritt des Mondes	Austritt	Beobachter	Fernrohr	Vergr.
M. Z. Hels.	18 ^h 59 ^m 5 ^s	19 ^h 42 ^m 13 ^s	Donner	Helimeter von 71 ^{mm}	200
»	18 58 48	19 42 31	Mag. G. Dreijer	Fernrohr von 57 ^{mm}	76
»	—	19 42 8	Stud. K. Eklund	Refractor von 174 ^{mm}	145

Die Luft war während der Beobachtung sehr gut und ruhig, der Himmel ganz ohne Wolken.

Helsingfors 1883 März 27.

Anders Donner.

Observations of Comet Pons-Brooks

made at the Halsted Observatory, Princeton N.J., with the 23inch Equatorial.

Communicated by Professor C. A. Young.

The observations of position were made by Professor Young, except those of Dec. 12th which were by Mr. Mc. Neill. The filar-micrometer was used, with bright wires and a power of about 160. On Jan. 16th and 17th however the observations were made by twilight, without artificial illumination, and consisted in measurements of distance, and position-angle, the position-zero being determined after the observations. In all other cases differences of α and δ

were measured. In reducing the star-places to 1883.0 the tables of m and n employed were those given in Oppolzer's »Lehrbuch der Bahnbestimmungen« (vol. I, 2^d Ed.), which are based on Struve's Constants. The comparison-stars on Jan. 16th and 17th were very small, not much above the 10th magnitude, and have not been identified.

The reductions and computations have all been made by Mr. Mc. Neill.

1883-84	M. T. Gr.	$\Delta\alpha$	$\Delta\delta$	Comp.	α app.	$\log p.\Delta$	δ app.	$\log p.\Delta$	Red. ad l. app.	*
Oct. 4	13 ^h 9 ^m 51 ^s	—	+ 2' 11".2	—8	—	—	+58° 7' 43".4	9.971	—0.64 +16".8	1
4	13 27 25	+ 1.517	—	8.—	16 ^h 29 ^m 12.14	9.910	—	—	—	1
5	13 5 22	—	— 6 10.2	—8	—	—	+57 54 52.8	9.971	—0.67 +16.2	2
5	13 23 32	+23.95	—	8.—	16 29 50.04	9.907	—	—	—	2
9	13 13 48	—	— 6 48.1	—12	—	—	+57 4 1.2	0.181	—0.67 +16.2	3
9	13 27 55	+ 2.09	—	12.—	16 32 53.13	9.905	—	—	—	3
Nov. 1	13 27 58	—	+ 5 13.3	—25	—	—	+52 33 38.2	0.567	—0.55 +15.4	4
1	13 29 31	—11.33	—	28.—	17 5 5.07	9.867	—	—	—	4
5	13 3 10	—	— 6 17.3	—12	—	—	+51 50 7.1	0.517	—0.51 +15.6	5
5	13 27 7	+15.41	—	13.—	17 13 28.24	9.859	—	—	—	5
17	13 7 53	+ 8.51	—	8.—	—	9.839	—	—	—0.29 +17.0	6
17	13 22 21	—	— 4 53.7	—8	—	—	—	0.628	—	6
28	10 34 17	—	— 2 21.1	—8	—	—	+47 15 48.7	0.026	+0.04 +19.7	7
28	10 55 0	—43.52	—	8.—	18 23 6.26	9.760	—	—	—	7
29	10 36 18	—	+ 2 24.8	—8	—	—	—	0.052	+0.08 +19.8	8
29	10 54 32	+59.02	—	10.—	—	9.757	—	—	—	8
Dec. 12	10 57 57	—	+ 4 1.0	—9	—	—	+42 13 49.4	0.261	+0.81 +24.6	9
12	11 16 53	+36.35	—	10.—	19 31 29.47	9.731	—	—	—	9
Jan. 16	10 39 20	— 2.41	— 1 28.3	7.7	—	9.348	—	0.794	—	10
17	10 43 17	—18.52	— 0 47.8	6.6	—	9.360	—	0.799	—	11

Mean Places of Comparison Stars.

*	α 1883.0	δ 1883.0	Authority	*	α 1883.0	δ 1883.0	Authority
1	16 ^h 29 ^m 11.61	+58° 5' 15".4	AOe. 16310	7	18 ^h 23 ^m 49.74	+47° 17' 50".1	AOe. 18273
2	16 29 26.76	+58 0 46.8	Krueger Z. A. N. 2553	8	18 26 13	+46 57	DM. 46° 2497
3	16 32 51.71	+57 10 33.1	Krueger Z. A. N. 2548	9	19 30 52.31	+42 9 23.8	Wash. Obs. 1875 N. 646
4	17 5 16.95	+52 28 9.5	AOe. 16858	10	—	—	Star not identified
5	17 13 13.34	+51 56 8.8	Lal. 31560	11	—	—	Star not identified
6	17 44 32	+49 42	DM. 49° 2692				

Star 2 is double, Mag. 8 and 9.5, Position angle 219° 35', Distance 1".55.

The comet was observed on several other nights when no stars good for comparison were near enough, the field with the lowest power being only 15' in diameter. For the last two months the weather has been very bad, there having been no nights when the seeing was really good, and only a few when anything at all could be seen.

A few notes on the physical observations are appended.

1883 Sept. 5 14^h Greenw. M. T. Comet first found. Very faint, with slight central condensation. Diameter 1½' or 2', but having no very definite outline and no tail.

Sept. 16. Diameter of nebosity 2' 46". A faint streak visible in nebosity at position angle 61°, probably the beginning of a tail.

Oct. 1. Tail about 8' long: position angle 57°. Nucleus almost stellar.

Oct. 2. Nucleus throws off small brush towards the sun. Tail brighter on southern edge: position angle 66°.

A small direct-vision spectroscope, without slit, shows two knots in the spectrum of the nucleus. Comet too faint for spectroscope with slit.

Oct. 3. Tail about 5' long, position angle 59°. Diameter of comet's head measures 57". Faint brush toward sun. Too faint for spectroscope; sky hazy.

Oct. 5. Comet fainter; tail hardly visible.

Oct. 24. Comet round; no tail. Nucleus stellar, like a star 9½ magnitude.

Nov. 1. Comet found in strong twilight. Nucleus 8½ mag.

Nov. 5. A faint brush, like an incipient tail, at 50° position-angle: very faint and indistinct.

Nov. 7. Tail unmistakable, but faint: position-angle 50°. Nucleus much condensed, but hardly stellar; total brightness about 7½ magnitude.

Nov. 29. At 11^h 5^m Greenw. M. T. Comet passed within 2" or 3" of star of 11th magnitude, without any visible effect upon star's appearance. Comet flattened out

and better defined on side next sun, the major axis of the nebosity being at right angles to the sun's direction and about twice the minor axis; tail straight and pointing from sun; the whole form like this: T

Dec. 28. Spectrum of Comet observed with Grubb spectroscope and two prisms. Spectrum of the ordinary type: the nucleus-spectrum rather more nearly continuous than usual with comets.

1884 Jan. 3. Used Clark spectroscope with one prism. Comet spectrum very fine. Three bands and a faint knot in the purple. The principal band in the green rather brighter in proportion to the others than ordinary, and broken into at least three, perhaps four or five, lines or narrow bands. Spectrum of the nucleus nearly continuous, but widened at the bands. No dark lines visible. With lowest eyepiece (power about 120) the appearance of the comet is very striking. The nucleus is about 6½ magnitude, reddish and nearly stellar. There are 4 or 5 brushes directed toward the sun, each 20" or 30" long. The tail near the head is cut out a little on the north side.

With field glass the tail is 4° long and nearly straight. Moon 4 days old, not far away.

Jan. 4. Comet's head shows three curved jets.

Jan. 16. Comet found in twilight, nucleus stellar, but with several brushes directed to sun. No tail visible in the 23 inch; but as seen in field-glass, it was conspicuous, and 4° long. Spectrum of nucleus nearly continuous: only very slightly knotted where it crosses the hydrocarbon bands.

Jan. 17. With low power the comet shows a bright and pretty well defined sector on east side of nucleus, nearly at right angles to the general trend of the tail. Tail now easily seen in large telescope, being much better defined on southern than on northern edge. In field glass the tail is 6° long. Comet estimated as equal to star of 2½ magnitude.

Stormy weather prevented any further observations.

Princeton N. J. 1884 February.

Beobachtungen des Cometen Pons-Brooks

angestellt von *B. von Engelhardt* am Fadenmicrometer des Aequatoreals (von 12" engl. Oeffnung) seiner Privatsternwarte zu Dresden.

(Fortsetzung von Nr. 2547 und Nr. 2560 der Astr. Nachr.)

1883-84	M. Z. Dresd.	$\Delta\alpha$	$\Delta\delta$	Vergl.	α app.	$\log p.A$	δ app.	$\log p.A$	*
Nov. 9	7 ^h 16 ^m 23 ^s	+6 ^m 14 ^s 65	+ 7' 49" 0	14.5	17 ^h 22 ^m 53 ^s 62	9.762	+51° 9' 37" 8	0.522	1
19	10 19 2	+1 31.33	— 2 27.0	3.1	17 50 25.30	9.666	+49 15 56.1	0.841	2
24	6 33 40	—1 35.70	— 0 45.3	24.8	18 6 55.02	9.720	+48 15 18.2	0.497	3
29	7 10 22	—4 27.43	+ 1 28.0	18.6	18 26 24.86	9.729	+47 3 4.4	0.592	4
Dec. 29	6 59 29	+0 40.19	+ 2 27.0	30.10	21 23 12.99	9.567	+27 8 55.0	0.690	5
Jan. 1	6 50 45	+0 11.65	— 1 32.0	35.13	21 44 50.22	9.530	+22 43 33.8	0.710	6
15	7 1 0	+1 24.25	—13 40.9	24.8	23 17 31.21	9.458	— 2 41 13.0	0.846	7
21	6 36 51	—0 58.11	+ 3 1.8	2.1	23 49 5.05	9.411	—13 4 53.8	0.881	8
25	6 26 32	+0 33.05	+ 3 25.4	18.6	0 7 4.05	9.397	—19 7 25.8	0.897	9
25	6 40 1	+0 35.41	+ 2 37.0	9.3	0 7 6.41	9.426	—19 8 14.2	0.893	9
25	6 49 29	+0 37.31	+ 2 4.0	9.3	0 7 8.31	9.444	—19 8 47.2	0.890	9