the injection through the second puncture into the skin and subcutaneous tissues well over the intercostal spaces above and below. The infiltration is easy to carry out and only takes a few minutes; the patient feels only the one prick of the needle. The operation can be started immediately it is finished. It is necessary to emphasise the importance of using the rib shears with the utmost care, as any wrenching will give rise to pain at the costo-sternal junction.

On opening the pleural cavity there may be some discomfort, especially should the patient have a fit of coughing, which is sometimes the case in large empyemas.

In nearly all of the 85 cases operated upon since May the patients were suffering not only from empyema but had signs of active pneumonia in both lungs, and many had a blood pressure of well below 100 mm. Hg. Notwithstanding their serious condition no patients were lost at the operation nor did their behaviour during it give rise to further anxiety.

We claim that local anaesthesia for these patients gives greater safety and increases the chance of ultimate recovery.

INFLUENZA AND BLACKWATER FEVER.

BY SAMUEL SAMUEL, M.B., CH.B. LEEDS, M.R.C.S.,
LATE CAPTAIN, R.A.M.C.; HONORARY MEDICAL AND SURGICAL
OFFICER, LEEDS JEWISH HOSPITAL.

MUCH has already been written in connexion with the present influenza epidemic, both in regard to its etiology and its complications. In the lay mind especially the present influenza has been looked upon by some individuals as a type of malaria, probably spread by the returning malarial soldier, and this, amongst other reasons, accounts, to a certain extent, for the great consumption of quinine, both as a prophylactic and as a curative drug.

The present influenza has been of a most virulent type, and the commonest of complications have been the various hæmorrhages which have included the various rashes, bleeding from certain organs—e.g., nose, uterus, stomach, lungs, &c. Though cases have occurred frequently where blood has appeared in the urine, it has chiefly been present when an acute nephritis has been a complication. I doubt, however, if many have experienced a case similar to the following, where the urine has appeared as blackwater, similar to what some of us who have seen active service in Macedonia have witnessed.

The patient, a male aged 45, previously quite well in health and never domiciled abroad, began on Monday, Nov. 25th, with typical influenza. He was treated with simple remedies such as spir. æther. nit. and simple alkalies, but no quinine was administered. His temperature began to subside on the third day, i.e., on the Wednesday. This same day he began to complain of violent colic in the epigastrium, which on the following day moved also to the flanks. Warm applications, simple warm-water enemata, and even small doses of morphia, had no effect in quieting the pain. On the Friday evening, whilst the pain was still severe, the patient passed quite an abnormally large quantity of urine as black as ink, with a reddish tinge showing just at the edge where the urine came in contact with the side of the chamber. An examination of the urine revealed the same features as are present in typical blackwater fever. On the Saturday morning the patient was seen in consultation with Lieutenant-Colonel G. W. Watson, who ordered tr. belladonnæ in 5 min. doses every three hours for the colic and warm applications to the flanks. The colic began to disappear almost with the second dose, and in the evening the patient was free from pain, though still passing the black urine. On the Sunday the black urine began to clear and to diminish in amount. The patient's conjunctivæ and skin then assumed a marked yellow tinge, which, however, disappeared gradually in the course of the next few days with the nightly dose of calomel gr. 1 and a morning seidlitz powder. Albumin and granular casts still persisted for about two days after the blackwater disappeared, when these also disappeared.

The patient to-day (Dec. 9th) is free from all symptoms, though weak and anæmic and earthy-looking. He is now allowed up for one hour a day. Just a week has now passed and there has so far been no recurrence of the blackwater, and the urine is quite clear and normal in all respects.

Leeds.

ROYAL SANITARY INSTITUTE.—The Henry Saxon Snell prize of 50 guineas for an essay containing the best suggestions for Improvements in Apparatus and Appliances for Dealing with House Refuse has been awarded, along with the bronze medal of the institute, to James Jackson, superintendent of the refuse disposal department at Birmingham.

Medical Societies.

ROYAL SOCIETY OF MEDICINE.

SECTION OF EPIDEMIOLOGY AND STATE MEDICINE.

A MEETING of this section was held on Dec. 13th, when Dr. E. W. GOODALL, Lt.-Col., R.A.M.C., delivered his Presidential Address on

Enteric Fever in Flanders in 1914-1915.

He said that enteric fever was prevalent in Belgium at or about the time war broke out and cases occurred in England during the autumn of 1914 among Belgian refugees. The rapid advance of the Germans and the flight of the inhabitants were likely to assist the spread of the disease, the more so if the nature of the invaded country be considered. Belgium, west of a line drawn through Antwerp and Brussels, is a plain, intersected with rivers, mostly canalised, the water in which can flow but very slowly. This plain is, in places, thickly populated and highly cultivated, and human excreta are largely used for the purpose of fertilisation. The water is usually derived from shallow wells. Further, the Belgians, as a defensive measure, flooded the country to the north-east of Dixmude, the effect of which was to accentuate the stagnation. When trench warfare began, in November, 1914, the trenches were consequently in a very insanitary condition. At this time, of the four armies engaged in Flanders, the British alone was protected adequately by antityphoid inoculation. He gave evidence to show that by the end of November, 1914, there was in the Flanders area a considerable number of cases of enteric in the Belgian, French, and German armies, and probably amongst Belgian civilians, and a comparatively small number in the British Army. The outbreak began to assume epidemic proportions soon after the armies settled down to trench warfare.

An account was then given of the Friends' Ambulance Unit. Nearly all the members belonged to the Society of Friends, many being professional men, so that whatever line of work was taken up there was no difficulty in finding, amongst themselves, expert organisers and advisers. Part of a large building just outside Ypres was given to them, and here the local headquarters of the unit and the hospital were set up. Besides the treatment of patients in hospital the medical officers of the unit carried on an extensive outdoor visiting practice amongst the remaining inhabitants of Ypres and the surrounding villages not occupied by the enemy, and the hospital had been recognised by the Ypres Committee of Safety as the Ypres Civil Hospital. The authorities of the British Army enlisted the services of this unit as a friendly link between themselves and the Belgian civil authorities to help them in their campaign against the epidemic. The area, not occupied by the enemy, affected by the epidemic was about 31 miles long and 12 miles broad, stretching from Nieuport on the north to Ploegsteert on the south. The epidemic probably spread rapidly towards the end of December, 1914.

With regard to the number of cases information was limited, but it was probably very considerable; there must have been about 4000 cases under treatment daily during January and February, 1915, in the French military hospitals in Dunkirk and its neighbourhood, and the number in the Belgian Army must also have been considerable. One thousand cases (Belgian civilians) passed through the hospitals of the unit at Ypres and Poperinghe, and about 1000 more came to the knowledge of the officers of the unit, which were treated elsewhere. The British Army furnished comparatively few cases—only 827 up to May 22nd, 1915.

Character of the Cases.

The disease differed in several points from the types the speaker had seen in England during the previous 25 years.

1. The fever ran a very irregular course, the oscillations being frequent and extreme, so that the chart at the height of the disease resembled in many cases that of pyæmia or phthisis. Temperatures of 104° to 105° F. were by no means uncommon.
2. A considerable number of the cases exhibited pronounced nervous symptoms, coma or semi-coma, or restlessness and delirium. These symptoms did not pass off as the general condition improved and the temperature fell. The comatose cases did worse than the others; in such cases death occurred without recovery of consciousness,