they depend on the unbending of the uterus. All that can be said is that they are curiously adapted to produce the effect which it is urged is so unnecessary. Nothing will convince me, at all events, that the straightening of a severely flexed uterus is a useless procedure, for the reason that I have found this procedure, accomplished in a variety of ways in different cases, so invariably productive of relief to the patient, and so remarkably efficient in removing the congestion which so frequently accompanies the flexion in painful troublesome cases. I am not unaware of the occasional difficulty in determining the precise therapeutic value of a particular procedure, and allowances must of course be made and sources of error in this respect guarded against. Dr. Herman thinks I have wrongly attributed benefit derived by my patients to treatment of the flexion of the uterus present in such cases, whereas that benefit was due to the general treatment pursued. I am not likely, as will be gathered from my writings, to undervalue the benefit of careful general treatment. It is just possible that I, having seen and watched these cases, may be a competent judge on this point. Cases are not very rare in which there has been a long trial of general treatment, most carefully adapted to the patient, with entire failure in procuring relief, while an immediate improvement has followed the adoption of local treatment. There is much to be learnt from the endeavour to relieve the suffering attendant on long-standing severe flexion of the uterus. One of the things I have learnt from some experience of such cases is that these cases cannot be cured without a judicious combination of general and local treatment, as I have fully stated on former occasions. But the real truth as to the flexion factor and its importance becomes evident in those cases where good general treatment doing its best for the patient, the difficulty in restoring the uterus to its normal state is unusually great. Then it is seen that the improvement is measured by the success of the local treatment, and that in proportion as the uterus is restored to its proper physical condition, the sufferings of the patient diminish.

Berkeley-square, W.

CASE OF

LARYNGEAL AND PULMONARY TUBERCULOSIS, ORIGINATING IN SIMPLE CHRONIC LARYNGITIS.

By G. HUNTER MACKENZIE, M.D.,

SURGEON FOR THROAT DISEASES TO THE EYE, EAR, AND THROAT INFIRMARY OF EDINBURGH.

THIS case is thought worthy of record for the following reasons: 1. The priority in origin of the laryngeal as compared with the pulmonary affection. 2. The engrafting of a tubercular on an initial simple inflammation of the larynx. 3. The presence of the tubercle bacillus in the sputum and laryngeal secretion, and its aid in diagnosis and prognosis. 4. Certain toxic effects following the use of iodoform.

The patient, a male aged fifty-five, was first seen by me on March 28th, 1882. He complained of loss of voice, great pain in the ears, especially the left ear, inability to swallow properly, and cough. His throat affection had lasted for about seven years, and was supposed to have been due to a cold from sleeping under an open window. Hoarseness came on at that time, from which he had never been able to free himself, and which finally merged into aphonia. Family history was good. The man's previous history was uneventful, with the exception of a slight cardiac affection of rheumatic origin some twenty years previously. He was tolerably healthy in appearance, of the rheumatic diathesis, with no indication of constitutional disturbance. On laryngoscopic examination, the arytenoid cartilages were swollen (not the typical pyriform swelling of phthisis), the ventricular bands and general mucous membrane of the larynx were irregularly thickened, and the vocal cords could not be seen. Some ashy-white secretion coated the central third of each ventricular band. Faucial sensitiveness was unexaggerated. The pulmonary symptoms were prolonged expiration, a slightly impaired percussion note, and a dry bronchitic click at the end of inspiration at the right apex anteriorly. The sputum was purulent, with a slight admixture of mucus, and contained the bacilli of tubercle in fair abundance in every field of the microscope. The patient was treated by local applications of chloride of zinc and subsequently of

nitrate of silver to the larynx, with at first considerable benefit to the laryngeal appearances and symptoms and to his general condition, for from being confined to the house he was soon able to go out of doors. The main indications for treatment seemed to be to endeavour to subdue the local affection by metallic astringents, but a return of pain in the throat, and especially in the ear, compelled an attempt to alleviate the local conditions by sedative sprays, inhalations, and insufflations. Sprays and inhalations proving comparatively ineffectual, recourse was had to an insufflation of morphia, at first alone, but subsequently combined with iodoform; these appeared to produce more benefit than any other local application. The course of the case tended gradually downwards. The laryngeal appearances and symptoms approximated more and more towards the tubercular, and signs of excavation at the right apex were detected towards the end of August. In January crepitant râles were abundantly present down the anterior surface of the right lung, with cavernous breathing at its apex. The pain in the throat and ear was occasionally very severe, especially on swallowing. The patient died on the 15th of February, 1884, of exhaustion from inanition.

A post-mortem examination of the throat only was permitted. There was considerable destruction of the tissues within the larynx. The arytenoid cartilages had undergone necrosis and dislocation, the mucous membrane in their vicinity was swollen and infiltrated, and the ventricular bands had disappeared from ulceration. The vocal cords were intact, except at their arytenoid attachments, where a loss of substance to the extent of one-sixth had taken place. The general mucous lining of the larynx was irregularly thickened, and at the posterior parts of the ventricles on both sides were deep excavating ulcers, leading to the necrosed arytenoid cartilages. Bacilli of tubercle were found in the following situations: (1) In a scraping of mucus from the lower part of thyroid and upper part of the cricoid cartilages. Here they were few, with an aggregation of about twenty in one field. There was no subjacent ulceration. (2) Underneath the anterior commissure of the vocal cords they were about as numerous as in the first situation. (3) In the secretion filling the laryngeal ulcers leading to the necrosed arytenoid cartilages they were from three to four times more abundant than in the other situations. Microscopical examination of sections from the region of the ventricular bands failed to show the presence of tubercle bacilli in the tissues.

As stated in the introduction, the features in the case worthy of notice are as follows:—

1. *The priority in origin of the laryngeal over the pulmonary affections.*—The history of the case undoubtedly points towards the larynx as being the part first affected. The commencement of a sore-throat, with hoarseness, could be traced to one particular evening when the patient, having placed himself in a draught, caught cold. He had hitherto enjoyed good health, and had no predisposition to tubercular disease. He was also assured by a medical practitioner whom he consulted about two years subsequently, in consequence of the persistent hoarseness, that his lungs were quite normal. The whole history and evidence pointed towards laryngitis, of a slow insidious form, as the starting-point of the disease. The examination on my first visit to the patient showed the laryngeal to be much more severe, and apparently of longer duration, than the pulmonary affection. A simple chronic inflammation of the larynx was undoubtedly the first link in the chain which led to such disastrous results to the patient.

2. *The engrafting of a tubercular on an initial simple inflammation of the larynx.*—It may reasonably be inferred that, not only at the commencement of the disease, but for some years subsequently, the condition present was one of simple inflammation of the larynx. Tuberculosis of the larynx lasting for seven years, and without being accompanied by extensive pulmonary changes, is, it is believed, quite unknown. In the present case a persistent chronic inflammation of the larynx disorganised and weakened the organ, and rendered it a suitable nidus for the reception and development of the tubercle bacilli, which readily reached it by inhalation. A progressive chronic inflammation appears to subserve this purpose more effectually than an acute inflammation, even where the latter leaves unmistakable traces of its presence in the form of permanent laryngeal changes. A case has been under constant observation during the last four years in which an acute laryngitis, at one time threatening suffocation, has played considerable havoc with the structures

and integrity of the larynx, but without evincing any tendency towards tuberculation. On the other hand, several cases, mostly in men, have been noted where, as in the present instance, a slow, gradually progressive inflammation of the larynx has undoubtedly ushered in tubercular disease both of the larynx and lungs. It is impossible to state with precision when the simple inflammation merges into the tubercular unless careful microscopical examination of the sputa be made from time to time with a view to the detection of the bacillus of tubercle. This is undoubtedly the variety of laryngeal tuberculosis in which the laryngeal precedes and leads to the pulmonary deposit, and may therefore justly be called primary laryngeal tuberculosis.

3. *The presence of the tubercle bacillus in the sputum and laryngeal secretion, and its value as an aid in diagnosis and prognosis.*—Of the laryngoscopic appearances in the case the utmost that could be stated was that they indicated a severe form of chronic laryngitis. The chest symptoms showed a bronchitic condition, with probably slight consolidation at right apex. The temperature chart afforded small indication, for the average of fifty observations was, morning 98°2', evening 98°6'. The discovery of the bacillus in the sputum supplemented in a most material way the other signs and symptoms, and afforded undeniable evidence that the case was of a more unfavourable character than was at first apparent. From March 28th, 1883, till the date of death (Feb. 15th, 1884), the sputum was examined on seventeen occasions, at about regular intervals, for the bacillus of tubercle, with the following results. On each occasion but one they were readily detected. In fair abundance in every field at first, their numbers afterwards seemed slightly to diminish, and in several fields they were not detected. They again slightly increased, and on Aug. 12th a very few were found in the laryngeal secretion on its removal by a brush. On Sept. 22nd, on testing the mucous and purulent portions of the same sputum, they were found from two to three times more abundant in the mucous than in the purulent portion. On Dec. 15th, after a hæmorrhage to the extent of one or two mouthfuls, marked signs of disease at both pulmonary apices, especially the right, and infiltration of the base of the right lung, they were present to the number of from fifteen to twenty in the field, rather isolated, but occasionally two lay in contact in a V-shaped arrangement, or three or four together, but not in contact. On January 16th, after steady persistent treatment by iodoform insufflation and inunction to the extent of inducing iodoform intoxication, the sputum, muco-purulent in character, contained as many as from forty to fifty in the field, generally not in contact. On Jan. 19th, under a continuance of the iodoform treatment, they could not be detected in the sputum, this being the only occasion when such was the case, but when the examination was repeated five days afterwards they were readily detected. It was probable, therefore, that some error in the preparation of the sputum interfered with their detection on the 19th. On Feb. 4th they were again found in the laryngeal secretion to the extent of fifteen in one field. The bacillary history of the case may therefore be shortly summarised by stating that these organisms assisted materially in the diagnosis, that they steadily persisted in the sputum, despite constant treatment by metallic astringents, antiseptic sprays, and iodoform insufflations to the larynx, and that their presence was not accompanied by febrile or other constitutional disturbance. Treatment by iodoform was pushed as far as was considered judicious, but without effecting any reduction in the number of the bacilli, although it diminished the putrefactive odour of the sputum. It is worthy of particular notice that on post-mortem examination, not only were they found in the secretion of the laryngeal ulcerations, but also in the mucus coating the crico-thyroid area, without subjacent ulceration. In the region of the ventricular bands and in the ventricles of the larynx, where they were from three to four times more abundant than in the crico-thyroid region, their presence was associated with deep destructive ulceration. This diversity may have been partly owing to the ventricles of the larynx affording a more secure nidus to these organisms, but probably partly also to the fact that the inferior regions of the larynx and the trachea are lined by a mucous membrane which is closely applied to the inner surfaces of the tube, and whose deeper layers (submucosa) contain strong bands of elastic longitudinal fibres. Whilst, in my experience, it is not uncommon to find erosion of the epithelium in the trachea and lower division of the larynx in phthisis, one

can never detect the same destruction as in the neighbourhood of the ventricular bands and arytenoid cartilages. The cartilaginous framework of the larynx and trachea has a resistant power which the softer tissues do not possess, and though, as would appear from the present case, the bacilli of tubercle cling to the whole of the respiratory tube, it is only in the softer tissues of the larynx and in the delicate alveoli of the lungs that their presence leads to deep destruction of tissue.

4. *Certain toxic effects following the use of iodoform.*—This drug was applied by insufflation and by inunction over the larynx and in the axilla. The difficulty in swallowing hindered its administration by the stomach. The formulæ used were, for insufflation, from one to five grains of iodoform with one-sixth to half a grain of acetate of morphia twice daily, and for inunction one part of iodoform to seven parts vaseline, also twice daily. When the larger doses of iodoform were reached, the patient became violent and quarrelsome, and a condition of excitement closely resembling the first stage of alcoholic intoxication was induced, which subsided on the discontinuance of the remedy. Considerable benefit to the local symptoms was experienced at the same time, but though the remedy was pushed as far as was conducive to safety, it was found that no effect was produced upon the bacilli of tubercle, which continued still to be found readily in the sputum. Inunction without insufflation also produced slight excitement and confusion of ideas. In a previous communication to a contemporary¹ I had already directed attention to the toxic action of the drug, and though the views then expressed by me did not find favour with certain continental authorities,² further experience has confirmed me as to their correctness. Dr. Batty Tuke, who saw the case now under notice with me, has informed me that, soon after seeing my patient, he was called to another case, in which the free use of iodoform produced the same symptoms as those above described, which disappeared on the discontinuance of the use of the drug. M. Maitre³ considers that, taken by man in doses of from five to six grains, it causes no notable symptoms; but my experience is that a smaller dose may readily induce toxic symptoms, particularly in weakly individuals.

With regard to the action of this remedy in laryngeal tuberculosis, although probably by its anæsthetic action it has a soothing effect on the throat, it does not seem to have any effect upon the disease. It does not diminish the number of bacilli in the laryngeal secretions, nor does it appear to promote the healing of the laryngeal ulcers. Little effect can, however, be expected from any remedies in such cases, unless use be made of them at very early periods of the disease, for its later stages are characterised by such an amount of laryngeal destruction and constitutional deterioration as invariably leads to a fatal termination.

Edinburgh.

A CASE OF LITHOTOMY WHERE THE NUCLEUS OF A LARGE PHOSPHATIC CALCULUS WAS THE WHALEBONE MOUTH-PIECE OF A TOBACCO-PIPE.

WITH REMARKS.

BY REGINALD HARRISON, F.R.C.S.,
SURGEON TO THE LIVERPOOL ROYAL INFIRMARY.

APART from the general interest attached to cases of this kind, which, curiously enough, are not very uncommon, there are, in this instance, two points upon each of which a few remarks may be made.

Robert H—, aged thirty-three, a seaman, was admitted on June 23rd, 1884, into the Liverpool Royal Infirmary suffering from symptoms of vesical irritation. The history elicited the following points:—Eight years previously the patient had gonorrhœa, from which he recovered quickly. In December last he had some scalding after micturition, and on the 13th of the same month he was admitted into a hospital in Wales with a fracture of the leg. Whilst in hospital he appears to have suffered from vesical irritation and passed two pieces of calculus; his symptoms continued

¹ Edinburgh Medical Journal, January, 1883.

² Centralblatt für klinische Medizin, 1883, No. ii.

³ Wood's Treatise on Therapeutics, p. 334. See also British and Foreign Medico-Chirurgical Review, vol. xxxiv., p. 247.