having ceased, the edges of the wound were placed in the most exact apposition, and retained in place by narrow pieces of tape attached through the medium of the adhesive solution. A small space was left at the lower extremity of the wound to permit the escape of any accumulation of fluid which might occur. Immediate union of the wound took place. On examination of the wound on the third day, it was found to be perfectly adherent throughout its whole extent, the line of the incision being scarcely perceptible a few feet distant, except at the point intentionally omitted, where the scar was of greater width.

JNO. PARKER MAYNARD.

COMMUNUTED FRACTURE OF THE PATELLA.

[PROF. P. F. EVE, of Augusta, Geo., relates the following case in a late number of the Southern Medical and Surgical Journal.]

This case produced great excitement at the time it occurred, or rather when agitated by a conflict with the law. It is the one, for which a suit of malapraxis was commenced against me in our Superior Court—damages laid at \$10,000—but which the plaintiff's attorney abandoned soon after my testimony was partially presented at the trial.

On the evening of the 16th of May, 1841, I was called to see Mr. B., in conjunction with my cousin, Dr. J. A. Eve. While driving a horse in a buggy, he had been kicked on the knee, and had fallen to the ground. The leg had therefore been flexed upon the thigh subsequent to the accident, which occurred about a mile out of the city, and an hour or two before we saw the patient. A transverse fracture of the patella was readily detected; as the patient was young and in good health, the immovable apparatus was proposed as the most certain means of securing union in the broken bone. This was accordingly applied at once, embracing the foot, leg and thigh. The usual number of daily visits were made, a dose or two of morphine and some opening medicines were administered, and on the 22d, six days after the accident, a roller bandage was applied over the whole apparatus to keep it in close contact with the limb, as it had become a little loose by drying and the shrinking in the soft parts. The next day the patient complained of pain, for the first time, in the knee-joint, and the whole dressing was removed. The limb was found cool and pale, owing probably to the compression, but chiefly to the position of it, it having been kept constantly elevated at an angle of about 45 degrees. The knee and the parts below soon after this began to swell, and Drs. Hook and Dugas were added to the medical council. The unfavorable symptoms continuing to increase, amputation of the thigh was performed on the 26th, ten days after the accident. The patient had a good recovery from this operation.

In examining the knee-joint, it was found filled with dark grunous blood, a portion of the cartilage of the internal condyle of the os femoris was chipped off, and the patella fractured into a number of fragments. The portion of the adductor magnus muscle which passes down to be

inserted into the internal condyle of the thigh bone, was moreover in a sphacelated condition.

On June 13th, I was called to attend to an infant in the father's family of this patient, which was so ill that it died. As late as August I was employed to adapt an artificial leg to the stump, and it was not until the next winter that the suit was brought against me.

A statement of the case, with interrogatories, were sent to eminent surgeons of Boston, New York, Philadelphia, Charleston, the interior of this State, &c., and by the aid thus obtained, through the efficiency of my attorneys, Messrs. Cumming and Jenkins, I was prepared for trial on the day appointed in June, 1842. My thanks are due to Dr. G. W. Norris, of Philadelphia, for putting at my disposition Seutin's work on the Im. Apparatus, just then arrived in this country; and I cannot refrain from inserting here the noble reply of Prof. Geddings, of Charleston, when the question was propounded to him, what is the result of such accidents as the one I was called upon to treat—"*death of the patient, either from mortification or tetanus.*" This controversy but the more fully convinced me that I belonged to an honorable profession, and was the associate of high-minded, dignified and liberal gentlemen. I have yet to learn that a want of success in every case is any disgrace, or that this trial ever injured me. My private class, the next winter after it, paid all my expenses, amounting to \$700. I have freely forgiven all who may have ever wished to slander or wrong me, and I can now allude to the circumstance without an unpleasant emotion.

THE BOSTON MEDICAL AND SURGICAL JOURNAL.

BOSTON, JUNE 21, 1848.

New Treatment of Bronchial Diseases.—Dr. William Gray, of New Hampshire, has constructed some curious apparatus for the purpose of inhaling certain medicinal vapors, with the object of allaying inflammations of the bronchial tubes, and lungs. A metallic box, some two inches in diameter and half an inch in thickness, contains a coiled steel wire. Over that he pours a volatile mixture—the wire affording extensive surface merely. When the box is closed, a flexible tube being attached to the side, the patient inhales the preparation. Of the benefits to be derived from the practice, those who have submitted to the process must be the best judges. We have merely examined the instrument, but have had no opportunity of conversing with gentlemen who have tested its properties. A leading idea of the inventor, if we understand him, is, that his method of applying a medicine to the inflamed membrane is an improvement on the common plan of introducing brushes charged with articles intended to produce the same result. And besides, Dr. Gray believes that he reaches farther, by his process, into the diseased lungs, than it is possible to do by any other means.