peared, and in few hours patient comfortable. In twenty-four hours the swelling of tongue and lips had disappeared. Case interesting because of locality; association with asthma, possibly with urticaria; and promptness of relief by morphine and atropine, *etc.* Suprarenal extract administration not supposed to have had any connection with causation of attack, because of immunity in its use, prior to, and since attack.

J. E. CLARK (New York).

INEQUALITY OF THE PUPILS IN THORACIC ANEURISM. R. C. B. Wall and E. W. A. Walker (Lancet, July 12, 1902).

After discussing the matter at some length and citing a number of cases, with evidence based upon clinical observation and experiment, the authors sum up their conclusions as follows: The explanation ascribing the anisocoria which sometimes occurs in cases of thoracic aneurism to interference with the sympathetic, is unsatisfactory. On anatomical grounds: (a) because in the majority of cases there is no evidence of implication of that portion of the sympathetic nerve trunk containing pupil dilator fibers, and (b) because it is not established that sympathetic filaments supply the sac wall of an aneurism. Further, alterations in vascular conditions may be associated with alterations in the size of the pupils, low arterial tension being associated with large pupils, and vice versa, phenomena for which the spiral structure of the vessels of the iris is probably responsible, and, therefore, local inequalities of blood-pressure may be associated with inequalities of the pupils. Clinical evidence has shown that enlargement of pupils is often associated with diminution of the temporal and radial pulses on the same side of the body, and obstruction of the carotid artery on one side of the neck is associated with enlargement of the pupil on the same side. By experiment upon the carotid artery in rabbits it was found that its obstruction caused enlargement of the pupils on both sides, owing to the freedom of circulation at the base of the brain, and that injection of water into the carotid artery of a dead rabbit caused enlargement of the pupil on that side. In the human subject digital compression of the carotid was also associated with enlargement of the pupil on that side. Their final conclusion was that inequality of pupils associated with thoracic aneurism is usually due to inequality of blood pressure in the ophthalmic arteries resulting from the abnormal vascular conditions.

JELLIFFE.

TREPHINING FOR BRAIN TUMORS. Joseph Ransohoff (Jour. of American Med. Asso., Oct. 11, 1902).

Operable brain tumors are rare. Only six per cent. of all cases are suitable for operation. The writer reports two successful cases. The first had suffered from convulsions beginning in the foot, rapidly extending over the side, followed by unconsciousness. He had violent headaches and frequent attacks of vertigo. Fundus shows some papillitis. Operation showed a tumor over the right foot center the size of a hen's egg. Recovery, with occasional light epileptic seizures, and a slight paresis of the arm and leg.

A second case of removal of a solitary tubercle from the ascending frontal convolution on account of muscular twitching and occasional loss of consciousness. No headache at any time. The patient recovered, with only a moderate amount of weakness of flexors of thumb and fingers. To make a diagnosis of cerebral tumor without headache is unusual. Against 104 cases of brain tumor in which operations were successful, there are 157 in which the operation was unsuccessful. Subcortical tumors particularly offer escape attention. If the dura and cortex are not involved headache often fails.

W. B. NOYES.