ness testifies that an ice-bag properly made, with drainage, produces "such a vasomotor dilatation, evidenced by an intense hyperemia, as no other means known to science can produce without injury to the parts covered." Baruch quotes from and endorses Woods Hutchinson to the effect that the whole arterial system participates in the rhythmic muscular action of the heart, of which it is a less specialized extension. He accepts the term "skin heart" as used by Hutchinson and holds that cold applications stimulate the vasodilators and increase the force of the rhythmic contraction of arterioles. This increased tonicity of the "skin heart," according to this theory, so greatly increases its propulsive power that the slow blood current in the skin is augmented to the relief of the central heart, and the more rapid elimination of waste and toxins.

It seems likely that the majority of the members of this society may not accept the theory of a "peripheral" or "skin heart." Baruch and Kellogg give a list of the men upon whose experimental work their faith is based. I would especially recommend the careful reading of Baruch's work because it is not only well written but is comparatively brief.

If our experience were your experience, I would think the reading of this paper hardly worth while. In response to the kind invitation of Dr. Gleason to lunch with you and talk to you I read you of those things which have proved helpful to us. We are not wedded to any theory or system of medicine. If the future shall reveal a specific against the Ebert bacillus or the Sternberg-Fraenkel diplococcus, we will welcome it gladly. In the light of our present experience, we hold we can best assist Nature by supplying liberally Nature's own life-sustaining remedies. As an east wind is to the heat-stricken children of the poor in Boston at the end of a torrid week in August, so pure cold water in liberal measure internally and intelligently applied externally* is to the fever-stricken children of the poor at all seasons.

NOTE 1. Acetozone is a mixture of equal parts of benzoylacetyle peroxide and an inert absorbent powder with a chemical formula of $C_{10}H_{12}O_4$. In the presence of water it undergoes hydrolysis and slowly decomposes. Such hydrolysis is said to be necessary to develop its full germicidal activity. This drug is one of the "organic peroxides" in which an excess of oxygen has been combined in such a way that it is somewhat slowly given off in a nascent condition. It is soluble in water at 77° F. in proportion of 1 to 1560.

NOTE 2. To obtain the best results, cold water given internally to fever patients should be at about 40° F. Cold compresses for adults should be dipped in water as near 60° F. as possible. For children under five, compresses at 80° are better. For children a full bath at 10° below the temperature of the body, with friction prolonged for twenty minutes, is very helpful. This general principle should be observed: the colder the water the shorter must be the application. Water is a potent remedy and may do harm if applied when the body is not in fit condition to react promptly to cold applications. That cold wet or cold dry applications should not be applied when the skin is cold or moist with perspiration goes without saying. When the skin is cold and clammy, the call is urgent for hot, dry flannels.

* See Note 2.

ARTERIO-VEINOUS ANASTOMOSIS.†

BY JOSHUA C. HUBBARD, M.D., BOSTON.

Assisting Visiting Surgeon, Boston City Hospital; Assistant in Surgery, Harvard Medical School.

My experience in arterial surgery has been obtained in two cases by doing an arterial venous anastomosis for reversal of the circulation. I feel that I have learned from them, although the number is small, certain facts which are worth recording. To make clear these points I will report the essential details of each case.

The first, already published in the *Annals of Surgery* for October, 1906, was that of a man of eighty years with senile gangrene of a portion of the right foot. Physical examination showed him to be a decidedly senile old man, with atheromatous arteries and a systolic heart murmur. No pulsation could be felt in the right dorsalis pedis artery. In May, 1906, he was operated upon. The femoral artery, which was decidedly atheromatous, and vein were isolated in Scarpa's triangle below the origin of the profunda and divided between Crille's clamps or elastic ligatures. The upper end of the artery was then invaginated into the lower end of the vein and the distal end of the artery into the proximal end of the vein. A complete reversal of the circulation was thus established. The technique of the invagination was as follows: Three double headed sutures were passed equally distant through the entire wall of the artery from inside out. The needles were then passed into the lumen of the vein about a quarter of an inch and at this point through its wall. When these sutures were drawn tight the artery was pulled into the vein. Re-enforcing sutures, including only the outer portion of the arterial wall, were placed here and there to catch the edge of the overlying vein to the artery. Number 1 Pagenstecher thread was used for all these sutures. When the controlling clamps were removed there was no leaking at either suture line and weak pulsations in the vein could be felt for a short distance below the anastomosis. There was absolutely no shock shown by the patient and recovery from the operation was satisfactory in spite of the fact that the senile condition of the patient made it difficult to keep him in bed or a dressing on the wound. After the operation the appearance of the leg did not change. There was no edema, dilatation of the veins or cyanosis. The gangrene which existed before the operation spread a little and then a line of demarcation formed. When the foot was later amputated, at the point of election on the tibia, both tibial arteries contained arterial blood. The stump healed satisfactorily but slowly. In March of this year, ten months after the operation, I saw the patient. He had a stump that was well supplied with circulation. There was no difference in appearance or size of this leg and the unoperated one.

Clinically this case was most successful, but the exact meaning of the presence of arterial

† Remarks made before the Academy of Surgery, Philadelphia, May 10, 1907.

blood in the tibial arteries at the time of the amputation I do not know. It seems that the arterio venous anastomosis must have increased in some way the amount of blood in the leg, for it is hard to believe that an amount of blood so small as to permit gangrene of the foot would be sufficient to nourish for ten months an amputation stump only a slight distance above the gangrenous area; and had clots formed at the sites of the anastomoses, it seems most probable that the gangrene would have extended rapidly up the leg instead of remaining localized.

The second case was that of an old woman of sixty years, with senile gangrene of the foot and a portion of the leg. The arteries were atheromatous. She was operated upon during February of this year. The femoral artery and vein were isolated in Scarpa's triangle and divided between clamps or ligatures below the origin of the profunda. The artery was much calcified and was so hard that some force was necessary to drive the needles through it and so brittle that the stitches tore out most easily. The intima formed a distinct lining to the vessel and was much like a second smaller tube inside a larger one. These characteristics of the wall complicated the technique immensely as I attempted to make my anastomosis according to Carrel's method, turning the walls so that at the suture line intima should come in contact with intima. The artery was so much like a pipe-stem that this was impossible, although a conscientious attempt was made. The ends of the artery and vein were therefore cut off freshly and the artery invaginated into the vein, as in the first case, the only difference being in the use of vaseline in this case to smear the ends of the vessels. The distal end of the artery and the proximal end of the vein were then ligated. The vein pulsated after the controlling clamps were removed. Pagenstecher No. 1 was used. There was no shock and no change in the appearance in the leg. Ten days later it was necessary to amputate above the knee for the gangrene which had been present before the operation. During these ten days it had become more pronounced, but its limits had extended only a little. During the amputation the anastomosis was cut down upon and removed. The artery was found filled with a loose, easily detached clot. This case was, therefore, a distinct failure.

From these cases I believe it is perfectly evident that there is no danger in continuing our investigations further, as there is no shock following the operation. Carrel's method is not applicable to a certain number of the cases where the operation is done on old persons with atheromatous arteries. In young persons and on experimental arteries it doubtless is most satisfactory, but as the operation has been proposed to cure conditions dependent upon lack of circulation in the extremities some other technique must be found, as practically all cases, except perhaps some due to trauma, will necessarily be in elderly persons. An objection which may be raised to the invagination method as employed in the above cases is

the fact that the divided end of the artery leaves a certain portion of its wall in the blood stream uncovered by intima, which favors clot formation. This method was introduced by Dr. Murphy * in 1897 for the repair of the continuity of an artery and was recommended only after experimentation. The slight modification of invaginating the artery into a vein instead of into another portion of the same artery would seem to me not to invalidate the method. However, as at present, this objection might be raised, I am working on some scheme to obviate this difficulty, but as yet unfortunately cannot report results. Two ways to avoid this have occurred to me. One is by smearing vaseline or some other substance onto the cut end of the invaginated artery to keep it out of the blood stream. The other way was suggested to me by the appearance of the arterial wall in the second case, where the intima formed a distinct layer inside the others and one which remained intact when the others cracked away from it. I have thought that it might be possible to cut the outer layers of the artery a quarter of an inch or so back of the intima and thus leave a greater length of intima as a cuff, the back of which could be covered with vaseline, so that when invaginated into the vein it might stick to the venous wall and cover over the cut end of the outer portion of the artery.

Clinical Department.

GLOTTIC SPASM OF NASAL ORIGIN IN ADULTS.

BY FRANK A. CONLON, M.D., LAWRENCE, MASS.
Former Hospitant in the Koschier Clinic, Vienna.

SPASM of the glottis of nasal origin in adults, a rare form of reflex laryngeal neuroses, was first described by Hack in 1882, and since that time fourteen cases have been published, making in all but fifteen recorded cases. To these I propose to add the sixteenth, but before doing so shall attempt to give a short résumé of the cases so far reported.

CASE I. Hack.¹ A healthy physician, whose disposition was not at all nervous, suffered at night severe attacks of dyspnea, caused by glottic spasm. Examination of posterior nares disclosed a marked congestion of the mucous membrane. After painting the posterior nares with silver nitrate, attacks disappeared for a long time, only to reoccur with the advent of a mild angina. Posterior nares again painted and attacks have not since reoccurred.

CASE II. Heryng reports two cases. Man,² forty-nine years old, who had been suffering from spasm of the glottis for three years, the attacks occurring during day and night. Examination of the larynx showed nothing abnormal. Posterior nares and pharynx normal. Anterior rhinoscopy showed polyp in left nares in region of middle turbinate. Polyp was removed and patient was entirely free from symptoms for a year. He then returned, suffering fresh attacks, and on examining the nose a new polyp was found.

* Medical Record, January 16.