in the premaxillary, as they are below or in front of the nasal aperture. The form of the teeth, both crown and root, is very similar to the teeth of *Hesperornis*. The fact that some teeth are scattered about near the jaw would suggest that they were implanted in a groove. No teeth are known from the lower jaw, but they were probably present.

The presacral vertebrae are all, or nearly all, biconcave, resembling those of *Ichthyornis* in general form, but without the large lateral foramina. There appear to be twenty-one presacral vertebrae, and the same, or nearly the same, number of caudals. The sacral vertebrae are fewer in number than in any known bird, those united together not exceeding five, and probably less.

The scapular arch strongly resembles that of modern birds. The articulation of the scapula and coracoid, and the latter with the sternum is characteristic; and the furculum is distinctly avian. The sternum is a single broad plate, well ossified. It probably supported a keel, but this is not exposed in the known specimens.

In the wing itself the main interest centres in the manus and its free metacarpals. In form and position these three bones are just what may be seen in some young birds of to-day. This is an important point, as it has been claimed that the hand of *Archaeopteryx* is not at all avian, but reptilian. The bones of the reptile are indeed there, but they have already received the stamp of the bird.

One of the most interesting points determined during my investigation of *Archaeopteryx* was the separate condition of the pelvic bones. In all other known adult birds, recent and extinct, the three pelvic elements, ilium, ischium and pubis, are firmly ankylosed. In young birds these bones are separate, and in all known Dinosaurian reptiles they are also distinct. This point may perhaps be made clearer by referring to the two diagrams before you, which I owe to the kindness of my friend Dr. Woodward, of the British Museum, who also gave me excellent facilities for examining the *Archaeopteryx* under his care. In the first diagram we have represented the pelvis of an American Jurassic Dinosaur allied to *Iguanodon*, and here the pelvic bones are distinct. The second diagram is an enlarged view of the pelvis of the *Archaeopteryx* in the British Museum, and here too the ilium is seen separate from the ischium and pubis.

In birds the fibula is usually incomplete below, but it may be coossified with the side of the tibia. In the typical Dinosaurs, *Iguanodon*, for example, the fibula at its distal end stands in front of the tibia, and this is exactly its position in *Archaeopteryx*, an interesting point not before seen in birds.

The metatarsal bones of *Archaeopteryx* show, on the outer face at least, deep grooves between the three elements, which imply that the latter are distinct, or unite late together. The free metacarpal and separate pelvic bones would also suggest distinct metatarsals, although they naturally would be placed closely together, so as to appear connate.

Among other points of interest in *Archaeopteryx* may be mentioned the brain-cast, which shows that the brain, although comparatively small, was like that of a bird, and not that of a Dinosaurian reptile. It resembles in form the brain-cast of *Laopteryx*, an American Jurassic bird, which I have recently described. The brain of both these birds appears to have been of a somewhat higher grade than that of *Hesperornis*, but this may have been due to the fact that the latter was an aquatic form, while the Jurassic species were land birds.

As the *Dinosauria* are now generally considered the nearest allies to Birds, it was interesting to find in those investigated many points of resemblance to the latter class. *Compsognathus*, for example, shows in its extremities a striking similarity to *Archaeopteryx*. The three clawed digits of the manus correspond closely with those of that genus; although the bones are of different proportions.

The hind feet also have essentially the same structure in both. The vertebrae, however, and the pelvic bones of *Compsognathus* differ materially from those of *Archaeopteryx*, and the two forms are in reality widely separated. While examining the *Compsognathus* skeleton, I detected in the abdominal cavity the remains of a small reptile which had not been previously observed. The size and position of this inclosed skeleton would imply that it was a *fetus*; but it may possibly have been the young of the same species, or an allied form, that had been swallowed. No similar instance is known among the Dinosaurs.

A point of resemblance of some importance between Birds and Dinosaurs is the clavicle. All birds have these bones, but they have been considered wanting in Dinosaurs. Two specimens of *Iguanodon*, in the British Museum, however, show that these elements of the pectoral arch were present in that genus, and in a diagram before you one of these bones is represented. Some other *Dinosauria* possess clavicles, but in several families of this subclass, as I regard it, they appear to be wanting.

The nearest approach to Birds now known would seem to be in the very small Dinosaurs from the American Jurassic. In some of these the separate bones of the skeleton cannot be distinguished with certainty from those of Jurassic Birds, if the skull is wanting, and even in this part the resemblance is striking. Some of these diminutive Dinosaurs were perhaps arboreal in habit, and the difference between them and the Birds that lived with them may have been at first mainly one of feathers, as I have shown in my Memoir on the *Odontornithes*, published during the past year.

It is an interesting fact that all the Jurassic birds known, both from Europe and America, are land birds, while all from the Cretaceous are aquatic forms. The four oldest known birds, moreover, differ more widely from each other than do any two recent birds. These facts show that we may hope for most important discoveries in the future, especially from the Triassic, which has as yet furnished no authentic trace of birds. For the primitive forms of this class we must evidently look to the Palaeozoic.

THE LIMITED BIOLOGICAL IMPORTANCE OF SYNTHETIC ACHIEVEMENTS IN ORGANIC CHEMISTRY.*

By PROFESSOR ALBERT B. PRESCOTT.

The solicitude shown for half a century as to the biological import of chemical synthesis arises from a misapprehension of the scope of chemical action. From all we know of chemism, it must be accepted, (1) that all the matter of protoplasm and cell is carried strictly in a state of chemical combination, but (2) it cannot therefore be accepted that chemical composition supplies the essential conditions or impulses for organization or other vital functions. The synthesis of all the chemical compounds of the living body may or may not be attainable in the laboratory, but its success would give no whit of promise for the development of organization. Chemical action is distinct from cell organization as it is from heat, cohesion, etc., and its correlations with all these forces have to await demonstration by experiment. Cell growth appears to be a necessary factor in the simple splitting of sugar into alcohol and carbon dioxide, and it may or may not be an essential factor in the chemical synthesis of proteids or of cellulose.

A GENTLEMAN of Milan, Signor Lorin, deserves high credit, for the public spirit of philanthropy he has shown in offering 20,000 francs to the municipal authorities to maintain a mortuary and post mortem room wherein the bodies of all persons dying of unexplained causes shall be rigidly examined before they are cremated.

* Read before the A. A. S., Cincinnati, 1887.