VI.—The “Red Snow” Plant (*Sphaerella nivalis*).

By G. S. WEST, M.A., D.Sc., F.L.S.

(Read February 17, 1909.)

THE red coloration of extensive tracts of perpetual snow in alpine and arctic situations is caused by a small alga known as *Sphaerella nivalis* Sommerf. It belongs to the sub-family Chlamydomonadeæ of the Volvocaceæ, and is very closely related to *S. lacustris* (Girod.) Wittr. (= *Hæmatococcus pluvialis* Flotow), the tiny alga which so often gives a red colour to drying-up rain-water pools and tanks.

It is in the resting stage of the alga that the red pigment is chiefly developed, and therefore the beautiful red coloration of the snow, which sometimes causes the admiration and astonishment of the traveller and explorer, is due to the presence of myriads of the resting-cells of *Sphaerella nivalis*.

There are no authentic records of “red snow” in the British Islands due to this alga, and, owing to the peculiar physiological conditions necessary for the completion of its life-cycle, it is scarcely likely that it will be found. It occurs in the Austrian, Swiss, and Italian Alps, and in Norway, the red tinge appearing towards the month of August. It has also been found amid the perpetual snows of other regions, more especially in the South American Andes, and in the arctic and antarctic snow-fields.

It possibly occurs throughout most of the arctic snow-fields, and the present article was written on the examination of some particularly fine resting-cells recently presented to the Society by the Peary Arctic Club.

Each resting-cell is a small globular body (17–24 μ in diameter), consisting of a protoplast inclosed in a fairly thick wall of cellulose. Pyrenoids and a nucleus are present, but are hidden by a red pigment known as “hæmatochrom,” which permeates the whole protoplasm, and exists partly in solution in minute drops of oil, and partly in a microcrystalline form. A small quantity of chlorophyll is also present, even in the purely red cells. The exact function of this hæmatochrom is a perplexing problem. It is certainly more stable than chlorophyll, and generally accompanies diminished vitality, which fact accounts for its presence in the resting-cells. The cells can thus probably exist through adverse circumstances much better than if they were merely green, not

being so easily destroyed by sudden changes in light and temperature. Both Kny and Kerner have shown that red colouring matters are probably concerned in the conversion of light-rays into heat, and the great value of such a function to the "red snow" plant is at once obvious. Moreover, Hazen has pointed out that in the closely related *Sphaerella lacustris*, more red pigment is habitually developed in cultures made from material collected from places frozen for a great part of the year, than in those made from material collected in milder climates.

The life-history of this organism is briefly as follows. The typical resting-cell forms four daughter-cells of a similar bright red colour, which remain for a time within the wall of the mother-cell. So long as unfavourable conditions supervene, these daughter-cells remain in the resting state, although often with a slight increase in size. Should the conditions become favourable, however, the daughter-cells very quickly escape from the mother-cell, develop a pair of cilia, and become motile. Each is now a zoogonidium (or asexual motile gonidium), and in this state the cell generally increases in size. The cell-wall is thin and distended, standing out from the main body of the protoplast; but much detailed observation is yet required on the motile state of *Sphaerella nivalis*. After a longer or shorter active period, the zoogonidium returns to the resting state, and the main body of the protoplast develops around itself a thick wall within the delicate wall it possessed in the motile state. A sexual method of reproduction by the conjugation of isogamous planogametes ("micro-zoogonidia") has been described. The resulting zygospores are said by Wittrock to be verrucose at maturity.

Sphaerella nivalis appears to have adapted itself to very low temperatures, but it does not stand alone in this respect. There are several algæ, belonging to widely separated groups, which live habitually in the snow, and form a small assemblage of plants known as the Cryoplankton.

Prolonged exposure to a low temperature is physiologically necessary for the "red snow" plant, otherwise it loses its vitality. Chodat has stated that this alga will not live for any lengthened period at a temperature above 4° C., but experiments conducted both in this country and the United States do not entirely support this statement, as the resting-cells can be kept alive in water for over twelve months at ordinary outside temperatures.

The distinctions between *Sphaerella nivalis* and *S. lacustris* are not very great, and many authors have considered them to be specifically identical. The only difference of importance appears to be the presence in the motile vegetative cells of *S. lacustris* of protoplasmic threads connecting the central mass of the protoplast with the outstanding cell-wall, whereas in *S. nivalis* these threads are absent. Wollenweber states that *S. lacustris* has numerous

contractile vacuoles which pulsate independently of one another, and that this is one of the chief distinguishing characters between *Sphaerella* (*Hæmatococcus*) and *Chlamydomonas*. He agrees with Wille that *Sphaerella nivalis* should be regarded as a species of *Chlamydomonas*, *C. nivalis*, but before this view can be definitely accepted the motile state of *Sphaerella nivalis* requires a good deal of detailed investigation.