not be the lot of the generality of patients to have a laboratory close at hand, and if by chance the bite is overlooked, and the disease actually gains its ascendancy in the blood, have we got any medicine in our Pharmacopœia on which the slightest reliance can be placed for treatment? The sad confession in the weakness of our resources will then be manifest, and, while brooding over our failures, it struck me that Nature could not be so improvident and callous as to leave us so thoroughly helpless and unprotected.

Starting from the acknowledged action of hydrophobic poison on the blood, can we not find another poison that has just an opposite virtue and likely to neutralise its effect? We have in the animal kingdom two deadly poisons, which differ entirely from each other in their physiological action. Thus, whilst the poison of snake-bite acts with the rapidity of an electric current, that of hydrophobia takes a long incubative period to mature itself. The effect of both is on the cerebro-spinal system, chiefly the medulla. Rabies produces death by causing spasm of the respiratory centres; snake-poison, paralysis. One causes hyperæsthesia of the cutaneous nerves; the other, numbness. One gives rise to excitement of the brain and furious delirium; the other, coma. In every detail, therefore, of its action, one is in physiological opposition to the other, and, applying the law of contraries, it stands to reason that the effect of one should neutralise the other in the system. Just in the same way as opium and belladonna, being two deadly poisons, their antagonistic virtue was discovered simply on the observance of their different effect on the pupils.

The suggestion may startle the readers as a bold and dangerous practice, but a deadly disease requires a deadly remedy, and we are apt to forget that in the whole range of therapeutics the most potent remedies are the most deadly poisons. But though poisons in ordinary doses, the attenuation of the quantity by dilution may yet bring them within the range of remedial measures. It is known that a snake-bite which is not effective to be fatal simply produces a deleterious effect, the symptoms varying in proportion to the quantity of the poison injected into the system. A smaller dose therefore would produce the physiological effect, instead of bringing on death and destruction. We have, consequently, to hit upon the happy medium, and this can be worked out satisfactorily by observing that a good bite of cobra instils ten to fifteen drops of the poison into the tissues. One drop diluted with glycerine and injected subcutaneously, and repeated every two hours according to the symptoms observed, affords a rational presumption for success in the treatment of hydrophobia: thus—snake-poison, twenty-four minims; glycerine, half an ounce; ten drops to be injected every two hours under the skin.

All the improvements in medical science in former days were deduced from empiricism. In fact, few remedies were discovered on rational philosophy. I have, on rational principle, suggested the treatment to which I would earnestly solicit the profession to give a trial, for it falls to the lot of few to treat one case of hydrophobia from one year's end to the other, and unless the co-operation of brother practitioners be accorded, our life is too short to enable us to establish one single truth in a generation. Cobra poison is plentiful in the rains in India, and, diluted with glycerine, it will keep its virtue intact for a year. I shall be glad to supply the mixture in August next to those who kindly feel disposed to put the method to the test.

SANTONIN IN AMENORRHOEA.

By AMAND ROUTH, M.D., B.S., M.R.C.P.,

ASSISTANT-PHYSICIAN TO CHARING-CROSS HOSPITAL, AND PHYSICIAN TO THE SAMARITAN FREE HOSPITAL FOR WOMEN AND CHILDREN.

BELIEVING that the re-establishment of the catamenia in cases of amenorrhœa with constitutional disease (especially chlorosis) is helpful in restoring the patient to health, and knowing also that general remedies may be some months in producing that result, I have tried santonin, as suggested by Mr. Walter Whitehead (THE LANCET, Sept. 5th, 1885)—viz., ten grains of powdered santonin to be taken on two consecutive nights. I omitted, however, the morning saline purge, wishing to avoid possible sources of error. I have tried the drug in twenty cases of amenorrhœa, thirteen of which were associated with anæmia or chlorosis, and failed

completely in seventeen. In all, restlessness, yellow vision, and discoloured urine followed the use of the drug. The following four cases are of interest:—

CASE 1.—Harriet V—, aged eighteen; plethoric and epileptic; has never menstruated. Eight hours after taking the first powder, violent colic with diarrhœa came on, with an unusually severe epileptic fit, during which the menstrual flow appeared, and continued painlessly for four days. She has not menstruated since (three months and a half), but has had one fit.

CASE 2.—Alice K—, aged nineteen; chlorosis; has never menstruated; much ovarian tenderness; slight leucorrhœa. After the two powders a profuse leucorrhœal discharge ensued, lasting six days, with much uterine colic, but no catamenia. Patient is now quite well and regular, after two months and a half of treatment with iron, arsenic, salines, and aloetic purgation.

CASE 3.—Annie B—, aged twenty-one; chlorosis; amenorrhœa three years. The powders produced no effect, but two weeks afterwards three grains of permanganate of potash, with kaolin in sufficient quantity, in form of pill, and taken three times a day after meals, brought on the flow after taking twelve pills. Patient is now nearly cured with the usual remedies.

CASE 4.—Mary E. R—, aged twenty-four; quite healthy; catamenia overdue eight days, though usually regular, and patient is to be married in six days. Catamenia came on during the night after one powder, and for the first time in her life painlessly, lasting the usual four days.

Remarks.—My twenty cases seem to indicate that santonin, as a direct emmenagogue, is useless in amenorrhœa associated with chlorosis, in which cases I have often found permanganate of potash useful. In healthy, or full-blooded women, santonin seems to have some power in starting a uterine flow, and apparently without causing much pain. Its mode of action, however, is quite unknown in this respect, and it seems not likely therefore to be much employed in gynecological therapeutics.

NOTE ON THE QUESTION OF THE POISONOUS PROPERTIES OF THE SALTS OF COPPER.

By CHARLES H. PIESSE, F.I.C., M.R.C.S.,
PUBLIC ANALYST.

THE important question as to whether the compounds of copper are poisonous or not has been recently investigated by M. Dumoulin, Professor of Toxicology in the University of Ghent. A report of the paper he read on the subject before the Belgian Academy of Medicine appeared in the *Semaine Médicale* of the 2nd ult., of which the following is an abstract:—"His first experiments were made on rabbits and dogs, to which he administered doses varying from four to six grammes (sixty-two to ninety-three grains) of copper-sulphate; in no case was the effect serious; the whole of the animals bore perfectly well the ingestion of the copper salts. He next tried the subacetate (verdigris) upon the same class of animals; they vomited at first, but after the administration of further doses the vomiting ceased, and they bore daily doses of the salt during a period varying from six weeks to one year. The carbonate, the oxide, and the oleopalmitate, of which doses varying from seven to fifteen grains were administered, gave equally negative results, the last-named compound being borne with the greatest ease. Toussaint and several German writers had administered to human beings doses rising to seventeen grammes (263 grains) daily of salts of copper without disturbance. It happened that one of the dogs experimented upon by M. Dumoulin was suffering from impetigo. He noticed that the disease disappeared, and attributed the cure to the copper salt. He therefore in the same way treated some children suffering from impetiginous eruption of the face and scalp, and without accident cured the eruption. He then successfully extended the treatment to all forms of scrofulous symptoms, to various superficial skin diseases, ophthalmias, &c. He has now under treatment some scrofulous children aged about five or six years, to whom he administers about one grain of copper sulphate daily, with the result that the