In line with the general policy of coordinating the state wide service, the state has just authorized the creation of a psychopathic department at the university with a special hospital and adequate support for research work in this field.

This enlarged consciousness on the part of the public as to its responsibility in regard to health is by no means confined to a few states. Indeed, steps are being taken to secure the leadership of the federal government in assuming this responsibility. There is little doubt that within the next decade health service under some such general plan as outlined above will be provided almost everywhere. The states will have to determine whether or not they wish to establish this service in isolated hospitals as has been done in the past in the case of sanatoriums for the tuberculous, hospitals for the insane and the like, or whether they wish to coordinate this work with that of the training of physicians in the college of medicine.

While it is true that in some schools the supply of clinical material is so large that these patients would be of little value in connection with the training of physicians, yet in many states it would be disastrous from the standpoint of the teaching clinic for the state to establish such a hospital apart from the college of medicine, necessitating, as it would, constant duplication of the staffs of the two institutions. The waste would be calamitous. The experience in Iowa is such as to suggest that this type of work can most successfully be done in connection with the college of medicine.

CONCLUSIONS

Our experience leads to these conclusions:

1. In view of the great rapidity with which the demands on our staff and hospital have grown, it seems to us important to note that any state in attempting to provide this type of service should make very comprehensive plans on the material side, which should include liberal provision in space and staff not only for adequate service to the indigents but also for adequate provision for the ever growing number of pay patients.

2. It is very important that future plans include ample provision for the vigorous prosecution of medical research. Otherwise the teaching staff may easily be overwhelmed with routine, with a consequent slump in growth. Furthermore, the unusual clinical demands serve as a constant challenge to the student of medicine.

3. Since the success of the work is absolutely dependent on skill and devotion of the staff, it is essential that many adjustments in the conditions of teaching must be made. The problem of full-time teaching becomes more acute.

Satisfactory adjustments of these problems must be worked out if medical teaching is to be kept at the highest standard.

Other departments in the field of technical education have had to meet the same situation. Indeed, in the field of agricultural education a very large part of the function has been the providing of service for the public. So much is this true that it would be hard to conceive at the present moment of a college of agriculture without its elaborate organization in the direction of special agricultural service to the state. Within the next few years we may expect in many state universities just such close coordination in the problems of the training of physicians, furthering research that will contribute to the knowledge of the field and extending health service to the public.

To the degree that the colleges of medicine of the state universities are alert to these new demands and effective in their responses will these colleges become real leaders in this present movement looking toward the highest type of physical and mental efficiency.

CHANGES IN THE BLOOD IN INFLUENZA *

RALPH A. KINSELLA, M.D.

AND

GORONWY O. BROWN, M.D.

ST. LOUIS

The tendency to nosebleed, the bloody sputum and the hemorrhagic character of the lesion in the lung, in an uncomplicated case of epidemic influenza, give rise to the inference that there are changes in the blood in this disease which predispose to hemorrhage.

TABLE 1.—EFFECT OF SODIUM CITRATE ON CLOTTING
TIME OF THE BLOOD

Name or Hospital Number	Date	Clotting Time, Minutes	Patient's Tempera- ture, F.	Remarks
D.	2/ 6/20	15	100.6	Before receiving citrate
D.	2/ 6/20	7		45 minutes after receiving 1 gm. sodium citrate intravenously
D.	2/ 7/20	7		18 hours after foregoing dose of citrate
D.	2/ 7/20	3½		1 hour after a second dose of 1 gm. sodium citrate intravenously
11278	2/ 6/20	10	105.0	Before receiving citrate
11278	2/ 6/20	5		¾ hour after receiving 1 gm. sodium citrate intravenously
11278	2/ 7/20	7		18 hours after the foregoing dose of citrate
11278	2/ 7/20	3½		1 hour after a second dose of 1 gm. sodium citrate intravenously
11350	2/20/20	12	99.0	Before receiving citrate
11350	2/20/20	3½		½ hour after receiving 1 gm. sodium citrate intravenously
11795	2/19/20	16	101.6	Before receiving citrate
11795	2/10/20	9		½ hour after receiving 1 gm. sodium citrate intravenously
11795	2/20/20	9		24 hours later
11795	2/20/20	9		½ hour after receiving a second dose of 1 gm. sodium citrate intra- venously
11844	2/20/20	14½	99.4	Before receiving citrate
11844	2/20/20	6¾		½ hour after receiving 1 gm. of sodium citrate intravenously

It was for the purpose of inquiring into this point that the present study was undertaken, and this report is a preliminary summary of the results that were obtained.

The special features studied were the clotting time, the platelet count, the fragility of the red cells, and the number of leukocytes.

METHODS

For studying the clotting time, blood was drawn from the median vein into a syringe wet with physiologic sodium chlorid solution, and 1 c.c. was immediately inserted in tubes 1 cm. in diameter also wet with physiologic sodium chlorid solution. It was found that if the syringe was one of small capacity and the quantity of blood drawn only a little more than that necessary for the test, the clotting time was much shorter than if a large syringe was used and the quantity of blood drawn was several cubic centimeters more than necessary for the test. The cases in Table 3 were studied by the latter method. Care was taken to use exactly the same technic for test cases and the normal controls.

* From the Medical Clinic, City Hospital, St. Louis University School of Medicine.

For counting the platelets, the technic described by Wright and Kinnicutt¹ was used. Counts of normal individuals were made as controls.

For estimating the fragility of red cells, the method described by Hill² was employed.

TABLE 2.—EFFECT ON CLOTTING TIME OF ADDITION OF VARIOUS AMOUNTS OF SODIUM CITRATE TO BLOOD IN THE TEST TUBE

Before Intravenous Administration of Sodium Citrate						
Amount of Citrate Added to 1 C.c. Blood	Clotting Time in Minutes					
	Patient 11350	Patient 11795	Patient 11795*	Patient 11844	Normal 1	Normal 2
Plain blood; no citrate.....	12	16	9	14½	9	10
0.005 mg.	12	..	9	16	..	8
0.0075 mg.	11½	..	11	14½	..	..
0.01 mg.	11	11	7	11½	..	11
0.025 mg.	11	..	10	11	..	9
0.05 mg.	12	..	15	14½	9	11
0.1 mg.	12	15	8	13½	9	11
0.5 mg.	17	19	10½	30	18	..
1.0 mg. (partial)...	45	Did not clot	Did not clot	Did not clot	Did not clot	20
2.0 mg.	Did not clot	Did not clot	Did not clot	Did not clot	Did not clot	Did not clot
2.5 mg.	Did not clot	Did not clot	Did not clot	Did not clot	Did not clot	Did not clot
Thirty Minutes after Administration of Sodium Citrate (1 gm.)						
Plain blood; no citrate.....	3½	9	9	6¾	..	..
0.005 mg.	3	..	7	7	..	..
0.0075 mg.	3½	..	9	5	..	..
0.01 mg.	3½	16	13	7	..	..
0.025 mg.	3½	..	13	10	..	..
0.05 mg.	3½	..	6	11	..	..
0.1 mg.	3½	17	10	10	..	..
0.5 mg.	3½	17	16	45	..	..
1.0 mg.	30	Did not clot	Did not clot	Did not clot	..	..
2.0 mg.	Did not clot	Did not clot	Did not clot	Did not clot	..	..
2.5 mg.	Did not clot	Did not clot	Did not clot	Did not clot	..	..

* The two readings of Patient 11795 were taken twenty-four hours apart, and after the first test, 1 gm. of sodium citrate was given as was done in each of the other cases.

RESULTS

Clotting Time.—Forty patients were studied. In all, the diagnosis of influenza was clear from the clinical manifestations. All occurred during the height of the epidemic. Some were in the second or third day of convalescence. The majority were studied between the third and sixth day of disease. The normal clotting time with the method used was definitely shorter than that of the patients. There seemed no relationship between degree of delay and severity of disease. Moreover, the delay was present even when secondary infections with pneumococci or hemolytic streptococci had occurred. It was as though the delay in clotting time represented an influence essentially associated with the intoxication of influenza, which persisted throughout the secondary pneumonia, and into convalescence.

Platelets.—The results are shown in Table 3. Counts were made in twenty-one cases. The reduction in the majority of the cases is very definite. As in the case of clotting time, the reduction of platelets was independent of the stage of the disease or of the presence of secondary infections. The reduction of platelets and the delay in clotting time occurred hand in hand. Both were apparently independent of the number of leukocytes present, as shown by Table 3.

Leukocytes.—In general, it may be said that influenza is attended by an absence of leukocytosis, often

extending to a degree of leukopenia. The presence of leukocytosis is good evidence of the presence of secondary infection.

COMMENT

As pointed out by Weil,³ intravenous injection of sodium citrate hastens clotting, although in the test tube the same salt in sufficient concentration will prevent clotting. Similar action is ascribed by Addis⁴ to sodium phosphate. To determine whether the same response to sodium citrate could be obtained in patients with influenza, possibly affecting the condition thereby, five patients were studied as shown in Table 1. The effect of intravenous injection was prompt. In one case (11278), the character of the sputum markedly changed from one of bloody appearance to one almost free from blood. It was thought, in the beginning, that patients improved after such treatment; but further tests made such a conclusion doubtful. In patients with streptococcus empyema, a peculiar reaction usually followed the injection of citrate, which consisted in a sudden, though transitory, gasping for breath. In the test tube, the addition of sodium citrate seems to inhibit the hemolytic action of serum of such patients on a suspension of sheep red blood corpuscles. Accordingly, the explanation of this clinical reaction was not found.

In seeking an explanation for the peculiar effect of sodium citrate on clotting time in vivo, it was thought that the salt might rapidly diffuse after entering the

TABLE 3.—COMPARISON OF LEUKOCYTE COUNT, PLATELET COUNT, CLOTTING TIME AND FRAGILITY OF RED BLOOD CORPUSCLES, IN INFLUENZA PATIENTS

Hospital Number	Leukocyte Count	Platelet Count	Fragility Test		Clotting Time, Minutes
			Complete Hemolysis, % NaCl	No Hemolysis, % NaCl	
10953	14,000	70,000	0.300	0.550	7½
11133			0.250	0.500	7
11149	3,000	38,000	0.250	0.500	6
11174	9,500	45,000	0.250	0.475	7
11250	11,000	66,000	0.300	0.425	11
11278	8,000	85,000	0.200	0.350	5
11405	9,000	78,000	0.300	0.425	12
11406	6,000	114,000	0.250	0.425	6
11420	7,000	89,000	0.325	0.425	11
11424	2,000	28,000	0.250	0.550	4
11428	3,000	47,000	0.300	0.475	7
11458	6,000	64,000	0.250	0.425	10
11469	10,000	67,000	0.250	0.400	10
11487	3,200	76,000	0.250	0.475	6
11367	4,200	34,000	0.300	0.475	4
11538	20,000	220,000	0.300	0.425	5
11546	4,500	20,000	0.300	0.425	9
11153	40,000	60,000			
11566	2,000	52,000	0.350	0.475	8
11595	15,000	133,000	0.300	0.425	12
11596	18,000	66,000	0.300	0.425	14
D.	18,000	285,000	0.250	0.450	7
Normals					
H.			0.250	0.475	
B.	8,000	258,000	0.250	0.450	4½
G.			0.250	0.475	
K.	8,000	235,000			4½
A.	12,000	224,000			5
Miss A.			0.300	0.425	

blood stream and be diluted so that only a minute amount remained after a few minutes, and that such a minute quantity might enhance the condition of "thrombin" and "prothrombin." Accordingly, tests were made of four patients and two normal controls, in which minute amounts of citrate in vitro seemed to have the effect of shortening clotting time, while the usual concentration of 0.2 per cent. prevented clotting. Further investigation must be made to substantiate this finding, shown in Table 2.

1. Wright, J. H., and Kinnicutt, Roger: A New Method of Counting the Blood Platelets for Clinical Purposes, J. A. M. A. **56**:1457 (May 20) 1911.

2. Hill, L. W.: The Resistance of the Red Blood Cells to Hypotonic Salt Solution in the Various Anemias, Arch. Int. Med. **16**: 809 (Nov.) 1915.

3. Weil, Richard: Sodium Citrate in the Transfusion of Blood, J. A. M. A. **64**: 425 (Jan. 30) 1915.

4. Addis Thomas: Proc. Soc. Exper. Biol. & Med. **14**: 1192, 1916.

SUMMARY

The clotting time of the blood of patients with influenza is delayed.

The number of platelets is reduced in such patients.

Neither of the foregoing conditions changes with the onset of secondary infection, although the number of leukocytes is usually greatly increased in secondary infection.

The fragility of the red cells is probably increased in influenza.

CAN THE TUBERCULOSIS TRANSMISSION RATE BE REDUCED?

JAMES G. CUMMING, M.D. (ANN ARBOR, MICH.)

Lieutenant-Colonel, M. C., U. S. Army

A. E. F., VLADIVOSTOK, SIBERIA

What are the possible avenues of tuberculosis transmission among individuals? Is it direct contact by the air route, though spray that is expelled by an act of coughing, or through dried sputum in public places? In answer to this question it may be said, that if the results of investigations of the influenza-pneumonia epidemic can be taken as a key to sputum-borne disease transmission, the transmission rate resultant from the distribution of tuberculosis by the air route among adults is of minor importance.¹ In the infant, however, another link may be added to this avenue of transmission: that of soiling the hands on a contaminated floor.

In contrast to direct contact, transmission may take place through several common routes by indirect contact: (a) through hand-to-mouth infection, the hands being soiled by contaminated inanimate objects, and (b) through inanimate object-to-mouth infection, the objects entering the mouth of the tuberculous as well as that of the nontuberculous. There are other ramifications in the routes of travel, but for the purpose of this paper the foregoing are the most important.

Here it might be said that our efforts to control the disease by the restriction of spitting and coughing have not given results that inspire a high degree of confidence. Should it be assumed that public spitting is a large factor in transmission, it is to be pointed out that this restrictive measure is of limited value because it is an attempt to deal with the individual. In this disease, as in all communicable diseases, a successful attack cannot be made through the individual—the animate—but must be made through the inanimate; or, as in insect-borne diseases, through insect destruction. It might be laid down as fundamental, from an epidemiologic standpoint, that the final solution in the eradication of any disease depends not on the control of the animate, but on the control of the inanimate through universal mass action.

On this basis there would be some encouragement in a plan of attack which could be universally applied to the most common *saliva*-contaminated inanimate objects. If this could be accomplished, the major avenue of transmission would be closed, the transmission rate would be immediately diminished, and on the basis of "the theory of case and source elimination" the majority of the tuberculous cases of the next human cycle of this organism would be eliminated as sources of infection. Likewise, by continuing this

blocking in subsequent cycles of the organism, the majority of cases and sources would be eliminated. Thus, it is only by application of this principle through its cumulative action that the disease will be ultimately eradicated.

What is the most common inanimate object which becomes sputum contaminated by one person and then, without being rendered aseptic, enters the mouth of another where the saliva and its organisms are deposited? The donor of the saliva may be tuberculous and the recipient nontuberculous. This exchange of inanimate objects with their attached organisms takes place three times a day between tuberculous and nontuberculous persons. These objects are eating utensils. Objection might be made to the possibility of any such transfer of tubercle bacilli, on the assumption that only the sputum contains large numbers of organisms, while the saliva of the oral cavity is comparatively free from contamination by this organism; also that eating utensils are washed, and furthermore that such utensils when used by the tuberculous, who have had hospital training, are boiled.

The saliva in the open case contains many tubercle bacilli; dishes when washed only in warm water are not rendered aseptic—the average removal of organisms is about 70 per cent.;² and the tuberculous who have had the advantages of training in a sanatorium are in the minority.

TRANSMISSION EXPERIMENTS

The field and laboratory investigations on the transmission of measles-pneumonia at San Antonio, as well as those on influenza-pneumonia at the port of embarkation, Newport News, Va., and in public institutions presented a lead as to the most likely major avenue of tuberculosis transmission.³ From these investigations it was deduced that direct contact through the air route played at best only a minor rôle in sputum-borne transmission, while indirect contact through hands and eating utensils was the major avenue of distribution.

The plan of the work was: (1) to determine the presence or absence of tubercle bacilli on eating utensils after they were used by tuberculous patients; (2) to determine the presence or absence of these organisms on eating utensils after these utensils were washed by the usual hand method in hot water; (3) to determine their presence on the hands of patients, and (4) to determine their presence in the air of tuberculosis wards.

In carrying out these researches it was recognized, from the standpoint of tuberculosis transmission, that of all eating utensils, the spoon would be the most likely transmitting object. This is because the spoon actually enters the mouth; the fork does, too, but of the two the spoon has the larger surface area to which mouth organisms may adhere. The spoon was, therefore, chosen as the transmitting agent in these experiments.

Spoon Wash Water.—In the first group of experiments, which were undertaken, May 1, 1919, the spoons used in two open cases of tuberculosis were washed after each meal with a cloth in 150 c.c. of hot water. About 50 c.c. of this wash water were centrifugated, and the centrifugate of eleven of these specimens was

2. Observations as yet unpublished.

1. Cumming, J. G., and Spruit, C. B.: Transmission of the Pneumonia Group of Organisms, to be published.

3. Lynch, Charles, and Cumming, J. G.: The Distribution of Influenza by Indirect Contact—Hands and Eating Utensils, *Am. J. Pub. Health* 9: 25 (Jan.) 1919.