its glaciation to any general ice-cap radiating from the Pole. He thought that the ice-sheet was general over the northern part of the British Isles, and on a much larger scale than was usually admitted—and that one of the obstacles to its recognition was the later glaciation along the valleys, which was more conspicuous than the older traces, and another the difficulty which some people had in ignoring the present coast-line.

Mr. J. CLIFTON WARD stated that in the northern parts of the Lake district he had found that the direction of the ice-flow must have been mainly to the north.

The AUTHOR, in reply, remarked that in Greenland, whatever might theoretically be the case, ice is pushed for scores or hundreds of miles down into the sea, until it gets out of its depth, and eventually floats off as icebergs. He pointed out the correspondence of the main valleys of Ireland with glaciations on the surface of rocks from Scotland, and exhibited specimens and rubbings in illustration of various characters of weathering and wear from different natural causes.

FEBRUARY 5, 1873.

Thomas Checkley, Esq., 70 Lichfield Street, Walsall; John Mackenzie, Esq., Government Examiner of Coal-fields for New South Wales, Newcastle, N. S. W.; John Ollenshaw Middleton, Esq., 1 Ebenezer Terrace, Plumstead Common, S.E.; Walter Rowley, Esq., 74 Albion Street, Leeds; and George William Shrubsole, Esq., Victoria Road, Chester, were elected Fellows of the Society.

The following communication was read:—

The OOLITES of NORTHAMPTONSHIRE. By SAMUEL SHARP, Esq.,
F.S.A., F.G.S. Part II.

[PLATES IX. & X.]

INTRODUCTION.

IN accordance with the intimation given in my First Part of a Memoir upon the Oolites of Northamptonshire, I now beg to offer the Second Part of that Memoir.

The limited district of which I treated on the former occasion afforded facilities for its division into four areas, and for describing the beds and the order of their superposition in each; and thus, by a comparison of the beds and their sequence in the several areas, for arriving at a right understanding as to the geology of the whole district.

The field to which I now direct attention is much more extended, and is of a character that will not allow of such a systematic division. I hope, nevertheless, to be able to convey clearly that which I have to communicate.

The main feature of my First Part was the description and con-

sideration of that formation, commercially most valuable, and geologically most interesting, the Northampton Sand.

The main feature of my Second Part will be the description and consideration of a series of beds grouped by Mr. Judd under the name of the "Lincolnshire Limestone"—of less commercial value than the former, it is true, but scarcely of less geological interest.

My endeavour has been to trace, through the county in a north-easterly direction, to Stamford and somewhat beyond, the continuity of beds occurring in the Northampton district; to describe the Oolitic beds and their sequence at certain points within the area considered (involving the interposition of a new and important formation); to examine the geological characteristics of the districts round Stamford, and very briefly those of the eastern portion of the southern border of the county; and thus, while giving a general idea of the geology of the Northern Division of Northamptonshire, to help to establish the soundness of certain views not yet held to be altogether conclusive.

My data were gathered and this Memoir was drafted before I had the advantage of examining Mr. Judd's Geological Survey Map, Sheet 64; while, unfortunately for geological science, but perhaps as well for the object I have in view, that gentleman's Memoir has not yet become the property of the public.

It may perhaps by some be deemed superfluous in me to have produced this treatise at all, seeing that Mr. Judd's map is already in the hands of geologists, and that his Memoir will shortly be published.

But some geologists still hesitate to accept the dictum that the beds of the Lincolnshire Limestone are Inferior Oolite; and, as Mr. Judd, after some years of close and systematic examination of those beds, and I, after a less systematic and exact although longer acquaintance with them, have independently come to the same conviction upon this point, I have thought that my second voice might not be without service, nor my local information without interest—especially as I have come armed from the beds to be discussed with an array of significant fossils, those once animate though now inanimate "oldest inhabitants," those silent yet eloquent witnesses to the truth of the conclusions at which we have arrived.

Before entering upon the task I have essayed to accomplish, as the Northampton district will, as it were, form my starting-point in tracing out the extension eastwards of the Oolitic beds of that district, it will be well to recall to notice the General Section given in my former communication.

It will be seen by this diagram that upon the clay of the Upper Lias are superimposed the series of beds of the Northampton Sand, having a maximum thickness of about 80 feet; which beds were divided by me (because of certain distinguishable characteristics) into "Lower," "Middle," and "Upper;" that upon the summit of the Northampton Sand occurs a plane of unconformity, indicated in the diagram as the "Place of the Great Limestone of the Inferior Oolite" (the Lincolnshire Limestone of Mr. Judd); and that above

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

227

this are three beds of the Great Oolite series, having a maximum thickness of 45 feet—the *lowest* consisting of clay having a persistent

Fig. 1.—Diagram of General Section, showing the position of the Inferior Oolite Limestone and Divisions of Northampton Sand.

ferruginous band at its base (which clay, containing wood, plants, and numerous bands with fresh-water shells, and exhibiting other estuarine characteristics, has been termed by Mr. Judd the “Upper Estuarine Series”); the *middle*, of a series of marly limestones; and the *uppermost*, of clay characterized by an abundance of *Ostrea subrugulosa*.

The Middle Division of the Northampton Sand was separated by me from the Lower Division (provisionally, as I stated) because of the intervention of a band of coarse Oolitic Limestone, and because of differences, throughout the district then treated of, in their mineral and stratigraphical characteristics.

To the north-east of Northampton, however, those differences are not so marked: for instance, over a considerable area, in the Lower Division, for ironstone is substituted an only slightly ferruginous limestone; and again, in other places, the Middle Division is quarried for ironstone, being more richly ferruginous than the Lower Division, while the Limestone band is frequently absent.

In contrast to the varying, and therefore uncertain, marks of distinction between the Lower and Middle Divisions of the North-

ampton Sand, those which define the separation of the Middle from the Upper Division everywhere obtain; for the presence of the sand-bed (generally with its characteristic vertical plant-markings, and designated by Mr. Judd the "Lower Estuarine Series") is observable wherever the Northampton Sand and higher beds are found in the same section.

It is a question, therefore, whether it would not be wise to abandon the hard line of separation between these Middle and Lower Divisions, and to class them together as the "Ferruginous" or marine beds of the Northampton Sand, retaining the distinction of the Upper Division of that formation under Mr. Judd's name of the "Lower Estuarine Series."

Exception has been made to the use of Mr. Judd's terms "Upper Estuarine" and "Lower Estuarine." I shall not enter upon this question: Mr. Judd is well able to maintain his own position. But, as these terms have been adopted in the maps of the Geological Survey, and will doubtless also be used in Mr. Judd's forthcoming Memoir, I have deemed it well, for the avoidance of confusion, to retain them in this treatise. For the same reason, I have adopted Mr. Judd's term of the "Great Oolite Clay" for the clay overlying the Great Oolite Limestone.

It is a fact worthy of notice that the two series of beds, the "Upper Estuarine" and the "Lower Estuarine,"—so widely separated in time and character, the one belonging to the period of the Great Oolite, and the other to that of the Inferior Oolite,—occur together in vertical juxtaposition in the neighbourhood of Northampton, throughout a large district including parts of both divisions of the county, and in Oxfordshire. In the latter county, the Upper Estuarine is traceable through to the Stonesfield Slate-bed; and the difficulty of separating the two Estuarines in Oxfordshire led to the conclusion arrived at, and published by the Geological Survey in 1860, that the Northampton Sand (few of its fossils being then known) was equivalent to the Stonesfield Slate.

An earlier conclusion prevalent among geologists (and perhaps still retained by some) was, that the Great Oolite limestone of the high grounds of the Northampton district, was identical with the limestone (the Lincolnshire Limestone) which occurs between Kettering and Stamford, and, characterizing the country about the latter town, extends on through Rutland and Lincolnshire into Yorkshire; that this Limestone was a member of the Great Oolite series; and that the calcareo-arenaceous slate of Collyweston and Easton, which bases this limestone, was the equivalent of the Stonesfield Slate of Oxfordshire: and this opinion has been maintained notwithstanding the discrepancy indicated by the very dissimilar fossil contents of the two formations and by the anomaly of their relative stratigraphical position.

I hope to be able to show the distinction between these two formations; to demonstrate, from the succession of beds at different points between Northampton and Stamford, and in the districts about Stamford, that, while the position of the Northampton Lime-

stone is that indicated in the diagram of the Northampton section, the true position of the Lincolnshire Limestone is below the Upper Estuarine series, and above the Lower Estuarine series of Mr. Judd's determination; and to prove, by palæontological evidence, that the former belongs to the Great Oolite, and the latter to the Inferior Oolite group of formations.

NORTHAMPTON TO KETTERING*.

In proceeding from Northampton in a north-easterly direction to Kettering (14 miles) *en route* for Stamford, are passed over (variously, according to the frequent alternations of hill and dale) the several beds of the Northampton section. The Lincolnshire Limestone is not encountered in any place on this line, nor indeed is any peculiarity worthy of comment here, excepting that, at a few miles south of Kettering, a wide tract of the Great Oolite limestone (of which Kingsthorpe offers a typical example) is hidden by overlying beds of Glacial Drift. The occurrence of such Drift-beds is very frequent in the south-eastern and southern districts of Northamptonshire; but, as my purpose is to treat only of the Oolitic beds, I shall not again allude to these more recent deposits.

Before passing beyond Kettering to districts in which the great expanse of the Lincolnshire Limestone may be explored, I will revert for a moment to parts of the county, west of that town, in which that formation occurs.

MAIDWELL, HARRINGTON, AND OLD.

At a distance of about 10 miles due north of Northampton, in the lordship of Maidwell, is a small patch of the Lincolnshire Limestone, having a spread of about a mile east and west, and half a mile north and south. This patch ranges about half a mile further west than the town of Northampton, and marks the extreme western extension of the area of the formation.

About a mile and a half due east of this, and separated from it by a north and south Liassic valley (through which the Blisworth and Harborough railway passes), is a much larger patch of this Limestone; which is the uppermost sub-surface rock throughout the greater part of the lordships of Harrington and Draughton.

Immediately beneath the Lincolnshire Limestone of these patches, as shown by the outcrop of the escarpment on either side of the valley, occur the Lower Estuarine and the Ferruginous beds of the Northampton Sand, without the intervention (be it particularly remarked) of the Upper Estuarine Clays, which, as B in the North-

* The route taken by the author in his descriptions and the areal disposition of the several beds he has described can be traced upon the maps of the Geological Survey—quarter-sheet LII. N.W., revised by Mr. Judd, and the admirable map sheet LXIV., by the same gentleman; but that route perhaps may be more easily followed by reference to the accompanying reproduced diagrammatic map (Pl. IX.), which was exhibited by the author on the occasion of the reading of this paper.

ampton Section, there separate the Great Oolite Limestone from those lower beds.

About three miles to the south of the Harrington area, and half a mile due east of the village of Old, is a small narrow strip of the Lincolnshire Limestone, only worthy of mention as being the most southerly point at which the presence of that Limestone has been detected.

KETTERING.

At about a mile due east of Kettering, the eastern escarpment of the valley of the small river Isc presents this section :—

Great Oolite Limestone.	
Upper Estuarine Series.	
Lower Estuarine Series,	} Northampton Sand.
Ferruginous Beds,	
Upper Lias Clay.	

This sequence (with the occasional superaddition of the Great Oolite Clay) is continued due east, right across the county to the Nene valley (a distance of 6 miles), and, with the further superadditions successively of Cornbrash, Kelloway Rock, and Oxford Clay, some three miles further still, to the boundary of Northamptonshire, and so on beyond, into Huntingdonshire.

Upon the outcrop of the same escarpment, at a point near Kettering Mill, and a very little north of the last-mentioned section, we find, as it were, the thin end of the wedge of the Lincolnshire Limestone coming in; and this section, and for the first time, is presented :—

Great Oolite Limestone.	
Upper Estuarine Series.	
LINCOLNSHIRE LIMESTONE (very thin).	
Lower Estuarine Series,	} Northampton Sand.
Ferruginous Beds,	
Upper Lias Clay.	

Upon the summit of the escarpment, upon the opposite or western side of the Isc valley (which here is about three quarters of a mile wide), and at about three quarters of a mile north-east of Kettering, the Lincolnshire Limestone, with a slightly increased thickness, again occurs, and at a little distance to the north is surmounted by the Upper Estuarine beds.

It is clear that at this point we cannot be very far off the original line of the thinning out of the Lincolnshire Limestone, and that we have touched upon the main continent (so to speak) of that formation, from which the Isc valley has here severed a portion of the southern margin; for, as the valley of the Isc trends northwards, the thin band exposed by the escarpment both thickens and spreads.

LEICESTERSHIRE.

From the point of its first appearance near Kettering, the formation extends for six miles in a north-westerly direction towards Leicestershire; in which county is a small outlier between Medbourne and Holt.

GLENDON, COTTINGHAM, ROCKINGHAM, &c.

Diverging from a north-east course, we will follow for a time the range of the formation northwards. At Raven Wood, near Glendon (two miles from Kettering), are ancient quarries in the Lincolnshire Limestone; which here has a thickness of about 8 feet, and is disposed in the same number of courses, some of the upper of which are more or less Oolitic. Towards the bottom is a hard band containing numerous *Nerinea cingenda*, Bronn, *N. triplicata*, Bronn, *Pinna cuneata*, Phil., *Serpula*, &c.

It was these quarries which, in 1869, I had the privilege of visiting (as stated in my First Part) in the company of Professor Ramsay, F.R.S., Mr. Etheridge, F.R.S., Mr. Howell, F.G.S., and Mr. Judd, F.G.S.; and here, upon the very threshold, as it were, of the formation, we found an abundance of fossils of sufficient significance to stamp the character of this Limestone, as being distinct from the Great Oolite Limestone of the Northampton district, and as being as certainly a member of the Inferior Oolite series.

Prominent among the fossils obtained and noted were:—

Gervillia acuta, Sow.
Lima proboscidea, Sow.
 —, large sp. (allied to *L. grandis*, Römer)?
Pecten personatus, Goldf.
Pinna cuneata, Phil.

Cardium Buckmani, Mor. & Lyc.
Ceromya Bajociana, d'Orb.
Lucina despecta, Phil., var. *cardioides*, d'Arch.
Myacites Scarburgensis, Phil., sp.
Pholadomya fidicula, Sow.

Pholadomya Heraulti, Ag.
Tancredia axiniformis, Phil.
Trigonia hemisphaerica, Lycett.

Natica Leckhamptonensis, Lycett.
Nerinea cingenda, Bronn.
 — triplicata, Bronn.

Serpula socialis, Goldf.
 — sp.?

Acrosalenia Lycettii, Wright.
Pygaster semisulcatus, Phil.

Although the Upper Estuarine Clays are not seen in contact with the Lincolnshire Limestone in the Raven-Wood section, they immediately overlie it at a short distance eastwards.

At a short distance to the west is a deep railway-cutting in the Lower Estuarine Sands; and at its north end, on either side, are shallow quarries in the Ferruginous beds of the Northampton Sand. So that the circumstances of the locality offer facilities for ascertaining certainly the vertical sequence of beds there; which is represented in this gathered section, drawn from data taken by Mr. Etheridge on the spot:—

Section near Glendon Railway-cutting.

		ft.	in.
Northampton Sand.	Lincolnshire Limestone	8	0
	Lower Estuarine Sands:—	ft.	in.
	a. Brown Sand	2	0
	b. Plant-bed, with vertical markings	1	6
	c. White Sands	8	0
	—	11	6
	Ferruginous beds	10	0
	Upper Lias Clay.		

The ironstone here is very fossiliferous in bands; the fossils noted presenting the same *facies* as those obtained from the Duston ironstone quarries.

Fossils from the Ferruginous Beds at Glendon.

<i>Avicula Münsteri</i> , Goldf.	<i>Astarte elegans</i> , Sow.
<i>Gervillia acuta</i> , Sow.	<i>Cardium Buckmani</i> , Mor. & Lyc.
— <i>lata</i> , Phil.	<i>Corbicella Bathonica</i> , Mor. & Lyc.
<i>Hinnites abjectus</i> , Phil. sp.	<i>Trigonia compta</i> , Lycett.
— <i>velatus</i> , Goldf. sp.	— <i>Phillipsii</i> , Mor. & Lyc.
Lima, large sp. (allied to <i>L. grandis</i> , Römer)?	— <i>Sharpiana</i> , Lycett.
<i>Pecten personatus</i> , Goldf.	— <i>V-costata</i> , Lycett.
<i>Pinna cuneata</i> , Phil.	<i>Unicardium</i> , sp.?
<i>Pteroperna</i> , sp.?	

Passing northwards, at Barford Bridge, about a mile from Raven Wood, we again cross the meandering valley of the Ise, and again encounter, on the northern escarpment, the same decisive sequence of beds as that near Kettering:—

Great Oolite Limestone.	
Upper Estuarine Series.	
LINCOLNSHIRE LIMESTONE.	
Lower Estuarine Series,	} Northampton Sand.
Ferruginous Beds,	
Upper Lias Clay.	

For two miles further north, we pass, upon the high ground, over the Great Oolite Limestone, and, with a slight incline, down a narrow outcrop of the Upper Estuarine Clays, and again on to the Lincolnshire Limestone. This thickens northwards; so that, at Cottingham, is a section which exposes a thickness (and this not the whole thickness) of 25 feet.

Low down in the section at Cottingham, and in the same position in other sections near Rockingham, is a band containing plants, mostly ferns, among which may be detected fronds in fructification of *Pecopteris polypodioides*, Lindley; which zone also occurs at Collyweston and Easton, near Wansford, and elsewhere, in the same position.

In the Park of Rockingham Castle* (the high ridge of which is nearly 250 feet above, and overlooks to the north, the valley of the river Welland, and commands an extensive view into the counties of Leicester and Rutland), I have noted the following succession of beds:—

	ft.	in.	ft.	in.
Lincolnshire Limestone—				
Coarse oolitic beds	10	0		
Marly beds (only partly exposed).....	5	0		
			15	0
Northampton Sand—				
Lower Estuarine Sands, with argillaceous bands and patches, and vertical plants	13	0		
Ferruginous beds.....	14	0		
			27	0
Upper Lias Clay.				

* At the western extremity of the line of my horizontal section. See Pl. X. fig. 2.

WEEKLY and GEDDINGTON.

The road from Kettering to Stamford skirts the valley of the Ise upon its north-western escarpment; which presents, beyond Weekly, the same succession of beds as before, but which in the interval have been somewhat disturbed by a small fault, having a south-west and north-east direction:—

Great Oolite Limestone.
Upper Estuarine.
LINCOLNSHIRE LIMESTONE.
Lower Estuarine, } Northampton Sand.
Ferruginous Beds, }
Upper Lias Clay.

A little further on, at a short distance south of Geddington, is a quarry in the Lincolnshire Limestone*. The Upper Estuarine Clay is just seen coming in at the top; below this, is a thickness of 15 feet of Lincolnshire Limestone (the lowest bed but one consisting of the hard *Nerinea* zone, as at Glendon); and at bottom, a reddish sand, representing the Lower Estuarine Series.

Section at the South Geddington Quarry.

1. Upper Estuarine Clay (just apparent).	
2. Lincolnshire Limestone—	ft. in.
<i>a.</i> Marly oolite, in thin bands, and much shattered, about ...	6 0
<i>b.</i> Soft marly limestone, containing few shells—burnt for lime	3 0
<i>c.</i> Hard sub-crystalline bed, blue-hearted, in three or four courses, containing numerous <i>Nerinea cingenda</i> , <i>N. triplicata</i> , &c.—the fossils standing out in relief upon the joint surfaces, from the action of water charged with carbonic acid.....	3 0
<i>d.</i> Shaly soft bed—"very rotten"—about.....	3 0
3. Reddish sand—Lower Estuarine.	

The village of Geddington is situated upon the Upper Lias Clay of the Ise valley; and is celebrated for its beautiful Cross, the most elegant of the three crosses which remain of the twelve erected in A.D. 1290, by Edward I., in memory of his Queen, Eleanor of Castile. It was built over a spring of ancient celebrity. Its material is Barnack Rag, and it is nearly as perfect as when erected, never having been restored.

Just north of Geddington, is Rippen's quarry in the Lincolnshire Limestone. The section is of the same character as that in the quarry south of Geddington; but the limestone beds have thickened to about 20 feet.

Fossils found in the two quarries:—

Hinnites abjectus, <i>Phil.</i> sp.	Natica Leckhamptonensis, <i>Lycett.</i>
Pecten personatus, <i>Goldf.</i>	Nerinea cingenda, <i>Bronn.</i>
Pinna cuneata, <i>Phil.</i>	— triplicata, <i>Bronn.</i>
Pteroperna pygmaea, <i>Mor. & Lyc.</i>	Pygaster semisulcatus, <i>Phil.</i> ("petrified mushrooms" of the pitmen).
Modiola Sowerbyana, <i>d'Orb.</i>	Pecopteris polypodioides, <i>Lindley.</i>
Trigonia hemispherica, <i>Lycett.</i>	
Unicardium impressum, <i>Mor. & Lyc.</i>	

* Up to this point, the Geological Survey Map, Quarter-sheet LII., N.W., revised by Mr. Judd, has served to supplement my personal observations.

Ascending the escarpment north of Geddington, we again pass over the same almost inevitable sequence:—

Great Oolite Limestone.	
Upper Estuarine Series.	
LINCOLNSHIRE LIMESTONE.	
Lower Estuarine,	} Northampton Sand.
Ferruginous Beds,	
Upper Lias Clay.	

In the discussion after the reading of my First Part, Professor Morris said he “had found a difficulty in reconciling the phenomena of the eastern and western Oolitic areas, but considered that the key of the arrangement was to be sought in the district between Northampton and Stamford.”

I agreed with Professor Morris in the opinion expressed in the latter part of his remarks; but I would submit that “*the key*” has been abundantly found, in the sections already noted, which display such a remarkably uniform and significant sequence of beds; which sequence occurs over and over again upon unnumbered escarpments throughout the area of the Lincolnshire Limestone.

LITTLE OAKLEY and STANION.

From Geddington, we pass over the Great Oolite Limestone to Little Oakley; between which village and Stanion, a valley is traversed which exhibits the usual series of beds, but complicated by the intricate intersection of several small crossing faults.

WELDON.

About four miles north-east of Geddington, are the Weldon quarries, whence is obtained the widely known Weldon freestone. The extent of the area which has been broken up into “Hills and Holes” (much of it planted over with trees of considerable growth) bears testimony to the antiquity of the excavations, and to the quantity of stone which has been raised. This freestone is a rather coarse-grained Oolite of a ruddy hue, having frequently a glittering fracture. When a disengaged block is struck with a hammer, “it rings like a bell” (to use a quarryman’s expression); a peculiarity which more or less characterizes the good freestones of this formation. It was much used in ancient times in the erection of churches and other prominent buildings throughout a very considerable district.

Mr. Judd obtained a fine series of fossils from the Weldon quarries, a list of which will be published in his Memoir. I only succeeded in noting the following:—

Pecten articulatus, *Schloth.*
Rhynchonella varians, *Schloth.*
Monodonta?
Nerinaea cingenda, *Bronn.*
 — *triplicata*, *Bronn.*

Phasianella elegans, *Mor. & Lyc.*
 — *parvula*, *Mor. & Lyc.*
 — *Pontonis*, *Lycett.*
Acrosalenia (spine).

The following is the section of the quarry at this time being worked ; with quarrymen's terms :—

	ft. in.	ft. in.
1. Baring, consisting of—		
<i>a.</i> Surface Soil,		
<i>b.</i> Boulder-clay,		
<i>c.</i> Upper Estuarine Clay	10	0 to 12 0
2. "Top Fine Stone"—pink oolitic freestone, having numerous comminuted but few perfect shells on the bedding surfaces: extracted in large blocks.....	1	6 to 2 0
3. "Rag"—a coarse red shelly stone, variegated horizontally, but not splitting in the planes of variegation; very soft in the bed, but hardening upon exposure; and very full of small shells. This stone, although good and durable for some purposes, is for the most part rejected, and buried with the infilling	2	0 to 2 6
4. Marly limestone, blue-hearted, and sometimes hard: it would make good lime, but is thrown away.....	2	0 to 2 6
5. "Lower Fine Stone"—also of a pink hue, very oolitic in grain, free in working, cut out in large blocks, and containing a few fossils		2 6
6. "Bottom Fine Stone"—a finer, whiter, and very oolitic freestone; the best stone of the section, and also raised in large blocks.....		2 6
7. Below these beds, as shown by a trial-hole, are several other strata; marly, sometimes shaly, and not worth quarrying; and having an aggregate thickness exceeding		14 0

About half a mile due north of the last quarry, is a shallow lime- and road-stone pit, exposing a section of from 6 to 8 feet of marly limestone, in very thin courses, but yielding no fossils. These are probably the continuation of some of the lower and unworked beds of the freestone quarry.

Weldon is about midway of my horizontal section between the Welland and the Nene valleys (Pl. IX. fig. 3); which has Rockingham at its western and Oundle at its eastern extremity, and to which I shall further allude when I come to describe the succession of beds at the last-named town. Although the Lincolnshire Limestone attains at Weldon to a thickness exceeding 25 feet, its south-western boundary cannot be very far removed; as it is nowhere found in directions due east, south-east, or south, at a greater distance than three miles. Its original margin on this side, however, may have been diminished by denudation prior to the deposition of the overlying beds.

KIRBY OLD SLATE QUARRIES.

Two miles due north of Weldon are the Kirby "Old Slate Quarries." These are not now worked, and no section is exposed. From such portions as I could find *in situ*, it would appear that the upper beds are oolitic, and the lower marly limestone. This arrangement seems very frequent in the Northamptonshire districts of the formation (as long since noticed by Professor Morris), although by no means uniformly prevalent; and it is probably merely the result of varying accidental and local conditions.

The marly beds contained:—

<i>Gervillia acuta</i> , Sow.	<i>Modiola Sowerbyana</i> , d' Orb.
<i>Pecten</i> , large (probably lens, Sow., or new sp.?)	<i>Pholadomya</i> , sp.?
<i>Ceromya Bajociana</i> , d' Orb.?	<i>Trigonia Phillipsii</i> , Mor. & Lyc.
— similis, <i>Lycett</i> ?	

The hard *Nerinea* zone was present, but I could not detect the presence of slate. At about half a mile south-west, however, near to Deene Lodge, are more modern pits, where slate is quarried; and these present the extreme south-western point at which slate equivalent to the Collyweston Slate has been found.

DEENE.

At Deene, two miles and a half north-east of Weldon, is a large stone- and brick-pit; which exhibits, under four feet of sand, soil, and rubble, about six feet of the Lincolnshire Limestone, having at its base three feet of a bed with "potlids," representing the Collyweston Slate-bed: below this, the Lower Estuarine occurs, of considerable thickness and very different in character from the same bed in the Northampton district—reddish sand, four feet; grey sand, with vertical plant-markings, three feet; a very dark chocolate almost black shaly clay, in very thin layers, five or six feet; beneath this, slate-coloured clay, containing a zone of iron-pyrites: then several feet of the ferruginous beds of the Northampton Sand; and below all, the Upper Lias Clay.

Upon my first examination of this pit, I was uncertain as to whether the Ferruginous Beds of the Northampton Sand were not wanting here, and as to whether the clays below the vertical plant-bed (as they yielded no fossils as an indication) were Lower Estuarine or Upper Lias. Mr. Judd has told me that he encountered the same difficulty in the first instance; but that, fortunately, a well was sunk, which passed through these clays for 10 feet, pierced several feet in thickness of the Ferruginous Beds of the Northampton Sand, and dipped into the Upper Lias Clay below.

Detailed Section at Deene Brick-pit.

		ft. in.	ft. in.
Lincolnshire Limestone.	1. Baring—soil, sand, and rubbly stone		4 0
	2. Marly limestone, of various degrees of hardness, in courses about 6 inches in thickness		1 6
	3. Sandy bed, containing rounded calcareous concretions	1 6 to 2 0	
	4. Hard subcrystalline limestone, sometimes blue-hearted, containing numerous <i>Nerinea</i>	2 6 to 3 0	
	5. Sand, with "potlids," the latter very hard—Collyweston Slate	3 0 to 3 6	
Northampton Sand.	6. Reddish Sand		4 0
	7. Grey Sands and Clay, with vertical plant-markings		3 0
	8. Dark chocolate, very shaly, thinly laminated clay, about		6 0
	9. Slate-coloured Clay, with masses of iron-pyrites, about		4 0
	10. Ferruginous Beds (several feet).		
	11. Upper Lias Clay.		

Fossils noted at Deene.

Cardium Buckmani, <i>Mor. & Lyc.</i>	Trigonia Sharpiana (?), <i>Lycett.</i>
Ceromya Bajociana, <i>d' Orb.</i>	Nerinaa cingenda, <i>Bronn.</i>
Goniomya V-scripta, <i>Sow.</i>	— triplicata, <i>Bronn.</i>
Lucina Wrightii, <i>Oppel.</i>	Serpula socialis, <i>Goldf.</i>
Modiola Sowerbyana, <i>d' Orb.</i>	Coral, sp. ?
Perna, sp. ?	Coniferous wood (large pieces of drift).
Pholadomya sp. ?	Ferns (Pecopteris polypodioides,
Trigonia Phillipsii, <i>Mor. & Lyc.</i>	<i>Lindley</i>), &c.

DEENE TO COLLYWESTON.

As between Northampton and Kettering the road passes over the Great Oolite Limestone and subjacent beds, varied by the occurrence and recurrence of hill and dale, so between Deene and Collyweston, passing through the villages of Bulwick and Duddington, the road, with a like variation, traverses the Lincolnshire Limestone for a distance of seven miles. No peculiarity worthy of notice occurs on this line; and I will, therefore, diverge, at Fineshade, for the sake of noticing a rather remarkable section at Wakerley, one mile to the west, and quarries, notable for the fossils they have yielded, at Morcot, two miles beyond.

WAKERLEY.

The section at Wakerley exposes a series of beds in the Lincolnshire Limestone, having an aggregate thickness of about 30 feet; the whole of which, from the top to the bottom, are exceedingly (some coarsely) oolitic; and several exhibit false-bedding. Five or six courses of the lower part of the section are like the "Rag" at Weldon, and full of imperfectly preserved shells. I obtained from the "Rag" beds, *Pseudodiadema depressa*, Ag., and a fine spine of *Cidaris Fowleri*, Wright*.

Details of Section of Lincolnshire Limestone at Wakerley.

	ft.	in.	ft.	in.
1. Rubbly oolite	3	0 to 3	6	6
2. Oolitic limestone, in three courses	3	6 to 4	0	0
3. The like, false-bedded, very inclined			4	0
4. The like, but in horizontal layers.....	3	0 to 3	6	6
5. The like, false-bedded			4	0
6. The like, horizontal			3	0
7. Very coarse oolite			1	0
8. "Rag" bed (like the "Rag" at Weldon), full of shells, chiefly minute, imperfectly preserved—in four to six courses, some very false-bedded	6	0 to 8	0	0

Between the nearly adjoining villages of Wakerley and Barrowden, flows, over the Upper Lias, the river Welland; which, throughout almost the whole of the southern boundary of Rutland, serves as the line of demarcation between that county and Northamptonshire.

* Mr. Judd, I believe, has obtained a large number of characteristic Inferior Oolite fossils from this quarry.

MORCOT.

At Morcot, in Rutland, two quarries are now being worked: these are at different elevations upon the escarpment (facing north-west) of the small river Chater. It is probable that the lower beds of the upper quarry are identical with the upper beds of the lower quarry. It will be necessary, therefore, in order to understand the whole section, to combine the two; and this view is confirmed by the fact that, while beds having blue cores are absent in the upper quarry, such beds occur towards the bottom of the section in the lower one.

The total thickness of the beds at the higher level is about 18 feet, and that of the beds at the lower level about 17 feet: but these figures do not indicate the whole thickness of the Lincolnshire Limestone at Morcot; for the presence of beds of the same formation is to be detected still higher on the escarpment than the summit of the higher quarry, and below the lowest beds of the lower quarry occur slate beds (not now exposed) from which characteristic fossils have been obtained: so that it is probable that the Lincolnshire Limestone at this place has a thickness of quite 60 feet.

The Morcot sections display the local variability of the Lincolnshire Limestone; for, while at Wakerley, only two miles distant, the beds from top to bottom are essentially oolitic, at Morcot, the sections are marked by the almost entire absence of the oolitic character, there being only one bed, of a foot in thickness, which is oolitic at all, notwithstanding that the numerous beds differ greatly in other respects, consisting of harder and softer bands of marly limestone in various alternations.

DETAILED SECTIONS.—Upper Quarry, Morcot.

	ft.	in.	ft.	in.
1. "Rammel"—broken limestone	3	0 to 3	6	
2. Soft calcareous bed, used for burning only.....			3	0
3. Brownish sand, containing calcareous masses—"used for building"			3	6
4. Compact limestone.....			1	0
5. Oolitic "Wallstone".....			1	0
6. Hard compact limestone, in two courses.....			1	0
7. Close marly limestone, sometimes rather crystalline.....				9
8. Hard marly limestone, thickness uncertain.				

Lower Quarry, Morcot.

	ft.	in.	ft.	in.
1. "Rammel"—broken limestone			3	0
2. Shattered limestone, in thin layers, becoming less broken and thicker in the layers towards the bottom of the bed				6 0
3. Soft shaly bed, with plants	2	0 to 3		0
4. Compact marly limestone, in two courses—with <i>Hinnites abjectus</i> , Phil.				2 0
5. Very hard blue-hearted limestone			9 to 1	0
6. Hard limestone, more or less crystalline, sometimes blue-hearted, in two courses			3	0

Fossils from Lincolnshire Limestone, Morcot.

Gervillia radians, *Mor. & Lyc.**
Hinnites abjectus, *Phil.*
Inoceramus obliquus, *Mor. & Lyc.*
Lima cardiiformis, *Sow.*
Ostrea (young).
Pecten lens, *Sow.*
Pinna cancellata, *Bean.*
 — *cuneata*, *Phil.*

Ceromya Bajociana, *d' Orb.*
 — *concentrica*, *Sow.*
 —, sp.?
Lucina Bellona, *d' Orb.*
 — *Wrightii*, *Oppel.*
Modiola Sowerbyana, *d' Orb.*
 — *sublævis* (?), *Sow.*
Pholadomya Dewalquea, *Lycett* (syn.
 of *media* of *Dewalque* and *Chapuis*).
 — *ovulum*, *Ag.*
 —, sp.?

Tellina, sp.?
Trigonia compta, *Lycett.*
Unicardium depressum, *Phil.*, sp.

Terebratula globata, *Sow.*

Natica adducta, *Phil.*
 — *Leckhamptonensis*, *Lycett.*

—, sp.?
Pterocera Bentleyi, *Mor. & Lyc.†*

Serpula, sp.?

Acrosalenia Lycettii, *Wright.*
Cidaris Fowleri, *Wright.*
Pygaster semisulcatus, *Phil.*
Stomechinus germinans, *Phil.*

Ferns:—*Pecopteris polypodioides*,
Lindley, &c.

To the south-east of Morcot, and to a high level, stretches the comparatively barren tract of Barrowden Field; in crossing which by the road to Tixover and Duddington, are passed over in succession (above the Lincolnshire Limestone)—the Upper Estuarine Series, the Great Oolite Limestone, the Great Oolite Clay, and, at the highest ground, a thin capping of CORNBRAsh.

LUFFENHAM.

Two miles north-east of Morcot, upon the Uppingham and Stamford road, near to the Luffenham Railway Station, is a quarry the limestone beds of which answer to those of the lower Morcot quarry, and yield some similar fossils.

KETTON.

Three miles further in the same direction, and beyond the village of Ketton, are the quarries whence is obtained the far-famed KETTON FREESTONE. This is the most excellent in quality of any freestone now obtainable throughout the entire area of the Lincolnshire Limestone.

When I refer to the church of St. Dunstan in Fleet-street, which is built of this stone, and point to the unimpaired condition of its exterior surfaces, after many years' exposure to the disintegrating action of the London atmosphere, I need say nothing of the durability of the Ketton freestone, even when exposed to unfavourable conditions.

It is a matter of observation that the "Ketton" freestone is of better quality when obtained from beneath a thick bed of clay, than

* Nearly allied to *Gervillia Hartmanni*, Goldf., a not unfrequent fossil in the Duston ironstone beds.

† Collected by Mr. Judd. See notice of this beautiful form in description of Fossils of Collyweston Slate Beds—forward, page 247.

when occurring nearer to the surface; and it is perhaps worthy of notice that, not only Ketton stone, but all stone having its origin in sedimentary deposition, is more enduring when in buildings it is placed in the same horizontal position which it occupied in its natural bed. If stone is placed in buildings with its planes of stratification in a vertical position, the crush of superincumbent weight will frequently split it; and this not occurring, weather action will probably scale the surface. Such phases of dilapidation are constantly to be observed in stone window mullions; in which, for the sake of economy of time and material, the naturally horizontal stratigraphical planes of the stone are perverted into a vertical position.

The Ketton quarries exhibit an immense area of broken ground; and, as no attempt has been made to level or restore the surface after excavation, it is left in a perfectly useless condition—offering a very unfavourable contrast to the admirable practice which obtains in some parts of Northamptonshire, of restoring, after quarrying, the surface to a state as good as, and sometimes, for the purposes of cultivation, even better than, before.

There are two places, not widely separated, where the Ketton stone is quarried at this time—one an old working, called “the Deeps,” from which comparatively only a small quantity of stone is obtained, but presenting the deeper section; the other, more recently opened, presenting a less deep but very lengthened section. The latter is divided into several holdings or proprietorships; a large number of men are employed; and a great quantity of stone is raised.

In the “Deeps,” the freestone underlies about 17 feet of the Upper Estuarine Clays, divided into three beds of various character, colour, and thickness, based by the usual ferruginous band. These clays contain shelly seams; some of which are characterized by the presence of the small peculiar rostrated bivalve *Nœra Ibbetsoni*, Morris, and others by *Cyrena*, &c. Three beds of stone are exposed, having an aggregate thickness of 11 feet: these consist of—a coarse red Oolite, good Ketton freestone, and a very hard, sometimes crystalline and blue-hearted, “Rag” stone.

Details of Section at “The Deeps,” Ketton.

1. Upper Estuarine beds—

	ft. in.	ft. in.
a. Grey clay	3 0	
b. White clay	5 0	
c. Variegated clay	8 0	
d. Ferruginous band..	0 9	
	—	16 9

2. Oolitic beds, Lincolnshire Limestone—

a. Coarse red oolite	2 6	
b. “Freestone”—good Ketton Stone, <i>Lima proboscidea</i> , frequent	3 6	
c. “Rag” stone—sometimes blue-hearted, very glistening on fracture, in several courses	5 0	
	—	11 0

Many years ago, in an adjacent working, I obtained during one of my pleasantly remembered excursions in the company of Prof. Morris, a fragment of the upper surface of the red oolitic bed of this section, which had been profusely perforated by a small boring bivalve.

In the Ferruginous beds of the Northampton Sand, casts of the crypts of *Lithodomus inclusus*, Phil., generally including the shell, are very abundant, and sometimes large, but are always found associated with masses of coral. On the other hand, in the limestones of the Great Oolite, casts of the crypts of *Pholas* occur, not so associated. As no coral has been found in connexion with these borings, I am disposed to conclude that they are the work of *Pholas* rather than of *Lithodomus*. At any rate, I would agree with the conclusion arrived at by Prof. Morris, as stated in his admirable article in the 'Geological Magazine' for March, 1869, that these perforations "clearly indicate a period of arrested deposition in this old sea-bed; during which these mollusks lived and inhabited the surface of the subjacent rock, already partly consolidated."

The newer and larger excavations at Ketton are near the base of an escarpment facing north-east. The section presents a thickness of beds of some 20 to 23 feet. The upper portion, to the depth of about 15 feet, consists of the clays of the Upper Estuarine Series, in three beds (as in the last section), viz. grey clay, whitish clay (with vertical yellow markings, as of root perforations), and a very compressed, thinly-laminated, and somewhat bituminous clay, intercalated with thin seams of brown and white sand, and having the ferruginous band at the base. The lower portion consists of Lincolnshire Limestone; the upper two beds of which, having together a thickness of from 5 to 6 feet, are oolitic freestone. The upper bed, being red, coarse, and perishable, and known as "Flesh" or "Crash" by the quarrymen, is rejected: the lower bed is the celebrated Ketton freestone. Beneath the freestone, is a shelly "Rag" bed, not worked, but probably about 4 feet in thickness; and underlying this, is a harder oolitic limestone having a glistening fracture, which also is not worked, and its thickness is uncertain. I have obtained from this bed, *Cypriocardia Bathonica*, d'Orb., and *Pholadomya fidicula*, Sow.

As the works proceed, and an advance is made into the mass of the hill, the section will deepen by an accession to the thickness of the Upper Estuarine clays. The high ground above has a capping of Great Oolite limestone.

Details of Section at the Newer Excavations, Ketton.

1. Upper Estuarine Series—

	ft. in.	ft. in.
a. Grey clay	3	6
b. Whitish clay, with yellow vertical markings, like root perforations	4	6
c. Very compressed and somewhat bituminous clay, in very thin layers, intercalated with thin seams		

	ft. in.	ft. in.
of brown and white sand, and containing shelly bands	7 0	
d. Ferruginous band.....	0 6	15 6
2. Lincolnshire Limestone—		
a. Coarse and very oolitic red stone, sometimes perforated on the upper bedding surface by <i>Pholas</i> borings—the “Flesh” or “Crash” of the quarrymen—crumbles upon exposure to frost, and is not quarried for “freestone”—varying in thickness up to	2 6	
b. “Freestone” bed—the celebrated Ketton stone, very oolitic, and freely worked, hardens upon exposure	3 0	
c. “Rag” stone—pisolitic bed, about	4 0	
d. Harder “Freestone”—more siliceous, presenting a glistening fracture, not worked, and thickness not ascertained	—	9 6
	—	

Fossils from the Upper Estuarine Series, Ketton.

Modiola imbricata, Sow.
Næra Ibbetsoni, Morris.

Cyrena, sp.?
&c.

Fossils from the Lincolnshire Limestone Beds, Ketton.

Lima bellula, Mor. & Lyc.
— *cardiiformis*, Sow.
— *proboscidea*, Sow.
— *Pontonis*, Lycett.
—, large sp. (allied to *L. grandis*, Römer)?
Pecten aratus, Waagen.
—, sp.?
—, sp.?
Pteroperna plana, Mor. & Lyc.
Cardium Buckmani, Mor. & Lyc.
Ceromya, sp.?
Lucina Wrightii, Oppel.
Modiola imbricata, Sow.
Myacites decurtata, Phil.
—, sp.?
Pholadomya fidicula, Sow.
Pholas, sp.?

— *Terebratula fimbria*, Sow.*
— *globata*, Sow.
— *sub-maxillata*, Sow.
— *perovalis*, Sow.
Natica Leckhamptonensis, Lycett.
Belemnites Bessinus (?), d'Orb.
— *Acrosalenia Lycettii*, Wright.
Echinobrissus clunicularis, Lhuys.
Pseudodiadema depressum, Agassiz.
Montlivaltia Delabechii, Edw. & Haime.
— *Strophodus magnus*, Ag. (palates).

KETTON, TO GEESTON AND COLLYWESTON.

Proceeding southwards, through Ketton, we pass its very beautiful church, portions of which are of the Norman, Semi-Norman, Early English, and Early Decorated styles (dating respectively in the 12th, 13th, and 14th centuries), all unimpaired, and all built of the neighbouring “Barnack Rag,” the famous church building-stone of a wide district in those early times.

We then cross upon the Upper Lias the river Chater, and ascend to the so-called Ketton station, which is in the hamlet of Geeston.

* *Terebratula fimbria* occurs also at Denton, Ponton, and Barnack, and Mr. Judd has met with it in the same beds near Lincoln.

A deep railway-cutting here trenches the Lincolnshire Limestone; the beds of which are much inclined and disrupted, having been traversed by a fault. The section exhibits no peculiarity: the upper beds, as usual, are oolitic, the lower more or less marly and crystalline. The Slate bed occurs near the bottom, overlying the Lower Estuarine and the Ferruginous beds of the Northampton Sand. Prof. Morris, in his "Notice," published in the 'Transactions of the British Association' for 1847, describes the section in this cutting as exposed during the formation of the railway, as follows:—

	ft.	in.
"Rubbly oolite in shivers	3	0
Compact marly limestone— <i>Nerinea</i> and <i>Ferns</i>	2	6
Marly rock, very fossiliferous— <i>Nerinea</i> , <i>Modiola plicata</i> , <i>Ferns</i> , and <i>Isocardia (Ceromya) concentrica</i> , <i>Pinna</i> , <i>Arca</i> ...	2	0
Sandy rock, with <i>Lima</i> , &c.	2	0
Crystalline ragstone, with <i>Nerinea</i> and patches of plants.....	3	0
Compact crystalline oolitic ragstone	8	0
Concretionary bed.....	2	6
Slate beds	3	0
Greenish clay.....	2	0
Ferruginous sand of inferior oolite at bottom."		

Proceeding southwards, we cross the Welland valley (again upon the Upper Lias) once more into Northamptonshire; and, in ascending the hill to Collyweston, see cropping out in succession upon the excavated road-side the Ferruginous beds of the Northampton Sand, the Lower Estuarine Sands, the Collyweston Slate, and some Limestone bands above.

COLLYWESTON.

The Collyweston Slate quarries commence immediately east of the village, and extend continuously for more than a mile north-east into the parish of Easton. It is known that they have been worked for 350 years, and possibly for a much longer period. The slate is of a fine-grained calcareo-arenaceous material, and occurs at the base of the Lincolnshire Limestone, in a thin band; the under surface of which frequently assumes the form of flattened hemispherical masses (like the "potlids" and "whinstones" of Stonesfield), which with the convex surfaces downwards, repose upon the white sand of the Lower Estuarine Series: in hollows upon the surface of which it would seem as if they had been moulded.

The "Slate," when first raised, is a very hard and solid stone, frequently blue-hearted; but, upon exposure to moisture and to frost, it becomes fissile, and its fine laminæ readily split into the "slates" so widely known and locally used for so long a period. It is remarkable, however, that, after exposure, if the frost has not been of sufficient intensity to split the stone, it becomes impervious to frost-action afterwards. I was informed by the pitmen during a recent visit, that only one night's effective frost occurred during the previous winter (1871-72); that, in consequence, an immense quantity of slate stone which had been raised was not available for

conversion into slates; and that as a result many pits for the present had been thrown out of work.

The superincumbent beds at Collyweston and Easton have an aggregate thickness varying according to situation from 15 to 25 feet, and present lithological and sectional peculiarities not to be found in any sections in the Lincolnshire Limestone which I have elsewhere had the opportunity of examining; but which (as Mr. Judd informs me) are not very unlike those which characterize some sections in Lincolnshire.

From the number and great extent of the slate quarries, it would be expected that some variations in the sections would be exhibited. This, to a certain extent, is the case—as will be found upon a comparison, one with another, of a section taken by Prof. Morris prior to 1847, and published in his “Notice” in the ‘Transactions of the British Association’ for that year, one noted by me several years ago, and others taken by me during the recent autumn. Beds present in some sections are wanting or have changed their character in others; and beds traceable through several sections are thicker or thinner according to locality.

Section taken at Collyweston by Professor Morris prior to 1847.

	ft.	in.
1. Compact oolite in shivers.....	7	0
2. Oolitic freestone	2	6
3. Marly oolite	2	6
“ Cream-coloured marly limestone, containing zones of shells of which the <i>Nerinea cingenda</i> and other species, <i>Lucina</i> , <i>Pholadomya</i> , <i>Modiola plicata</i> , are abundant; with patches of fragments of ferns (<i>Pecopteris poly-podioides</i>) and other plants (<i>Zamites</i>), not occurring in regular layers.”		
4. Concretionary sand bed (“beds of sand and sand-rock”) ...	1	6
5. Compact crystalline ragstone (“with <i>Nerinea</i> and <i>Ferns</i> ”)...	1	6
6. Brown rubbly incoherent sand, with concretions.....	2	0
“ A concretionary bed of sand, with irregular cylindrical ramose bodies of sandstone, resembling furoid stems.”		
7. Slate beds—“Calcareo-siliceous beds, grey and brown.”		

Former Section taken by S. S.

	ft.	in.	ft.	in.
1. Flaggy limestone, slightly ferruginous, and more arenaceous and siliceous than the ordinary Lincolnshire Limestone of the district. [This bed represents bed 3 of Professor Morris's section.]...	3	0	to	4 0
2. Sand bed—somewhat ferruginous, and containing fantastic concretionary masses. [4 of Morris.]...	3	0	to	4 0
3. Flaggy limestone—sometimes blue-hearted. [5 of Morris.]	3	0	to	4 0
4. Peculiar laminated bed, made up of alternate layers of sand and calcareous sandstone, each about half an inch in thickness; these are from 40 to 50 in number, and the edges of the sandstone layers stand out in relief upon the surface of the bed. In one section this bed presents the most remarkable wavy distortion. [6 of Morris?]	2	0	to	2 6
5. Hard compact blue-hearted limestone in courses ...	3	0	to	5 0

6. Slate bed—the lower surface moulded into flattened hemispherical forms. [These beds (5 & 6) represent the slate beds (7) of Morris.]
7. Sand (Lower Estuarine)
8. Ferruginous beds

} Northampton Sand.

During my recent visit, I carefully took notes of two sections somewhat widely separated, adopting for the several beds the familiar names given them by the quarrymen; and I will here parenthetically suggest that it would be desirable generally to adopt the plan of taking such names, as these would serve for the purposes of identification and comparison upon future visits of geologists, and in after times, even though the exact sections described may have long since disappeared. These two sections so nearly tallied in every respect, except as to thickness, that one table will serve to describe both.

Section of Slate Quarries, Collyweston, Sept. 1872, with Quarrymen's Terms.

	Perkins's Pit.		Hill's Pit.	
	ft.	in.	ft.	in.
1. Broken stone, in thin layers (<i>Natica Leckhamptonensis</i> , Lycett)	1	0 to 2	0	2 0 to 3 0
2. "Red Cale"—very oolitic stone, like bad Ketton freestone, in thin courses. [These beds together represent beds 1 & 2 of Professor Morris's section.]		6 0	6 0 to 7 0	
3. "White Cale"—compact marly limestone (<i>Pecopteris polypodioides</i> , in densely filled patches and with fronds in fructification, <i>Hinnites velatus</i> , <i>Lima</i> , sp., large <i>Pecten aratus</i> , <i>Pecten lens</i> , <i>Cypricardia nuculiformis</i> , <i>Unicardium impressum</i> , <i>Nerinea cingenda</i> , &c.). [Bed 3 of Morris, 1 of my former section.]	1	0 to 1	6	1 6 to 2 0
4. Sandy soft bed. [4 of Morris, 2 of my former section.]		0 9		1 0
5. "Ringstone"—large oolitic grains in a very hard crystalline matrix (<i>Modiola Binfieldi</i> ?). [5 of Morris, 3 of my former section.]		0 6		0 8
6. "Top Sand"—the peculiar laminated bed of my former section. The layers are somewhat inclined in Perkins's pit, but more so in Hill's pit, and very distorted in others. [6 of Morris?, 4 of my former section.]		2 0	2 0 to 2 6	
7. "Brood"—a hard oolitic limestone, in courses of about 6 inches thickness, a good building-stone (<i>Natica Leckhamptonensis</i> , large new <i>Lima</i> *). [Wanting in Morris's section.]		3 0		1 6
8. "The Hard Sand"—a variable bed, sometimes much thicker (large <i>Natica Leckhamptonensis</i> , <i>Cardium Buckmani</i> , <i>Ostrea</i>). [6 of Morris? This and the last beds are wanting in my former section.]		0 9		1 0

* Allied to *L. grandis*, Römer?

	ft. in.	ft. in.	ft. in.	ft. in.
9. "The Hard Limestone"—in three or four courses, glistening in the fracture from the presence of small crystals of carbonate of lime, frequently blue-hearted, and a "first-rate building-stone" (<i>Avicula Münsteri</i> , <i>Gervillia acuta</i> , <i>Hinnites velatus</i> , <i>Lima</i> , sp., <i>Pecten aratus</i> , <i>P.</i> , small sp., <i>Pteroperna pygmæa</i> , <i>Cardium Buckmani</i> , <i>Ceromya concentrica</i> , <i>Lucina Wrightii</i> , <i>Natica Leckhamptonensis</i> , <i>Nerinea</i> , sp., crustacean, <i>Pseudophyllia</i> , sp., <i>Ferns</i> , <i>Pecopteris polypodioides</i> , wood, &c. A zone filled with <i>Annelides</i> traverses this bed	2	0 to 3	0	2 0
10. "Bitch"—a soft marly material, good for lime-burning, but not for building-stone (<i>Pecten aratus</i>). [This and the last beds represent 5 of my former section.]		0	0	0 6 0 9
11. "Slates." [These beds, 9, 10, 11, would seem all to be included in slate-bed 7 of Morris.]	2	0 to 2	6	1 6 to 2 6
12. "Sand"—Lower Estuarine with plants.				

In all the sections, some of the beds are divided by thin argillaceous seams.

Fossils from the Beds above the Slates, Collyweston.

<i>Avicula Münsteri</i> , Goldf.	<i>Unicardium impressum</i> , Mor. & Lyc.
<i>Gervillia acuta</i> , Sow.	
<i>Hinnites velatus</i> , Goldf. sp.	<i>Natica Leckhamptonensis</i> , Lycett.
<i>Lima</i> , large new sp. (allied to <i>L. grandis</i> , Römer)?	<i>Nerinea cingenda</i> , Bronn.
—, sp. ?	—, sp. ?
<i>Ostrea</i> , sp. ?	
<i>Pecten aratus</i> , Waagen.	<i>Dentalium</i> (?), new species.
— lens, Sow.	
<i>Pteroperna pygmæa</i> , Dunker, sp.	<i>Serpula</i> , sp. ?
<i>Cardium Buckmani</i> , Mor. & Lyc.	<i>Crustacean</i> —
<i>Ceromya concentrica</i> , Sow.	<i>Pseudophyllia</i> , sp. ?
<i>Cypriocardia nuculiformis</i> , Römer, sp.	
<i>Lucina Wrightii</i> , Oppel.	<i>Ferns</i> —
<i>Modiola Binfieldi</i> (?), Mor. & Lyc.	<i>Pecopteris polypodioides</i> , Lindley.
	Wood.

Connected with the organic contents of the upper beds, the abundance of *Pecopteris polypodioides*, sometimes showing the delicate fructification beautifully preserved, is significant, as it seems to imply that the land surface upon which the *Pecopteris* grew could not have been very distant. The presence of the characteristic Inferior Oolite fossil *Natica Leckhamptonensis* throughout the section, the large new *Lima* with the shell beautifully preserved, and the small new *Dentalium* (?), are also worthy of particular notice.

Among the numerous fossils of the Slate beds, the little rare shell *Alaria Phillipsii* (first found, I believe, in the "Grey Limestone," Inferior Oolite, of Scarborough) claims attention; but the beautiful gasteropod *Pterocera Bentleyi* (named after my friend and former fellow worker, Mr. Bentley, of Stamford) seems chiefly to characterize these beds at Collyweston. The latter is by no means rare,

although the known examples have mostly been found in one quarry (Mr. Hill's); I think, however, that this is chiefly attributable to the fact that the intelligent quarryman, Mr. Lomax, preserves and dispenses to geologists all the fossils which he obtains. I have one good specimen of this peculiarly local fossil, found in the Lincolnshire Limestone at Denton, near Grantham (to which I shall presently allude); it occurs also in the shelly beds at Ponton, and I have already mentioned one other obtained by Mr. Judd from Morcot: but I believe it has not been detected in any other locality. On some of the slates, peculiar large worm-tracks are found.

Fossils from the Slate-beds, Collyweston.

<i>Avicula Braamburiensis</i> (?), <i>Sow.</i> — <i>clathrata</i>, <i>Lycett.</i> — <i>Münsteri</i>, <i>Goldf.</i> — <i>subcostata</i>, <i>Römer.</i> —, sp. ? <i>Gervillia acuta</i>, <i>Sow.</i> — <i>radians</i> (?), <i>Mor. & Lyc.</i> (young). <i>Hinnites abjectus</i>, <i>Phil.</i> sp. — <i>tegulatus</i>, <i>Mor. & Lyc.</i> — <i>velatus</i>, <i>Goldf.</i> sp. <i>Lima cardiiformis</i>, <i>Sow.</i> sp. — <i>gibbosa</i>, <i>Sow.</i> sp. —, sp. ? <i>Pecten clathratus</i>, <i>Römer.</i> — <i>lens</i>, <i>Sow.</i> — <i>personatus</i>, <i>Goldf.</i> <i>Pinna cancellata</i>, <i>Bean.</i> — <i>cuneata</i>, <i>Phil.</i> <i>Placunopsis socialis</i>, <i>Mor. & Lyc.</i> <i>Pteroperna costatula</i>, <i>Deslongch.</i> — <i>pygmæa</i>, <i>Dunker</i>, sp. <i>Astarte elegans</i>, <i>Sow.</i> — <i>excavata</i>, <i>Sow.</i> — <i>depressa</i> (?), <i>Goldf.</i> — <i>compressiuscula</i> (?), <i>Mor. & Lyc.</i> <i>Cardium Buckmani</i>, <i>Mor. & Lyc.</i> — <i>Stricklandi</i>, <i>Mor. & Lyc.</i> — <i>subtrigonum</i>, <i>Mor. & Lyc.</i> <i>Ceromya concentrica</i>, <i>Sow.</i> —, sp. ? <i>Cucullæa cancellata</i>, <i>Phil.</i> — <i>Goldfussii</i>, <i>Römer.</i> —, sp. ?	<i>Goniomya literata</i>, <i>Sow.</i> sp. <i>Homomya</i> (<i>Myacites</i>) <i>unioniformis</i>, <i>Mor. & Lyc.</i> —, sp. ? <i>Lucina Wrightii</i>, <i>Oppel.</i> —, sp. ? <i>Modiola gibbosa</i>, <i>Sow.</i> — <i>Sowerbyana</i>, <i>d'Orb.</i> <i>Myacites decurtata</i>, <i>Phil.</i> sp. <i>Trigonia costata</i>, var. <i>pullus</i>, <i>Sow.</i> — <i>compta</i>, <i>Lycett.</i> — <i>spinulosa</i> (?), <i>Young & Bird.</i> —, sp. ? <i>Unicardium impressum</i>, <i>Mor. & Lyc.</i> —, sp. ? <i>Alaria Phillipsii</i>, <i>d'Orb.</i>, sp. <i>Cylindrites</i>, new species, like <i>turriculatus</i> of <i>Lycett</i> (found at Ponton), but having fewer whorls. <i>Delphinula</i>, sp. ? (see <i>Phil. Geol. Yorks.</i> i. tab. ix. f. 32). <i>Natica</i> (<i>Euspira</i>) <i>canaliculata</i>, <i>Lycett.</i> —, sp. ? <i>Pterocera Bentleyi</i>, <i>Mor. & Lyc.</i> <i>Astropecten Cotteswoldia</i>, var. <i>Stamfordensis</i>, <i>Wright</i> (from Slate-bed, St. Martin's, Stamford). Fish—small teeth and scales. Plants—small fragments.

The Collyweston Slate is now very much in request for roofing either restored or new buildings in Mediæval styles. It assorts well with such buildings; and, while fresh, is very eyeable. When the material and the workmanship are both of the best, it is tolerably durable; but if the slates are not picked, and if the workmanship is not better than ordinary, the roofs soon become dilapidated, ugly inequalities appear, the wind and the rain penetrate, and a slate roof then is neither sightly, efficient, nor lasting.

Sir Gilbert Scott has used this slate in his exquisite restoration of the beautiful Round Church at Northampton, and also for the roof

of the new Chapel of St. John's College, Cambridge; from the chip-pings at which place, through the kindness of Mr. Keeping of the Woodwardian Museum, I obtained a fine specimen of the *Pterocera Bentleyi*.

EASTON.

The Ferruginous beds of the Northampton Sand, underlying the Lower Estuarine, thicken in Easton parish, and may be seen in some trial-holes, north of the village, on the hillside looking towards Tinwell. They are here some 15 feet in thickness, and consist of cellular ironstone of a rich character.

WOTHORPE TO ST. MARTIN'S.

Just east of Easton, reached by a slight descent, is Wothorpe Grove. Here, to the right, on the excavated side of the road, and high up on the southern escarpment of the Welland valley, may also be seen the Ferruginous beds of the Northampton Sand, overlying the Upper Lias Clay. The Lias on this side attains to a much higher elevation than on the Rutland side of the valley, and extends, in a kind of ridge, still capped with the ironstone, for a mile and a half to the north-east, into Burghley Park; at which point, it considerably overtops the town of Stamford. A copious spring, the drainage of the Lincolnshire Limestone and Northampton Sand to the south, occurs near to the Wothorpe ruins, at the junction of the latter bed with the Lias Clay; and the water is caught in reservoirs, which are of sufficient elevation to supply by gravitation every part of the town of Stamford.

FAULTS.

This discrepancy of level is the result of faults; one of which I have mentioned as traversing the section of the Geeston cutting near Ketton. Another strikes athwart the Welland valley and the Stamford parish of St. Martin, and extends some miles to the east. To the latter fault, and to its very remarkable and apparently anomalous consequences, I shall again allude*.

I would explain here, that, in the route I have taken for the purpose of description, many discrepancies of level, the effects of dip, of the bending of strata, and of faults, might have been noticed; but as these are not necessarily pertinent to my object, and have been mapped, and will be described, by Mr. Judd, I have deemed it not desirable to add to the length of my Memoir by particularly referring to them.

AREA OF STAMFORD FIELD AND NEIGHBOURHOOD.

Crossing the Welland valley, to the north of the town of Stamford, and into a nook of Lincolnshire, we come to the high ground well known as the Stamford Open Field, now in the process of being enclosed.

The highest point of Stamford Field is nearly 200 feet above the level of the river Welland. In this one hill-mass, we have the

* See Diagram of Section, Plate X. fig. 1.

hardly to be over-estimated advantage of a complete sequence of beds, ranging downwards from the Cornbrash to the Upper Lias inclusive. Here the beds tell their own tale, leaving little for surmise or for question as to their identity or order*.

Recent operations in connexion with enclosure led to the opening, at the highest point, of a shallow pit; thus exposing a section of Cornbrash, having a thickness of about four feet, seldom greatly exceeded in the district. From the stone of this pit, I obtained some usual Cornbrash fossils:—large *Ostrea Marshii*, Sow., *Terebratula sub-lagenalis*, Dav., *Dentalium entaloides*, Deslong., *Echinobrissus orbicularis*, Phil., &c. &c.

The Great Oolite Clay underlies the Cornbrash, although not exposed in any section. Below this, are beds of the Great Oolite Limestone; from which I have obtained:—

Avicula echinata, Sow.

Ceromya concentrica, Sow.

Modiola imbricata, Sow.

Ostrea Sowerbyi, Mor. & Lyc.

— sub-rugulosa, Mor. & Lyc.

Rhynchonella concinna, Sow.

Ammonites, sp.?

— *gracilis*, Buckm. (large),

Nautilus Baberi (?), Mor. & Lyc. (large septum).

Clypeus Müllerii, Wright.

Lower beds of this Limestone are seen at the top of the section in Torkington's brick-pit, overlying the extremely various and variegated beds of the Upper Estuarine Series; which consist here of—a reddish sandy bed, coarse shelly clay with plants, pale grey clay streaked with exceedingly shelly seams (sometimes containing *Cyrena* and similar freshwater or estuarine shells, and sometimes *Neæra Ibbetsoni*, *Pholadomya acuticosta*, *Modiola imbricata*, &c.), dark blue clay (shells with their tests preserved), and dark chocolate laminated clays.

These Estuarine beds have an aggregate thickness of about 27 feet, and are based by the ferruginous band; here strongly pronounced, and peculiar in its rather open structure, and in the presence of much wood.

Section in Torkington's Brick-pit, Stamford Field.

1. Great Oolite Limestone—

	ft. in.	ft. in.
a. Limestone	1 0	
b. Sandy stone	1 0	
c. Grey clay	1 0	
d. Very soft marly white stone (<i>Rhynchonella concinna</i> , <i>Modiola imbricata</i> , <i>Ceromya concentrica</i>)	0 9	3 9

2. Upper Estuarine Series—

a. Reddish sandy bed, in layers, sometimes passing into a "caley" stone	4 0
b. Coarse shaly clay, with plants	3 0
c. Grey clay, in layers full of shells (<i>Cyrena</i> , <i>Neæra Ibbetsoni</i> , <i>Pholadomya acuticosta</i> , <i>Modiola imbricata</i> , <i>Ostrea</i> , &c.), and containing plants and wood	2 0

* See Diagram of Section, Plate X. fig. 1.

	ft. in.	ft. in.
d. Dark blue clay, with shells, the tests occasionally preserved	8 0	
e. Zone of chocolate-coloured indurated clay, thinly laminated	1 0	
f. Softer bed of chocolate-coloured clay—"the best for bricks"	4 0	
g. Yellowish clay	5 0	
h. Ferruginous band.....	0 9	27 9

At the base of this section, is the Lincolnshire Limestone—an oolitic freestone, similar to, and in the same position as, the upper beds of freestone in the Ketton quarries.

Fossils from the Upper Estuarine Beds, Stamford Field.

Cyrena, sp. ?	Ostrea Sowerbyi, <i>Mor. & Lyc.</i>
Modiola imbricata, <i>Sow.</i>	Pholadomya acuticosta, <i>Sow.</i>
—— Lonsdalei, <i>Mor. & Lyc.</i>	Tancredia angulata, <i>Lycett.</i>
Neera Ibbetsoni, <i>Morris.</i>	Plants and wood.

The beds of the Great Oolite Clay and Great Oolite Limestone crop out upon the escarpment of a very considerable circuit about this high ground; and the presence of the Upper Estuarine clays below is indicated by the occurrence of several brick-pits (in work or abandoned), and by a considerable stretch westward of an argillaceous moorland called the "Stamford Lings"—all upon the same level.

The freestone which appears at the base of the section in Torkington's brick-pit, is traceable westward, along the face of the escarpment, to the public Cemetery (where it has a thickness of some 10 or 12 feet), and on to the foot of the "Lings," where it was formerly quarried.

The northern boundary of the "Lings" is part of the southern boundary of Rutland; and north of this, just over and below the ridge, are the well-known Simpson's Little Casterton freestone quarries; the beds of which are geologically the same as those at Ketton, and the stone only second in quality and reputation to that of the older site: the fossils also nearly correspond. The freestone beds are surmounted by about 15 feet of the Upper Estuarine clays, which correspond exactly, in character and fossil contents, with the same series in Torkington's brick-pit.

Section at Simpson's Freestone Quarries.

	ft. in.	ft. in.
1. Upper Estuarine—Clay in various bands, and shelly zones, containing <i>Neera Ibbetsoni</i> , <i>Modiola imbricata</i> , <i>Pholadomya acuticosta</i> , crystals of selenite, &c.		15 0
2. Lincolnshire Limestone—		
a. "Crash"—a soft freestone, crumbling upon exposure to frost and atmospheric action, and useless as building-stone, in two courses	4 0 to 5 0	
b. "Rag"—an occasional pisolitic and shelly band, like the "Rag" at Weldon and Ketton.....	nil to 0 9	
c. "Top Course"—good freestone	4 0	

	ft. in.	ft. in.	ft. in.	ft. in.
d. "Bottom Course"—good freestone, but rather coarser in grain	2	0	to	2 6
e. Hard band of crystalline limestone, full of comminuted shells		0	6	
f. Marly beds, not worked.	—	—	10	6 to 12 9

Fossils from Freestone Bed, Simpson's Quarries.

Avicula, smooth species.	Terebratula globata, Sow.
Lima proboscidea, Sow. sp.	— perovalis, Sow.
Ostrea gregaria, Sow.	Chemnitzia vetusta (?), Mor. & Lyc.
— sulcifera, Phil.	Natica Leckhamptonensis, Lycett.
Lucina Wrightii, Oppel.	Nerinaa cingenda, Bronn.
Homomya (Myacites) crassiuscula, Mor. & Lyc.	— Cotteswoldia, Lycett.
Trigonia costata, var. pullus, Sow.	—, sp.? (like funiculus, Deslong.).
Unicardium depressum, Phil.	Trochus spiratus (?), d'Arch.
	Pseudodiadema depressum, Ag.

At a little less than a mile north of Simpson's quarries, after, by a gradual descent, successively passing over the outcrop of the lower beds of the Lincolnshire Limestone, the Slate beds, and the Lower Estuarine and the Ferruginous beds of the Northampton Sand, (beds of the Great Oolite Limestone and the Upper Estuarine constituting the high tract to the east,) we come, at Casterton Parva, to the valley of the small Rutland river, the Gwash (*Gausenna* of the Romans); which flows over the Ferruginous beds of the Northampton Sand.

Nearly thirty years ago, a well of no great depth was sunk at this village, and a green and very fossiliferous phase of these beds was reached. From this bed Mr. Bentley and I obtained specimens containing numerous shells in excellent condition. The mineral character of these beds is very much like that of beds in the same position at the Duston ironstone quarries near Northampton, and is probably attributable (as I explained in my First Part) to the fact that the thickness of the overlying beds has prevented atmospheric oxidation. The fossils as a group correspond with those obtained from the upper portion of the Ironstone beds at Duston.

Fossils from Green Beds of Northampton Sand, from Well at Casterton Parva, Stamford, 1844.

Avicula inæqualvis, Sow.	Lima, sp.
— sp.? (young).	Ostrea flabelloides, Lam.
Gervillia, new sp.	— sulcifera, Phil.
Hinnites abjectus, Phil. sp.	Pecten arcuatus, Sow.
Lima bellula, Mor. & Lyc.	— articulatus, Schloth.
— duplicata, Sow. sp.	— demissus, Phil.
— Dustonensis, Etheridge, M.S.	— lens, Sow.
— pectinoides, Sow. sp.	Perna rugosa, Goldf.
— Pontonis, Lycett.	Plicatula tuberculosa, Mor. & Lyc.
— proboscidea, Sow. sp.	
— punctata, Sow. sp.	Astarte elegans, Sow.
— Rodburgensis, Lycett, M.S.	Cardium (near to) Stricklandi, Mor. & Lyc.
—, large sp. (allied to L. grandis, Römer)?	Ceromya concentrica, Sow.
—, large sp.?	Cucullæa cancellata, Phil.

Cucullæa, new sp.?
Lucina, new sp.?
Modiola imbricata, *Sow.*
 — *Lonsdalei*, *Mor. & Lyc.*
 — *subreniformis*, *Mor. & Lyc.*
 — sp.?
Opis lunulatus, *Sow.* sp. (a variety).
Unicardium, sp.?

Rhynchonella sub-decorata, *Dav.*
Terebratula ovoides, *Sow.*
 — *perovalis*, *Sow.*
 — *sub-maxillata*, *Sow.*

Actæon Sedgwicki, *Phil.* sp.

Alaria trifida, *Phil.*
Natica, sp. ? (small).
Nerinæa cingenda, *Bronn.*
 — *triplicata*, *Bronn.*
Trochus, sp. ? (young).

Ammonites Murchisonæ, *Sow.*
 — —, var. *corrugatus*, *Sow.*
Nautilus clausus, *d' Orb.*
Belemnites Bessinus, *d' Orb.*

Serpula, sp. ?

Cidaris Wrightii, *Desor.*

Returning towards Stamford—below the freestone bed of the “Lings,” is a very close and brittle marly limestone, in which *Rhynchonella Crossii*, Walker, occurs. Below this, are the series of marly and crystalline beds of the Lincolnshire Limestone exposed in Tinkler’s and Squires’s quarries, having a thickness of about 29 feet; some of them very fossiliferous, and containing zones of coral. A particular bed (containing much coral, many *Nerinæa*, and other fossils), very crystalline, and taking a high polish, was formerly called the “Stamford Marble,” and was much used for chimney-pieces. The bed is still present in the section, but its mineral conditions are so altered as to unfit it for its former uses. In Squires’s quarry (which nearly adjoins Tinkler’s), a soft marly bed is thickly developed, and yields a very fine cream-coloured stone, easily worked, and (under the name of the “Stamford Stone”) much used for chimney-pieces and for the interior carved work of churches. It contains many fossils, often in fine condition; many examples of a large *Natica* and of a very large *Lima*, and a beautifully preserved frond of a cycadaceous plant, have been obtained.

Fossils from the Marly Bed of Squires’s Quarry.

Hinnites abjectus, *Phil.* sp.
Lima cardiiformis, *Sow.*
 — *Etheridgii*, *Wright.*
 — *impressa*, *Mor. & Lyc.*
 — *proboscidea*, *Sow.*
 — *Pontonis*, *Lycett.*
 — *Rodburgensis*, *Lycett.* M.S.
 —, large sp. (allied to *L. grandis*, *Römer*)?
 —, large sp.?
Pecten aratus, *Waagen.*
 — *arcuatus*, *Sow.*
 — *clathratus*, *Römer.*

Arca, large sp.?
Astarte elegans, *Sow.*
 — *recondita*, *Phil.*
Cardium Buckmani, *Mor. & Lyc.*
 — *subtrigonum*, *Mor. & Lyc.*
Ceromya Bajociana, *d' Orb.*

Ceromya similis, *Lycett.*
Cucullæa elongata, *Sow.*
Cypriocardia Bathonica, *d' Orb.*
Cyprina Jurensis, *Goldf.* sp.
 — *Loweana*, *Mor. & Lyc.*
Lucina Bellona, *d' Orb.*
 — *despecta*, *Phil.*
 — *Wrightii*, *Oppel.*
Macrodon Hirsonensis, *d' Orb.* sp.
Modiola Sowerbyana, *d' Orb.*
Myacites securiformis, *Phil.* sp.
Mytilus furcatus, *Goldf.*
Pholadomya Dewalquea, *Lycett.*
 — *Heraulti*, *Ag.*
Pholadomya lyrata, *Sow.*
 — *ovalis*, *Sow.*
 — *ovulum*, *Ag.*
 —, sp.?
Tancredia axiniformis, *Phil.*

Rhynchonella sub-tetraëdra, *Dav.*
Terebratula perovialis, *Sow.*

Natica Leekhamptonensis, *Lycett.*
— (like) *Michelini*, *d'Arch.*
—, sp. ?

Nerinea cingenda, *Bronn.*
— *Cotteswoldiæ*, *Lycett.*
— *Jonesii*, *Lycett.*

Trochotoma obtusa, *Mor. & Lyc.*
— *tabulata*, *Mor. & Lyc.*
Turbo depauperatus, *Lycett.*

Belemnites acutus, *Miller.*

Clypeus Michelini, *Wright.*

Stomechinus germinans, *Phil.*, large
var.

Calamophyllia radiata, *Lamx.*
Latimæandra Flemingi, *Edw. &*
Haime.

Thecosmilia gregaria, *M^cCoy.*

Hybodus (dorsal spine).
Strophodus magnus, *Ag.* (palates).
— *subreticulatus*, *Ag.* (palates).

Fruond of cycadaceous plant.

Ferns—
Pecopteris polypodioides, *Lindley.*
Wood.

Section in Lincolnshire Limestone, Tinkler's Quarry, Stamford.

	ft. in.	ft. in.
1. Rubbly and broken limestone.....		4 0
2. Soft concretionary marly limestone, containing Coral zones, with <i>Perna</i> , <i>Lithodomus inclusus</i> , &c.....		3 6
3. Marly limestone in thin layers, shivered		4 0
4. Compact marly limestone, in thin and irregular layers [I counted seven].....		3 6
5. Very hard limestone, containing oolitic grains, sparsely distributed, occasionally very blue- hearted	2 0 to	2 6
6. Earthy shale bed, in very thin <i>laminæ</i> , containing numerous <i>Pecten</i> s and other shells, with tests beautifully preserved, but crushed by compres- sion	1 6 to	2 0
7. "Stamford Marble"—a very hard limestone, blue-hearted, and containing much <i>Coral</i> , <i>Ne- rinæa cingenda</i> , <i>N. triplicata</i> , numerous other shells, and teeth and palates of <i>Pycnodus Buck- landi</i> and <i>Strophodus magnus</i> and <i>S. subreticu- latus</i> —formerly much more crystalline than in the present section, and then, when polished, a favourite material for chimney-pieces, &c., hence its name—in two courses	1 0 to	1 6
8. Very hard limestone, coarsely grained, in two courses	2 6 to	3 0
9. Compact marly stone, rather hard		1 6
10. Compact marly stone, softer, and containing <i>Nerinea</i> —in three courses	3 0 to	3 6
11. Rather oolitic limestone, a good building-stone		1 0

Fossils from Freestone and Shelly Beds near to Tinkler's Quarry,

Area pulchra, *Sow.*
Lima proboscidea, *Sow.* sp.
Pecten lens, *Sow.*
Pholadomya fidicula, *Sow.*

Serpula ?
Stomechinus germinans, *Phil.*
Strophodus magnus, *Ag.* (palates).

Fossils from Tinkler's and neighbouring Quarries.

Avicula clathrata, *Lycett.*
— *echinata*, *Sow.*
Gervillia acuta, *Sow.*

Lima bellula, *Mor. & Lyc.*
— *Etheridgii*, *Wright.*
— *Pontonis*, *Lycett.*

Lima Rodburgensis, *Lycett*, M.S.
 —, large sp. (allied to *L. grandis*, *Römer*)?
Ostrea flabelloides, *Lam.*
 —, large flat species.
Pecten aratus, *Waagen*.
 — *arcuatus*, *Sow.*
 — *clathratus*, *Römer*.
 — *demissus*, *Phil.*
 — *lens*, *Sow.* (or large new sp.?).
 — *personatus*, *Goldf.*

Arca Prattii, *Mor. & Lyc.*
Astarte elegans, *Sow.*
 — *minima*, *Phil.*
Ceromya Bajociana, *d'Orb.*
 — *similis*, *Mor. & Lyc.*
Cyprina Jurensis, *Goldf.*
 — *Loweana*, *Mor. & Lyc.*
 — *trapeziformis*, *Römer*.
Cypriocardia Bathonica, *d'Orb.*
 — *nuculiformis*, *Römer*.
Goniomya V-scripta, *Sow.*
Lithodermus inclusus, *Phil.*
Lucina Bellona, *d'Orb.*
 — *Wrightii*, *Oppel*.
Modiola Sowerbyana, *d'Orb.*
Myacites calceiformis, *Phil. sp.*
 — *decurtata*, *Phil. sp.*
 — *securiformis*, *Phil. sp.*
Pholadomya Dewalquea, *Lycett*.
 — *fidicula*, *Sow.*
 — *Heraulti*, *Ag.*
 — *ovalis*, *Sow.*
 —, *sp.?*
 —, *sp.?*
Tancredia axiniformis, *Phil.*
Trigonia costata, var. *pullus*, *Sow.*
 — *sculpta*, *Lycett*.
 — *V-costata*, *Lycett*.
 —, *sp. (new)?*
Unicardium, *sp.?*

Rhynchonella crucis, *Walker*.
 — *sub-decorata*, *Dav.*
 — *sub-tetraëdra*, *Dav.*
Terebratula globata, *Sow.*
 — *perovalis*, *Sow.*
 — *sphaeroidalis*, *Sow.*
 — *sub-maxillata*, *Sow.*

Actæonina, large species?
Natica (Euspira) canaliculata, *Mor. & Lyc.*
 — *formosa*, *Mor. & Lyc.*
 — *grandis*, *Goldf.*
 — *Leckhamptonensis*, *Lycett*.
Nerinea gracilis, *Lycett*.
 — *cingenda*, *Bronn.*
 — *Jonesii*, *Lycett*.
 — *Oppelii*, *Lycett*.
 — *triplicata*, *Bronn.*
Phasianella elegans, *Mor. & Lyc.*
Pterocera, *sp.?* (like *ignobilis*, *Mor. & Lyc.*).

Ammonites Murchisonæ, *Sow.**
 — *subradiatus*, *Sow.*
 — *terebratus*, *Phil.*
 — (large septa, very like species found in Ferruginous beds at Dutton).
Nautilus obesus, *Sow.*
 — *polygonalis*, *Sow.*
Belemnites Bessinus, *d'Orb.*

Serpula convoluta, *Goldf.*

Galeropygus (Hypocypus) agariciformis, *Forbes*.
Holcypus hemisphaericus, *Ag.*
Pygaster semisulcatus, *Phil.*
Pentacrinus, *sp.?*

Anabacia orbulites, *Edw. & Haime*.
Cladophyllia Babeana, *Edw. & Haime*.
Isastræa limitata, *Edw. & Haime*.
Montlivaltia tenuilamellosa, *Edw. & Haime*.
Thamnastræa, *sp.?*
Thecosmilia gregaria, *M'Coy*.

Ilybodus (large spine).
Pycnodus Bucklandi, *Ag.* (teeth).

Telosaurus (tooth).

Pecopteris polypodioides, *Lindley*.
 Coniferous wood.

Some years ago, in a rather wide crevice, formed by an open joint, in the upper beds of Tinkler's quarry, were found—canine and molar teeth of two individuals of *Hyæna* of different ages, portions of a very diminutive tooth of *Elephas*, large cervine molar (perhaps *Megaceros*?), small cervine molar, and some gnawed bones. These are in my possession, and have been kindly identified for me by Professor Rolleston, F.R.S. So far as I know, this is the only

* In the Museum of the Stamford Institution.

instance of the remains of carnivorous *mammalia*, or of any thing allied to a cave-deposit, which has occurred in this part of England.

By a well sunk at the distance of a few yards from Tinkler's quarry, limestone and other beds are shown to extend downwards for some 20 feet below the floor of the quarry, Slate appearing at the bottom. It is probable that some of these beds are analogous to the superincumbent beds of Collyweston.

Professor Morris, in his well-known Paper on the Lincolnshire Oolites, published in the Society's Journal for 1853, gives, on page 336, the following as a foot-note:—

“At Tinkler's quarry and the adjoining lands near Stamford, a typical series of the whole district may be observed. In a higher part of the hill, the stratified and bituminous clays, with the ferruginous band, may be observed, overlying the freestones (Ketton and Casterton): the lower parts of the freestones form the top of the quarry; below which—

	ft. in.
1. Compact oolitic rock, few shells	2 0
2. Concretionary compact marly oolite, full of shells, and zones of corals, the bottom more compact, the upper part marly, and decomposes more rapidly, containing shells in great abundance	4 0
3. Compact hard shelly oolitic rock, <i>Nerinea</i> , &c.....	2 6
4. Compact oolitic rock, somewhat crystalline	1 6
5. Shaly bed, irregular laminated fragments of plants, and many compressed shells, <i>Lucina</i> , <i>Pecten</i> , &c.	2 0
6. “Stamford Marble”—very compact marly limestone, full of shells and corals, <i>Nerinea</i> abundant.....	2 6
7. Indurated, somewhat marly rock	3 0
8. Compact rock	1 6
9. Compact, marly, coarse-grained oolitic rock	2 6
10. Fine-grained oolitic rock.....	1 0
11. Cream-coloured marly rock*; with <i>Nerinea</i> abundant, <i>Lima</i> , <i>Terebratula</i> , <i>Isocardia</i> (<i>Ceromya</i>), <i>Modiola</i> , <i>Lucina</i> , &c....	1 6
12. Coarse oolitic rock.....	2 feet to 26 0

Probably resting on the sands with slaty beds, which have been found in sinking lower down the hill, overlying the ferruginous rock, which covers the Upper Lias.”

This series, thus noted by Professor Morris twenty years ago, may still be considered, so far as it goes, to be “typical” of the district; but, as might be expected, the then section in Tinkler's quarry does not exactly agree with that now exposed; which, of course, is at some distance from the former site. A comparison will exhibit differences, and yet a remarkable coincidence. Although entirely different sets of figures represent the various thicknesses of the beds of the two sections, these figures, when summed up, give total thicknesses for the two sections almost identical. Thus, the section recently measured by me exposes a thickness of about 29 feet, to which may be added the further thickness of 20 feet penetrated by the well, giving a total thickness of 49 feet. Professor Morris's measured and estimated thicknesses amount together to 50 feet; the difference being only one foot.

The coincidence seems to me very significant. However variable

* This represents the marly bed, the “Stamford Stone,” of Squires's quarry.

and discrepant the rate of deposition at the two points during the passage of time represented by the whole thickness of beds, the aggregate amount of deposit at both points only differed in the proportions of 49 to 50.

The measurements of the beds of the Lincolnshire Limestone exposed at Simpson's Quarry, on the "Lings," in Tinkler's Quarry, and in the well, give 65 feet as the total thickness of the formation here.

A well at Torkington's brick-pit (half a mile to the east) pierces through a thickness of 74 feet of the same beds; this about tallies with the thickness pierced by a well sunk by Mr. Browning the architect, at a somewhat lower level in North-street, allowing for a diminution of thickness at the top.

Some excavations were recently made, at a lower level than Tinkler's Quarry, near to the Scotgate entrance to Stamford; which exhibited the Slate beds reposing on the Lower Estuarine Sands.

For the foundations and cellars of the houses of the Rock Terrace, hard by, excavations were made in the Ferruginous beds of the Northampton Sand; so that the surface of the Upper Lias Clay cannot be many feet below the level of the street at this point.

Thus it has been shown that the escarpment of the high ground north of Stamford presents a complete sequence of beds, from the Cornbrash at the summit to the Upper Lias at the base; the Lincolnshire Limestone being by far the most important formation of the whole group*.

THE DISTRICT EAST AND NORTH-EAST OF STAMFORD.

To the east and north-east of Stamford, the various beds are considerably depressed. On the road to Uffington, immediately north of the bridge which passes over the Stamford and Essendine Railway, and abutting upon the deep cutting here, is Mr. Eldret's quarry; in which is a fine section, exposing a thickness exceeding 30 feet of beds of the Lincolnshire Limestone.

The floor of this quarry is only a few feet higher than the level of the Welland river at this point, although the base of the limestone has not been reached.

This is a very good typical section of the middle beds of the formation; which here has thickened considerably. It consists of a series of fifteen distinct beds of limestone, of varying character: some are oolitic (one being a true "freestone"), and these are in an unusual position, at the bottom of the section, while others, and by far the greater part, are marly, and devoid of oolitic grains; some are soft, like the Squires's-pit "Stamford Stone," while others are hard, and sometimes crystalline and blue-hearted; some are very fossiliferous, while others are slightly so, and some apparently bare of fossils. In my detailed notes of this section, I have recorded the peculiar names by which the several beds are identified by the quarrymen.

* See Diagrammatic Section, Plate X. fig. 2.

Section at Eldret's Quarry, with Quarrymen's Terms.

	ft.	in.
1. "Rammel"—broken stone, about	4	0
2. "Clinkers"—compact marly whitish stone, very good for lime-burning: has a glistening fracture	2	0
3. "Pendle"—hard, flaggy limestone, rather oolitic, deeper in colour than the last, in thin layers	2	0
4. "Shelly Course"—composed wholly of shells with corals, very hard, ("Stamford Marble"?)	2	6
5. "Bulymong"—soft white marly limestone, containing numerous fossils, (like the "Stamford Stone" of Squires's quarry), more compact and harder towards the bottom... ..	4	0
6. "Blue Limestone"—hard compact stone, blue-hearted, good rubble walling-stone	4	6
7. Course of cream-coloured clay	0	2
8. Hard limestone, with oolitic grains	1	0
9. "Bastard Freestone"—an oolitic limestone, in two courses	3	0
10. Hard and compact marly course	0	7
11. Soft white marly limestone (like the "Stamford Stone" of Squires's quarry), in four courses of different thickness	5	0
12. "Caley" oolitic bed (like some of the upper beds at Collyweston)	1	0
13. "Bastard Freestone," containing concretionary masses of very hard limestone	1	6
14. "Freestone"—a good oolitic freestone	1	1
15. Limestone—thickness not ascertained.		

BELMISTHORPE.

Two miles north of Eldret's Quarry, the railway, after crossing the valley of the Gwash, passes, by a cutting, through the Great Oolite Limestone and the underlying Upper Estuarine Clays. This cutting is distant only two miles to the north-east from the summit of the Stamford Field, and yet the beds here exposed are nearly 100 feet lower in level than the same beds in their elevated position at the last-named site. From these beds at Belmistorpe, I have an abundance of Great Oolite fossils of forms rendered familiar by their frequent occurrence in the Great Oolite beds of the Northampton district.

Fossils from Great Oolite Limestone at Belmistorpe.

<i>Lima cardiiformis</i> , Sow. sp.	<i>Homomya crassiuscula</i> , Mor. & Lyc.
<i>Ostrea sub-rugulosa</i> , Mor. & Lyc.	<i>Isocardia tenera</i> , Sow.
<i>Pecten annulatus</i> , Sow.	<i>Macrodon Hirsonensis</i> , d'Arch. sp.
<i>Pinna cuneata</i> , Phil.	<i>Modiola imbricata</i> , Sow.
—	<i>Myacites calceiformis</i> , Phil. sp.
<i>Cardium Buckmani</i> , Mor. & Lyc.	— <i>securiformis</i> , Phil. sp.
— <i>subtrigonum</i> , Mor. & Lyc.	— <i>Terquemea</i> , Buv. sp.
<i>Ceromya Symondsii</i> , Mor. & Lyc.	<i>Næra Ibbetsoni</i> , Morris.
—, sp. ?	<i>Pholadomya deltoidea</i> , Sow.
<i>Corbis Bathonica</i> , Mor. & Lyc.	— <i>lyrata</i> , Sow.
<i>Cucullæa</i> , sp. ?	— <i>oblita</i> , Mor. & Lyc.
<i>Cypricardia Bathonica</i> , Mor. & Lyc.	— <i>socialis</i> , Mor. & Lyc.
<i>Cyprina depressiuscula</i> , Mor. & Lyc.	— <i>solitaria</i> , Mor. & Lyc.
<i>Gresslya peregrina</i> , Phil.	—, new round sp. ? with irregular <i>costæ</i> .*
<i>Homomya Vezelayi</i> , d'Arch. sp.	

* Found also at Danes' Hill, near Peterboro', and near Northampton—all in the Great Oolite Limestone.

Trigonia costata, var. *pullus*, *Sow.*

—, var. *elongata*, *Sow.*

— *Moretoni*, *Mor. & Lyc.*

Unicardium varicosum, *Sow.*

Natica formosa, *Mor. & Lyc.*

— *grandis*, *Goldf.*

— *intermedia*, *Mor. & Lyc.*

Nautilus Baberi, *Mor. & Lyc.* (very numerous and large).

— *subtruncatus*, *Mor. & Lyc.*

Crustacean—

Eryma (allied to) *elegans*, *Oppel.*

Fossils from the Upper Estuarine Clays, Belmistorpe.

Terebratula intermedia, *Sow.*

Crustacean—

Eryma (allied to) *elegans*, *Oppel.*

Hybodus, large dorsal spine.

Teleosaurus, large vertebrae and bones.

—, large and numerous scutes.

—, small jaw.

—, small atlas.

Fossils from Great Oolite, Uffington, adjoining Belmistorpe.

Lima duplicata, *Sow.* sp.

Isocardia tenera, *Sow.*

Pholadomya deltoidea, *Sow.*

Pholadomya Heraulti, *Ag.*

Ammonites macrocephalus, *Schloth.*

(a large example).

GREAT NORTHERN RAILWAY.

Half a mile beyond the Belmistorpe cutting, the Stamford and Essendine Railway joins the main line of the Great Northern Company. The sections exposed in the cuttings of this line between Peterboro' and Grantham have been rendered classical by their description by Professor Morris in a Paper published nearly twenty years ago in the Quarterly Journal of this Society*; and which Paper constitutes the only existing reliable record of these valuable sections, long since covered up.

In this long interval, some advance has been made in the knowledge of the geology of this part of England; and, as additional light has been thrown upon the character and sequential position of the beds then described, I shall perhaps be justified in redirecting attention to these sections, and, while adopting Professor Morris's descriptions, in venturing to assign some of the beds to other formations than those to which he referred them. It must not be forgotten, however, that in his interesting article in the 'Geological Magazine' for March, 1869, Professor Morris indicated that the views which he had formerly entertained had undergone much modification†.

By taking these sections in succession in a direction from south to north, a very complete sequence of the beds (from the Oxford Clay to the Lincolnshire Limestone inclusive) of this Midland part of the country will be obtained; and I shall endeavour to apply them as one group of evidence as to what that sequence really is,

* Nov. 1853.

† As far back as the date of Professor Morris's Paper in 1853, Dr. Lycett had arrived at the provisional conclusion that the beds now known as the Lincolnshire Limestone were Inferior Oolite; and even three years previously, the Rev. P. B. Brodie, F.G.S., in a paper published in the 'Proceedings of the Cotteswold Field Club,' expressed an opinion that some beds of the limestones of Lincolnshire should be classed rather with the Inferior than with the Great Oolite.

Fig. 2.—Section at Casewick Cutting*.

Length 39 chains. Vertical scale 120 feet to 1 inch.

Fig. 3.—Section at Danes'-Hill Cutting*.

Length 22 chains. Vertical scale 120 feet to 1 inch.

* Taken by the kind permission of Professor Morris, F.G.S., from his Paper "On the Lincolnshire Oolites," Quart. Journ. Geol. Soc., Nov. 1853.

and as to the true geological character of the beds of which it consists.

During the progress of the excavation of these cuttings, Mr. Bentley and I had the advantage of traversing them more than once in the instructive company of Professor Morris; and I am fortunate in possessing, and in being enabled to exhibit a selection from, fossils gathered on those occasions by Mr. Bentley and myself.

CASEWICK CUTTING.

The most southerly of these cuttings is at Casewick, about three miles to the south-east of Stamford. The upper part of the section consists of 8 feet of Quaternary Drift beds; between which and the Secondary beds, is intercalated, for the space of about 30 yards, a freshwater deposit: these are of sufficient interest in themselves, but are foreign to my subject. Professor Morris describes the uppermost Secondary bed in this cutting as being Oxford Clay: he then gives Kelloway Rock, and then Oxford Clay again: the lowest bed of the section and the floor of the cutting consist of Cornbrash.

I have characteristic fossils from each of the beds of this section, including—*Ammonites Herveyi*, *A. biplex*, and *A. Gowerianus*, from the Oxford Clay; *Gryphæa bilobata*, in every stage of growth, and the very characteristic *Avicula expansa*, from the Kelloway Rock; and two fair specimens of the rare and peculiar *Terebratula Bentleyi* from the Cornbrash.

Fossils from the Casewick Cutting.

OXFORD CLAY.

Ammonites biplex, Sow.

Am. Gowerianus, Sow.

Am. Herveyi, Sow.

KELLOWAY ROCK.

Avicula expansa, Phil.

Gryphæa bilobata, Sow.

— *dilatata*, Sow.

Ostrea, sp.?

—, sp.?

Pecten, sp.?

Gresslya peregrina, Mor. & Lyc.

Modiola pulchra, Phil.

Myacites calceiformis, Phil. sp.

Myacites recurva, Phil. sp.

Myopsis dilatata, Phil. sp.

(or new sp.?).

— sp.?

Trigonia, sp.?

Serpula intestinalis, Phil.

Belemnites hastatus, Blainv.

Megalosaurus, sp. ? (bone).

CORNBRASH.

Lima pectiniformis, Schloth.

— *rigida*, Sow. sp.

— *rigidula*, Sow. sp.

Ostrea, large flat sp.

Goniomya V-scripta, Sow. sp.

(large var.)

Pholadomya acuticosta, Sow.

Rhynchonella concinna, Sow. sp.

— *Morieri*, Dav.

Terebratula Bentleyi, Morris.

Serpula intestinalis, Phil.

—, sp.?

Strophodus magnus, Ag. (palates).

BANTHORPE CUTTING.

At less than a mile north of Casewick, is the Banthorpe cutting.

The Cornbrash which based the last section appears at the top of this: beneath the Cornbrash, appears the Great Oolite Clay; and underlying it, to the depth of 7 feet, is the Great Oolite Limestone of Belmistorpe and the Stamford Field. From the Clay, I obtained a fine caudal vertebra of *Cetiosaurus*, that largest of all known Saurians; and this Clay, be it noted, is in the exact relative position as that at Kirklington in Oxfordshire, whence Professor Phillips obtained the wonderful and unique skeleton with which he has enriched the Oxford Museum. From the Limestone, I collected a large example of *Nautilus Baberi*, which occurs so abundantly in the Great Oolite Limestone of Northampton, and which equally distinguishes the equivalent bed at Belmistorpe. The Cornbrash at Banthorpe has yielded *Pecten vagans*, Sow., and *Myacites calceiformis*, Phil. sp.

ESSENDINE, AUNBY, AND DANES' HILL.

A mile from Banthorpe, is the Essendine cutting, and at short intervals the Aunby and the Danes'-Hill cuttings. Professor Morris has given the details of these sections with considerable minuteness; but, as they are not widely separated, and present no differences that will affect my position, I will include them in one general description:—

The Cornbrash, which bases the Casewick cutting, and caps the Banthorpe section, occurs in patches upon the summit of the section at Danes' Hill, and thus serves to preserve the continuity of the sequence through this widely extended, interesting, and significant series of sections. (See *antè*, figs. 2 and 3.)

Below the Cornbrash, (and occurring at Danes' Hill only,) are compact marly and sandy beds, with shells, about 5 feet in thickness, which perhaps may be considered equivalent to Forest Marble.

These overlie the Great Oolite Clay, containing bands of *Ostrea sub-rugulosa* in a normal horizontal position, and occasional patches of *Serpulæ*. This clay, in the several sections, varies in thickness from 5 feet to 8 feet.

Next, in descending order, is the Great Oolite Limestone—8 feet to 12 feet—described by Professor Morris as a “sandy and marly rock, becoming occasionally very compact, calcareous, bluish, and sometimes shaly”: at Danes' Hill, it is divided by argillaceous bands. From this bed, many Great Oolite fossils have been obtained.

The Upper Estuarine series consists here of numerous bands of very varying sands and clays; which are sometimes very shaly, sometimes bituminous, and occasionally assume the character of a soft marly rock. The fossils noted are of well-known Great-Oolite forms, interspersed with some of undescribed species. The presence of layers of *Cyrena*, other estuarine shells, much wood (almost converted into jet), and of other vegetable matter, characterizes this bed; which is traversed at Essendine and Danes' Hill by one band, and at Aunby by two bands, of peculiar plant-growth (precisely

like that described by me as occurring in the sand-bed at Kings-thorpe), consisting of vertical root-perforations surmounted by horizontal bituminous layers. This Upper Estuarine series was not wholly exposed at Essendine: at Danes' Hill, it had a thickness of about 22 feet; while at Aunby, it had thickened to no less than 32 feet. The usual ferruginous band, of the thickness of about a foot, based these clays.

At the southern end of the Danes'-Hill cutting, and immediately underlying the ferruginous band, an upper bed of the Lincolnshire Limestone was exposed.

In the Essendine cutting, and probably in the Great Oolite Clay (the same clay as that at Banthorpe and at Kirklington), immense bones of *Cetiosaurus brevis* and of *C. longus* were found; which bones are now in the Museum in Jermyn Street.

As no useful end can be served in keeping distinct the fossils of these sections, I have grouped them together in the following list, in order to economize space and to avoid repetition.

Fossils from the Essendine and Danes' Hill Cuttings.

CORNBRASH.

Terebratula ornithocephala, Sow.

Terebratula sub-lagenalis, Dav.

GREAT OOLITE LIMESTONE.

Gervillia Islipensis, *Lycett*.

Lima cardiiformis, Sow. sp.

— *duplicata*, Sow. sp.

Ostrea Sowerbyi, Mor. & Lyc.

— *sub-rugulosa*, Mor. & Lyc.

—, sp. ?

Pecten annulatus, Sow.

— *lens* (?), Sow. (or new sp. ?).

—, sp. ?

Perna foliacea, *Lycett*.

— (like) *mytiloides*, Lam.

— *quadrata*, Sow. sp.

— *rugosa*, Goldf.

—, sp. new ?

Pteroperna emarginata, Mor. & Lyc.

—, *plana*, Mor. & Lyc.

—, sp. ?

Arca Eudesii, Mor. & Lyc.

—, sp. ?

Cardium Buckmani, Mor. & Lyc.

— *cognatum*, Phil.

— *lingulatum*, *Lycett*.

— *Stricklandi*, Mor. & Lyc.

— *subtrigonum*, Mor. & Lyc.

—, sp. ?

Ceromya Symondsii, Mor. & Lyc.

Cypricardia caudata, *Lycett*.

Cypricardia nuculiformis, Rümer.

Cyprina Davidsoni, *Lycett*.

— *Lowciana*, Mor. & Lyc.

—, sp. ? (var. *elongata* ?).

Isocardia tenera, Sow.

Modiola cuneata, Sow.

— *gibbosa*, Sow.

— *imbricata*, Sow.

— *Lonsdalei*, Mor. & Lyc.

— *subreniformis*, Mor. & Lyc.

Myacites calceiformis, Phil. sp.

— *securiformis*, Phil. sp.

Nevra Ibbetsoni, Morris.

Opis, new sp.*

Pholadomya deltoidea, Sow.

— *Heraulti*, Ag.

— *lyrata*, Sow.

— *Phillipsii*, Morris.

— *socialis*, Mor. & Lyc.

—, new sp. ? with a few coarse *costæ*†.

—, new round sp. ? with irregular *costæ*‡.

Tancredia axiniformis, Phil.

Trigonia costata, Sow.

— *Moretoni*, Mor. & Lyc.

Unicardium varicosum, Sow.

Rhynchonella concinna, Sow. sp.

* Found also at Stowe-Nine-Churches, near Weedon.

† Found also near Peterboro' and at Kingsthorpe.

‡ Found also at Belmishthorpe, near Peterboro', and near Northampton, all in Great Oolite Limestone.

Terebratula intermedia, Sow.
 — *maxillata*, Sow.
 — *obovata*, Sow.

Amberleya nodosa, Mor. & Lyc.
Delphinula?
Natica formosa, Mor. & Lyc.
 — *globosa*, Römer.
 — *intermedia*, Mor. & Lyc.
 — (*Euspira*) *canaliculata*, Mor. & Lyc.
 — — *pyramidata*, Mor. & Lyc.
 — — *Sharpei*, Mor. & Lyc.
Trochotoma, sp.?

Nautilus Baberi, Mor. & Lyc.
 — *subtruncatus*, Mor. & Lyc.

Serpula, new sp.

Acrosalenia hemicidaroides, Wright.
 — *pustulata*, Forbes.
 — *Wiltoni*, Wright.
Clypeus Mülleri, Wright.

Crustacean—
Eryma (allied to) *elegans*, Oppel.

UPPER ESTUARINE CLAYS*.

Avicula, sp.?
Lima, sp.?
Ostrea Sowerbyi, Mor. & Lyc.
 —, sp.?
Pecten retiferus (?), Mor. & Lyc.
 —, sp.?
Pinna, new species?
Arca rugosa, Mor. & Lyc.
 —, sp.?
Astarte angulata (?), Mor. & Lyc.
Cardita?
Cardium Stricklandi, Mor. & Lyc.
 — *subtrigonum*, Mor. & Lyc.
Cucullaea concinna, Phil. sp.
 — *triangularis* (?), Phil. sp.
 —, sp.?
Cypricardia Bathonica, d' Orb.
 — *caudata*, Lycett.
 —, sp.?
Cyprina Loweana, Mor. & Lyc.
 —, var. *elongata*?
 —, sp.?
Cyrena, sp.?
Gresslya, sp.?

Modiola imbricata, Sow.
 — *gibbosa*, Sow.
 —, sp.?
Myacites, sp.?
Negara Ibbetsoni, Morris.
Pholadomya acuticosta, Sow.
Tancredia axiniformis, Phil.
 — *planata*, Mor. & Lyc.
Tellina, sp.?
Thracia, sp.?
Unicardium gibbosum, Mor. & Lyc.
 — *impressum*, Mor. & Lyc.
 — *varicosum*?, Mor. & Lyc.

Rhynchonella concinna, Sow. sp.

Cerithium, sp.?
Natica, sp.?

Acrosalenia, sp. ? (spines).

Plants—in both horizontal and vertical positions.
 Wood—converted almost into jet.

Identity of the Fossils of the Great Oolite of Stamford with those of the Great Oolite of Northampton.

As I shall not again refer in detail to the beds of the Great Oolite, I will here point to the identity of the organic forms which are most abundant or most significant in the beds of that formation in the Northampton district, with those of the greatest frequency or of equal significance in the beds which I have termed "Great Oolite" in the Stamford district; by which identity, as I think, the equivalency of these beds in the two localities is palæontologically established.

The most prominent fossil probably in the Great Oolite Limestone of Northampton is the large *Nautilus Baberi*; of which numerous examples have been found in the limestone of Belmishthorpe,

* The fossils in the Upper Estuarine Clays are for the most part very imperfectly preserved; and it must be understood, therefore, that the fossils in the list marked with a note of interrogation are only probably identified.

and others in similar beds at Banthorpe, Essendine, and near Peterboro'.

Large specimens of *Ammonites gracilis*, Buckm., occur in the Northampton beds; I have a less large example from Stamford.

Natica formosa, *N. intermedia*, and other species of *Natica* having elevated spines, occur in great numbers at Kingsthorpe and Blisworth; they are prevalent forms also at Belmistorpe, Essendine, and Danes' Hill.

Pholadomya (deltoidea, Heraulti, lyrata, socialis, &c.) is a most abundant genus in the Northampton beds: it is as abundant and various at Belmistorpe, and also occurs at Essendine and Danes' Hill.

Modiola imbricata is equally numerous in the beds of both districts, as are also *Myacites calceiformis*, *Cypriocardia Loweana*, and *Isocardia tenera*.

Terebratula maxillata and *Rhynchonella concinna*, such marked forms at Blisworth and Kingsthorpe, distinguish also Danes' Hill and Essendine.

Other forms, of less abundance but of much significance, are common to the beds which I have termed "Great Oolite" in both localities; such as the rare *Amberleya nodosa* of Mor. & Lyc., the still more rare *Necera Ibbetsoni* (known only, I believe, in the Great Oolite beds of these localities), and the beautiful undescribed crustacean allied to *Eryma elegans* of Oppel, so abundant at Buttock's Booth near Kingsthorpe, and also found at Belmistorpe, Essendine, and Danes' Hill.

Besides these, are the abundant and puzzling series of shell-less fossils, so difficult of recognition; and which the great experience and persistent patience of Mr. Etheridge alone enabled him approximately to identify. Among these, are *Ceromya Symondsii*, *Cypriocardia Bathonica*, *Cardium Buckmani*, *Unicardium*, and others; the similarity of which, both as to condition and form, in the groups obtained from the two localities, also serves to mark the parallelism of the beds whence they have been derived.

LITTLE BYTHAM.

At Little Bytham, two miles north of Danes' Hill, the Great Oolite Limestone has disappeared; but the Upper Estuarine Clays exhibit a thickness of about 30 feet, and the ferruginous band at the base is still present, as it is in several other cuttings further north. Below the Clays, the Lincolnshire Limestone appears in the form of good oolitic freestone; and this (for the sake of the stone) was excavated, during the making of the railway, to a considerable depth below the floor of the cutting. Under 10 feet of this freestone, was found a compact marly limestone, 3 feet in thickness; and beneath this again, 8 feet of good freestone.

THE PONTONS.

Four miles south of Grantham, is the Great Ponton cutting. Under about 6 feet of the Upper Estuarine Clays, are 37 feet of the

beds of the Lincolnshire Limestone:—Beneath 5 feet of rubbly oolite, are a series of shelly, coralline, and oolitic bands (having an aggregate thickness of 15 feet), which have yielded a perfect harvest of fossils, including many forms not previously known; some of which were described and figured by Dr. Lycett in an Appendix to Professor Morris's Paper. Under these shelly bands, is a soft marly limestone, 3 feet; and below this, 15 feet of "coarse shelly oolites and freestones."

Nearer Grantham, is the Little-Ponton cutting; in which is exposed 50 feet of the beds of the Lincolnshire Limestone. A well in the neighbourhood pierced through some 60 feet of the same beds.

Fossils from the Shelly Beds, Ponton.

Gervillia acuta, Sow.
Hinnites velatus, Goldf. sp.
 — *regulatus*, Mor. & Lyc.
Lima bellula, Mor. & Lyc.
 — *cardiiformis*, Mor. & Lyc.
 — *duplicata*, Sow. sp.
 — *gibbosa*, Sow. sp.
 — *Pontonis*, Lycett.
 — *punctata*, Sow. sp.
Ostrea flabelloides, Lam.
 — *gregaria*, Sow.
 — *sulcifera*, Phil.
Pecten lens, Sow.
 —, sp.?
Plicatula tuberculosa, Mor. & Lyc.
 —, sp.?
Pteroperna costatula, Deslong. sp. (young).
 — *plana*, Mor. & Lyc.
 — *pygmæa*, Dunker, sp.

Arca æmula, Phil.
 — *Prattii*, Mor. & Lyc.
 — *pulchra*, Sow.
Astarte depressa, Goldf.
 — *elegans*, Sow.
 — *excentrica*, Mor. & Lyc.
 — *minima*, Phil.
 — *Pontonis*, Lycett.
 — *Wiltoni*, Mor. & Lyc.
 —, sp.?
Cardium, sp.?
Ceromya similis, Lycett.
Cypricardia Bathonica, d'Orb.
Cyprina nuciformis, Lycett.
 —, sp.?
Cytherea (Cyprina) dolabra, Phil.
Homomya crassiuscula, Mor. & Lyc.
Lucina Bellona, d'Orb.
 — *despecta*, Phil.
 —, var. *cardioides*, d'Arch.
 — *rotundata*, Römer.
 — *Wrightii*, Oppel.
Macrodon Hirsonensis, d'Arch. sp.
Modiola Binfieldi, Mor. & Lyc.

Modiola gibbosa, Sow.
 —, sp.?
Myoconcha crassa, Sow.
Opis gibbosus, Lycett.
 — *lunulatus*, Sow. sp.
 — *similis*, Desh. (or new sp.?).
 —, smooth new sp.?
Tancredia angulata, Lycett.
 — *axiniformis*, Phil.
Trigonia costata, Park.
 —, var. *pullus*, Mor. & Lyc.
 — *hemispharica*, Lycett.
Unicardium depressum, Phil. sp.
 — *parvulum*, Mor. & Lyc.

Rhynchonella, sp.?
 — *Crossii*, Walker.
 — *sub-decorata*, Dav.
Terebratula Buckmani, Dav.
 — *fimbria*, Sow. (young).
 — *globata*, Sow.
 — *ovoides*, Sow.
 — sp.? (numerous young).

Actæonina glabra, Phil. sp.
 —, sp.?
Alaria armata, Mor. & Lyc.
 — *hamulus*, Deslong. sp.
 — *hamus*, Deslong. sp.
 — *Phillipsii* (?), d'Orb. sp.
 — *sub-punctata*, Goldf. sp.
 — *trifida*, Phil. sp.
Ceritella acuta, Mor. & Lyc.
 —, sp. (like Sowerbyi, Mor. & Lyc.)?
Cerithium Beanii, Mor. & Lyc.
 —, sp. (like *costigerum*, Piette)?
 — *gemmatum*, Mor. & Lyc.
 — *limæforme* (?), Römer.
 — *quadrinectum*, Goldf.
 —, new sp.?
 —, sp.?
Chemnitzia vetusta, Phil. sp.
Cylindrites acutus, Sow. sp.
 — *brevis*, Mor. & Lyc.

<i>Cylindrites bullatus</i>, <i>Mor. & Lyc.</i> — <i>cylindricus</i>, <i>Lycett.</i> — <i>turriculatus</i>, <i>Lycett.</i> <i>Delphinula</i> (<i>Crassostoma</i>) <i>Prattii</i>, <i>Mor. & Lyc.</i> —, sp.? <i>Lithodomus inclusus</i>, <i>Phil.</i> <i>Monodonta Lyellii</i> (?), <i>d'Arch.</i> —, sp.? <i>Nerina</i> <i>Cotteswoldiæ</i>, <i>Lycett.</i> —, sp. (like small <i>Eudesii</i>, <i>Mor. & Lyc.</i>)? —, sp. (like <i>funiculus</i>, <i>Desl.</i>)? — <i>punctata</i>, <i>Voltz.</i> — <i>triplicata</i>, <i>Bronn.</i> — <i>Voltzii</i>, <i>Deslong.</i> <i>Onustus</i>, sp. (like <i>Burtonensis</i>, <i>Lycett.</i>)? <i>Phasianella elegans</i>, <i>Mor. & Lyc.</i> — <i>latiuscula</i>, <i>Mor. & Lyc.</i> — <i>parvula</i>, <i>Mor. & Lyc.</i> — <i>Pontonis</i>, <i>Lycett.</i> — <i>tumidula</i>, <i>Mor. & Lyc.</i> <i>Pterocera Bentleyi</i>, <i>Mor. & Lyc.</i> <i>Pileolus plicatus</i>, <i>Sow.</i> (an acute var.). <i>Rimula Blotii</i>, <i>Deslong.</i> <i>Rissoina obliquata</i>, <i>Sow.</i> sp. —, sp.? <i>Solarium</i>, sp. (somewhat like <i>varicosum</i>, <i>Mor. & Lyc.</i>)? <i>Trochotoma extensa</i>, <i>Mor. & Lyc.</i> <i>Trochus Dunkeri</i>, <i>Mor. & Lyc.</i> — <i>bijugatus</i> (?), <i>Quenst.</i>	<i>Trochus Ibbetsoni</i>, <i>Mor. & Lyc.</i> — <i>Leckenbyi</i>, <i>Mor. & Lyc.</i> — <i>monilitectus</i>, <i>Phil.</i> —, a var. — <i>ornatissimus</i>, <i>d'Orb.</i>, var. <i>Pontonis</i>, <i>Morris.</i> — <i>spiratus</i>, <i>d'Arch.</i> —, a var. — <i>squamiger</i>, <i>Mor. & Lyc.</i> —, sp.? <i>Turbo bijugatus</i> (?), <i>Quenst.</i> — <i>gemmatus</i>, <i>Lycett.</i> — sp. (like <i>Gomondei</i>, <i>Mor. & Lyc.</i>)? — <i>ornatus</i> (?), <i>Quenst.</i> (<i>Littorina ornata</i> (?), <i>Sow.</i>). — <i>Phillipsii</i>, <i>Mor. & Lyc.</i> — new sp.? <i>Serpula socialis</i>, <i>Goldf.</i> — <i>sulcata</i>, <i>Sow.</i> — large sp.? <i>Vermicularia nodus</i>, <i>Phil.</i> <i>Pseudodiadema depressum</i>, <i>Ag.</i> <i>Pentacrinus</i>, sp. ? (joints). Claws of Crustacea. <i>Cladophyllia</i>, sp. ? <i>Isastræa limitata</i>, <i>Edw. & Haime.</i> <i>Thecosmilia</i>, sp. ? <i>Polyzoa</i>, sp. ?

Besides the foregoing, many minute young or new forms have been obtained from the Ponton shelly beds; but which, although often well preserved, it has been found impossible to identify with known species.

DENTON.

I will not attempt to trace the occurrence of these beds further north; but will incidentally draw attention to the fact that at Denton (6 miles south-west of Grantham, and 18 miles due north of Morcot), is a bed of the Lincolnshire Limestone; which has yielded nearly the same suite of fossils as the last-named place (including the fine example of *Pterocera Bentleyi* mentioned in my notice of the Collyweston Slate beds), with the addition of several univalve and bivalve forms, and some beautiful and probably new varieties of Coral.

Fossils from Lincolnshire Limestone, Denton.

<i>Gervillia lata</i>, <i>Phil.</i> <i>Hinnites velatus</i>, <i>Goldf.</i> sp. <i>Lima cardiiformis</i>, <i>Sow.</i> sp. — <i>impressa</i>, <i>Mor. & Lyc.</i> — <i>Luciensis</i>, <i>d'Orb.</i> — <i>Pontonis</i>, <i>Lycett.</i> — <i>proboscidea</i>, <i>Sow.</i> sp. <i>Perna quadrata</i>, <i>Phil.</i>	<i>Pinna cuneata</i>, <i>Phil.</i> <i>Placunopsis ornatus</i>, <i>Mor. & Lyc.</i> <i>Ceromya similis</i>, <i>Lycett.</i> <i>Cucullæa Goldfussii</i>, <i>Römer.</i> <i>Lucina Wrightii</i>, <i>Oppel.</i> <i>Macrodon Hirsonensis</i>, <i>d'Arch.</i> sp. <i>Myacites decurtata</i>, <i>Goldf.</i> sp.

Pholadomya fidicula, *Sow.*
 — *ovalis*, *Ag.*
 — *Zeiténii*, *Ag.*
Trigonia hæmisphærica, *Lycett.*
 — *Phillipsii*, *Mor. & Lycett.*
 — *striata*, *Miller* (large).
Unicardium depressum, *Phil.*

Terebratula fimbria, *Sow.* (young of).
 — *perovalis*, *Sow.*
 — *sub-maxillata*, *Sow.*

Alaria armata, *Mor. & Lyc.*
Chemnitzia Wetherellii, *Mor. & Lyc.*
Natica adducta, *Phil.*
 — *Leckhamptonensis*, *Lycett.*
 — *punctura*, *Bean*, sp.
Nerinaea cingenda, *Bronn.*
 —, sp. (like *funiculus*, *Deslong.*)?
 — *triplicata*, *Bronn.*
Pleurotomaria armata, *Münst.*

Pleurotomaria, sp. (allied to *Marcou-
 sana*, *d'Orb.*)?
 — *ornata*, *Defrance.*
 — *sulcata*, *Sow.* sp.
 —, sp.?
Pterocera Bentleyi, *Mor. & Lyc.*

Serpula, new sp. *?

Acrosalenia Lycettii, *Wright.*
Clypeus Michelini, *Wright.*
Galeropygus agariciformis, *Forbes.*
Pygaster semisulcatus, *Phil.*

Calamophyllia, sp. (like *Stokesii*, *Ed-
 wards & Haime*)?
Montlivaltia tenuilamellosa, *Edw. &
 Haime.*
Thecosmilia gregaria, *M'Coy*—a var.
 Genus? sp.? †
 Genus? sp.?

BOURN.

From Cornbrash beds near Bourn, about 5 miles north-east of Essendine, I have the following fossils:—

Exogyra, nov. sp. (like *E. virgula*,
Def.)?
Lima impressa (?), *Mor. & Lyc.*
 — *rigida*, *Sow.* sp.
 — *rigidula*, *Phil.* sp.
Ostrea (flat round large sp.?).
Pecten vagans, *Sow.*

Cardium subtrigonum, *Mor. & Lyc.*
Goniomya V-scripta, *Sow.* sp.
Homomya crassiuscula, *Mor. & Lyc.*
 (young).
Isocardia tenera, *Sow.*
Lucina Lycettii, *Oppel.*
Modiola gibbosa, *Sow.*
Myacites modica, *Bean*, sp.
 — *securiformis*, *Phil.* sp.
 — *sinistra*, *Ag.* sp.
 — *Terquemea*, *Buv.* sp.

Myopsis Jurassi, *Ag.*
Pholadomya, sp.?
Trigonia Scarburgensis, *Lycett.*
Unicardium, sp.?

Rhynchonella varians, *Schloth.*
Terebratula Bentleyi, *Morris.*
 —, (young examples).
 — *coarctata*, *Park.*
 — *intermedia*, *Sow.*
 — *lagenalis*, *Schloth.*
 — *sub-lagenalis*, *Dav.*

Ammonites macrocephalus, *Schloth.*

Serpula intestinalis, *Phil.*
 — *tetragona*, *Sow.*

Echinobrissus orbicularis, *Phil.*

FAULT AT STAMFORD ‡ AND EASTWARD.

Crossing the Welland, from the old Lincolnshire town of Stamford to the Northamptonshire parish of St. Martin, the important fault laid down in Mr. Judd's map and its consequences are encountered. Of the precise line taken by this fault, I was unaware until I saw that map, although well acquainted with its peculiar and anomalous results; which extend some miles to the east, and some of which were

* From a band, with plants and wood, similar to that overlying Slate-bed at Collyweston.

† See remarks upon these Corals by Professor Duncan, F.R.S., in discussion at the conclusion of this Paper.

‡ See Diagram of Stamford Section, Plate X. fig. 1.

described by me twenty years ago. These, however, I had not associated with *one* fault, as it is not uncommon in districts familiar to me to find beds near the brow of an escarpment much dislocated and their levels disturbed.

I have stated that the Upper Lias Clay, capped by the Northampton Sand, occurs at a considerable elevation south of Stamford, quite overtopping the town. The effect of the forces producing this fault, whilst probably elevating the beds which appear at so high a level, was to depress other beds throughout a considerable area.

Thus, at Stamford Bridge, the Upper Lias Clay is only just up to the level of the bed of the river; and in ascending the hill from this point through St. Martin's, will be passed over in succession—the Ferruginous beds of the Northampton Sand, the Lower Estuarine sands and clays, the Collyweston Slate and Lincolnshire Limestone beds, and the Upper Estuarine Clays; then again, in reiterated sequence, a great thickness of Upper Lias Clay, the Ferruginous beds (worked for ironstone at the top of the hill), the Lower Estuarine beds, the Collyweston Slate, and further on the rock beds of the Lincolnshire Limestone. So that the Collyweston Slate occurs both at the foot and at the top of the escarpment, with a difference of level of some 150 feet. (See Plate X. fig. 1.)

A cross fault has divided the sunken mass; for, in a section at the back of the Midland Railway Station (levelled out of the side of the hill), the Lincolnshire Limestone is seen in lateral juxtaposition with the Ferruginous beds of the Northampton Sand. From an excavation in the Station-yard, I obtained, from a calcareous band in the latter, fragments of a zone containing numerous bivalves, the hollows of which being filled with calcite offer a sparkling contrast to the ferruginous matrix—an effect exactly paralleled by the *Astarte-elegans* zone in ironstone quarries at Harlestone, near Northampton.

The railway passes, by a tunnel under St. Martin's, through the subsided mass of Lincolnshire Limestone, the beds of which have preserved their horizontal position, with little apparent disturbance. At the east end of the tunnel, the railway (very little above the level of the river) passes over beds of the Collyweston Slate; from which, at this point, in 1853, I obtained the beautiful and unique *Astropecten Cotteswoldiae*, var. *Stamfordensis*, described, named, and figured, by Dr. Wright, in his Monograph upon the *Asterioidea*, published in the volume of the Palæontographical Society for 1862.

BURGHLEY PARK.

A mile east of Stamford, on the road to Pilsgate, a view of Burghley House, the palatial residence of the Marquis of Exeter, is obtained through the vista of a fine avenue of trees. In the low intervening ground, a few years ago, a temporary excavation exposed a section, to the depth of 15 feet, in horizontal beds of the Lincolnshire Limestone. At a higher elevation, the House itself, and the almost encircling ornamental sheet of water, are situated upon the clay of the Upper Lias: the ground still rises beyond, and

the beds between the Upper Lias and the Lincolnshire Limestone, in regular sequence, crop out upon the face of the escarpment.

PILSGATE.

In ascending Pilsgate Hill, half-a-mile further east, are passed over, in strange succession at this level (little above the river Welland)—the narrow outcrop of the Upper Estuarine beds, the Great Oolite Limestone (from a quarry in which many years ago I obtained fossils), the Great Oolite Clay, and further on in the village Cornbrash; with which the fault has brought into lateral contact the Lincolnshire Limestone.

BARNACK.

South of the road near Barnack, is Mr. Shelton's quarry, presenting a section in the marly beds of the Lincolnshire Limestone of about 17 feet. At a little distance to the north of this quarry, and at a lower level, is a wide flat of Oxford Clay, which in Barnack, for a considerable distance, is brought into lateral contact with the Upper Lias Clay; and again, on the other side of the Welland, at Uffington, about half-a-mile to the north, the Cornbrash occurs nearly upon the same level.

Fossils from the Cornbrash at Uffington.

Avicula echinata, Sow.
Lima duplicata, Sow.
Ostrea Marshii, Sow. (large).
Pecten, sp.
Rhynchonella concinna, Sow.
 — *varians*, *Schloth.* sp.
Terebratula intermedia, Sow.
 — *lagenalis*, *Schloth.*
 — *sub-lagenalis*, *Dav.*
 — *obovata*, Sow.

Terebratula ornithocephala, Sow.
Chemnitzia vittata, *Phil.* sp.
Ammonites macrocephalus, *Schloth.*
Serpula tetragona, Sow.
Cidaris Bradfordensis, *Wright.*
Echinobrissus clunicularis, *Lhwjd.*
 — *orbicularis*, *Phil.* sp.
 — *quadratus*, *Wright.*
Holactypus depressus, *Leske* (a var.?).

Immediately south of Barnack, the road to Ufford passes through the site of the widely spread quarries whence was obtained the ancient and famous "Barnack Rag". Roman works are known to have been executed in this stone, and Roman coins bestrew the neighbourhood. Several Cathedrals and Abbeys, and most of the early churches throughout a very wide extent of country, are built of this time-enduring stone. A carved fragment from Crowland Abbey, of the 12th century, shows few symptoms of decay. These quarries have been exhausted for more than 400 years, and no section of the ancient beds exists. They may be equivalent to the worthless "Rag" beds of Weldon, Wakerley, and Little Casterton, or to the shelly beds of Ponton: from the stone of the last, the Barnack Rag lithologically is certainly frequently undistinguishable. Mr. Judd informs me, however, that the Barnack Rag immediately overlies the Northampton Sand, and I have found a corresponding shelly bed elsewhere occupying the same position; so that the Barnack Rag is possibly situated quite low down in the general section of the Lincolnshire Limestone, its peculiar character being attributable

(like the many other extreme variations in the beds of the formation) to local and exceptional conditions.

Many years ago, in the carrying out of some agricultural drainage works in fields immediately west of the old quarries, a shallow trench was excavated in the Barnack Rag; from which source, Mr. Bentley and I obtained rock specimens, and numerous, generally minute, but frequently well-preserved, fossils.

From Walcot, a hamlet to Barnack, I obtained years ago, a fine and exceptionally symmetrical specimen of Coral—*Thamnastræa concinna*, Edw. & Haime; probably an undescribed variety*.

Fossils from the "Barnack Rag," Barnack.

Anomia (Placunopsis) semistriata, <i>Bean.</i>	Opis similis, <i>Desh.</i> (or new sp.?).
Avicula Braamburiensis, <i>Phil.</i>	— smooth new sp.?
— echinata, <i>Sow.</i>	Tancredia, sp.?
Gervillia ornata, <i>Lycett.</i>	Trigonia costata, <i>Park.</i> , var. pullus, <i>Mor. & Lyc.</i>
Gryphæa mima, <i>Phil.</i>	Unicardium depressum, <i>Mor. & Lyc.</i>
Hinnites velatus, <i>Goldf.</i> sp.	— parvulum, <i>Mor. & Lyc.</i>
Lima bellula, <i>Mor. & Lyc.</i>	— varicosum, <i>Sow.</i>
— cardiiformis, <i>Mor. & Lyc.</i>	
— duplicata, <i>Sow.</i>	Rhynchonella Crossii, <i>Walker.</i>
— gibbosa, <i>Sow.</i>	— quadruplicata, <i>Ziet.</i>
— Pontonis, <i>Lycett.</i>	— spinosa, <i>Schloth.</i> sp.
— semicircularis (?), <i>Goldf.</i>	— (young form).
Ostrea flabelloides, <i>Lam.</i>	Terebratula fimbria, <i>Sow.</i> (young).
— grogaria, <i>Sow.</i>	— perovalis, <i>Sow.</i>
— sulcifera, <i>Phil.</i>	— simplex, <i>Buckm.</i>
—, sp.?	— sub-maxillata, <i>Sow.</i>
Pecten aratus, <i>Waagen.</i>	
— articulatus, <i>Schloth.</i>	Actæonina parvula, <i>Römer</i> , sp.
— lens, <i>Sow.</i>	—, sp.?
— personatus, <i>Goldf.</i>	Alaria armata, <i>Mor. & Lyc.</i>
— vagans, <i>Sow.</i> , var. peregrinus, <i>Mor. & Lyc.</i>	— hamulus, <i>Deslong.</i> sp.
Perna quadrata, <i>Phil.</i>	— hamus, <i>Deslong.</i> sp.
Pteroperna plana, <i>Mor. & Lyc.</i>	— trifida, <i>Phil.</i> sp.
	Ceritella acuta, <i>Mor. & Lyc.</i> , var.
Arca æmula, <i>Phil.</i>	— parvula, <i>Römer</i> , sp.
— Prattii, <i>Mor. & Lyc.</i>	—, (like Sowerbyi, <i>Mor. & Lyc.</i>)?
— pulchra, <i>Sow.</i>	Cerithium Beanii, <i>Mor. & Lyc.</i>
Astarte elegans, <i>Sow.</i>	—, sp. (like costigerum, <i>Piette</i>)?
— fimbriata, <i>Walton</i> , M.S.	— gemmatum, <i>Mor. & Lyc.</i>
— minima, <i>Phil.</i>	— limæforme (?), <i>Römer.</i>
— Pontonis, <i>Lycett.</i>	— quadricinctum, <i>Goldf.</i>
—, sp.?	— strangulatum, <i>d'Arch.</i>
Cardium incertum, <i>Phil.</i>	—, sp.?
Cypricardia, sp.?	Chemnitzia Wetherellii, <i>Mor. & Lyc.</i>
Cyprina Loweana, <i>Mor. & Lyc.</i>	Cylindrites brevis, <i>Mor. & Lyc.</i>
Cytherea (Cyprina) dolabra, <i>Phil.</i>	— bullatus, <i>Mor. & Lyc.</i>
Lucina crassa, <i>Sow.</i>	— cylindricus, <i>Lycett.</i>
— despecta, <i>Phil.</i>	— turriculatus, <i>Lycett.</i>
Macrodon Hirsonensis, <i>d'Arch.</i>	—, sp.?
Mytilus lunularis, <i>Lycett.</i>	Delphinula alta (?), <i>Mor. & Lyc.</i>
Opis lunulatus, <i>Sow.</i> sp.	— (Crossostoma) Prattii, <i>Mor. & Lyc.</i>

* See description of this specimen by Professor Duncan, F.R.S., in discussion at the conclusion of this Paper.

Delphinula, sp.?
 Monodonta Lyellii (?), *d'Archiac*.
 Natica Leckhamptonensis, *Lycett*.
 Nerinaea cingenda, *Bronn*.
 ——— Cotteswoldiae, *Lycett*.
 ———, sp. (like Eudesii, *Mor. & Lyc.*, young)?
 ———, sp. (like funiculus, *Deslong.*)?
 ——— Jonesii, *Lycett*.
 ——— punctata, *Voltz*.
 ——— triplicata, *Bronn*.
 ——— Voltzii, *Deslong*.
 ———, sp.?
 Onustus, sp. (like Burtonensis, *Lycett*)?
 Phasianella conica, *Mor. & Lyc*.
 ——— elegans, *Mor. & Lyc*.
 ——— parvula, *Mor. & Lyc*.
 ——— Pontonis, *Lycett*.
 ——— tumidula, *Mor. & Lyc*.
 ——— variata (?), *Lycett*.
 ———, sp.?
 Pleurotomaria, sp. (like composita, *Mor. & Lyc.*)?
 Rissolina obliquata, *Sow.* sp.
 Solarium, sp. (something like varicosum, *Mor. & Lyc.*)?
 Trochus Dunkeri, *Mor. & Lyc*.
 ——— bijugatus (?), *Quenst*.
 ——— Ibbetsoni, *Mor. & Lyc*.
 ——— Leckenbyi, *Mor. & Lyc*.
 ——— monilitectus, *Phil*.
 ———, a var.
 ——— ornatissimus, *d'Orb.*, var. Pontonis, *Morris*.
 ——— spiratus, *d'Arch*.
 ———, var.

Trochus squamiger, *Mor. & Lyc*.
 ———, sp.?
 Turbo bijugatus (?), *Quenst*.
 ——— gemmatus, *Lycett*.
 ——— ornatus (?), *Quenst*. (*Littorina ornata*, *Sow.*)?
 ——— Phillipsii, *Mor. & Lyc*.
 ———, sp.?
 Ammonites Blagdeni (?), *Sow.* (young).
 ——— Murchisonae (?), *Sow.* (young).
 Serpula plicatilis, *Goldf*.
 ——— socialis, *Goldf*.
 ——— sulcata, *Sow*.
 Vermicularia nodus, *Phil*.
 Acrosalenia Lycettii, *Wright*.
 Cidaris Fowleri, *Wright*.
 ——— Wrightii (?), *Desor*.
 Pseudodiadema depressum, *Ag*.
 Pentaerinus subsulcatus (?), *Goldf*. (joints).
 Crustacea (claws).
 Isastræa limitata, *Edw. & Haime*.
 Montlivaltia, sp.?
 Thecosmilia gregaria, *Edw. & Haime*.
 Thamnastræa concinna, *Edw. & Haime* (a remarkable variety)*.
 Hybodus? (dorsal spine).
 Strophodus magnus, *Ag.* (palates).
 ——— subreticulatus, *Ag.* (palates).

In addition to the fossils tabulated above, many other minute specimens have been obtained from the Barnack Rag; which, from their being new or young forms (although frequently well preserved), it has been found impossible to identify with described species.

UFFORD.

In ascending the hill into Ufford, the almost tiring reiterated series of all the beds from the Lias upwards to the Great Oolite Limestone inclusive is again crossed. At less than a mile east of Ufford, and at a very slightly lessened elevation, are quarries in the Lincolnshire Limestone. Under about 16 feet of very oolitic stone, are 2 feet 6 inches of slates and "poltids," which repose upon the Lower Estuarine Sands,—seen to the thickness of 9 feet, disposed in thin seams, exhibiting much variation in colour, and containing horizontal and vertical plant-markings.

HELPESTONE.

At a little distance east of this, and within an area not greatly

* See Professor Duncan's remarks in the "Discussion."

exceeding half-a-mile in radius, the various formations assume positions exceedingly anomalous and complicated. On the high ground of Helpstone Heath, is a large quarry in the Lincolnshire Limestone; which is exposed to the thickness of 18 to 20 feet. All the beds are very oolitic, and one is very much inclined from false-bedding: this is separated, by a horizontal flaggy bed, from another bed as much inclined in the reverse direction. No Slate is found here; but a thin band of blue clay separates the Limestone from the Lower Estuarine Sands beneath.

Immediately north of this quarry, is Knight's brickfield. Here, beneath some 10 or 12 feet of oolitic limestone, and separated from it by a foot of sandy clay, are about 18 feet of the Lower Estuarine Sands—in very thin and variegated layers, much inclined to the north (in the direction of the neighbouring escarpment), and containing the characteristic plant-bed in one of the upper bands. Beneath, are ironstone beds of the Northampton Sand, not exposed, but probably having a thickness of 10 or 12 feet. These overlie the clay of the Upper Lias, which is worked for bricks, and is at a considerable elevation above the neighbouring low lands. A few hundred yards to the north of the brick-pit, and at a slightly lower elevation, a temporary opening, a few years ago, disclosed a very solid and shelly rock of Cornbrash: a little to the south-east of this point again, is the abandoned Old Coppice Green parish pit, in Lincolnshire Limestone, the beds of which are coarsely oolitic and false-bedded; while, at the bottom of the hill, at a much lower level, right and left of the southern entrance to the village of Helpstone, are quarries in the Great Oolite Limestone, which is surmounted by a thin covering of Great Oolite Clay, with layers of *Ostrea subrugulosa* in a normal horizontal position.

Half a mile further east than the Old Coppice Green pit, is the most eastern point of the Lincolnshire Limestone area in this direction; and about two miles further, in the parish of Marholme, or in that of Walton, is the last that is to be seen of the Limestones of the Great Oolite.

In directions, north, north-east, east, and south-east, of this locality, all these Northamptonshire beds dip, and are lost, under the extended field of the Oxford Clay; which constitutes the great low flat of the Lincolnshire, the Cambridgeshire, and the Huntingdonshire Fens.

ST. MARTIN'S, STAMFORD.

On the summit of the hill south of and over-looking Stamford (to which I have already twice alluded), are the Marquis of Exeter's excavations for ironstone, just within the Burghley Park Wall. (See Pl. X. fig. 1).

At the top of the section, in patches, answering to the surface contour, appears the Collyweston Slate, weathered into slate from lying so near the surface: beneath this, are the Lower Estuarine Sands and Clays, having a thickness of from 6 to 7 feet, the lowest band containing vertical plant-markings: immediately under these,

is the "Best Black" ironstone (cellular), then the "Second," together from 4 to 5 feet in thickness: a calcareous band of 6 inches comes in here; and below it, succeed—the "Bottom" ironstone (so called), also cellular, 2 feet; a green ferruginous bed, $1\frac{1}{2}$ foot; and a thin ferruginous band, "full of water," and containing small pebble-like nodules (as in the same bed in different ironstone quarries about Northampton), 9 inches; and under all the Upper Lias. As far as I have been able to ascertain, no fossils have been found in these beds.

Section at Burghley-Park Ironstone-Quarry.

	ft.	in.	ft.	in.
1. Soil and Rubble, with patches of Collyweston Slate at bottom			2	6
2. Lower Estuarine Series—				
a. Sand—pale yellow, becoming redder towards the bottom	5	0		
b. Blue clay, with vertical plant-markings	1	6		
3. Ferruginous Beds—			6	6
a. "Best Black" Ironstone, cellular	2	0		
b. "Second"—less cellular, and more sandy	2	0		
c. Calcareous band	0	6		
d. "Bottom" ironstone, cellular	2	0		
e. Green ferruginous stone, about	1	6		
f. Thin red ferruginous band, with pebble-like nodules (as at Duston and Kingthorpe)	0	9		
			8	9

(The last two beds were "full of water.")

4. Upper Lias Clay.

Within a few hundred yards to the west, are Lumby's Terra-cotta Works. A band in the Lower Estuarine Clays supplies an excellent material (mixed with some other ingredients) for this manufacture, and a very durable cream-coloured *terra-cotta* is produced. Similar clay is found at other places in the same bed, and is largely used in the well-known Terra-cotta Works of Mr. Blashfield of Stamford.

At a quarter of a mile further south, on the roadside opposite Whincup's Farm, is the old stone quarry of the abolished Trustees of the Great North Road. The Lincolnshire Limestone is here seen in section to the depth of 18 feet: it is divided into eight distinct beds, varying in mineral condition; some are marly and others oolitic, those near the bottom having much of the character of Barnack Rag, being coarsely oolitic, and containing numerous small shells.

Section in the Lincolnshire Limestone in the Old-Road Pit, near Whincup's Farm.

	ft.	in.
1. Rubble and broken limestone	1	6
2. Compact cream-coloured marly limestone, in thin layers much broken	3	0
3. Soft white marly limestone, surfaces and angles rounded by weathering (<i>Lima bellula</i> , Mor. & Lyc.)	2	6
4. Hard cream-coloured limestone, rather oolitic	3	0
5. Oolitic limestone, like the "cale" of Collyweston	2	0
6. Soft crumbling "caley" oolite	2	6
7. "Rag" bed—coarse oolite, containing numerous shells, <i>Lucina Wrightii</i> , Oppel, <i>Opis</i> , &c.	1	6
8. Hard oolitic stone—not bottomed	2	6

WHITTERING.

A mile south of Whincup's Farm, the road descends a small valley, crosses the White Water brook upon the Upper Lias, and, after passing for a mile over various beds of the Lincolnshire Limestone, traverses the area of the old "Whittering Pendle" quarries. These were very shallow, and, having fallen into disuse, the old familiar pits have long since been levelled down and ploughed over. The "Whittering Pendle," although it has been considered identical with the Collyweston Slate, is very different in its mineral character, being very hard, crystalline, and sometimes almost cherty in texture. It was excavated in large irregular slabs, varying in thickness from one inch to two inches, and was used, without being squared, for the door-slabs and rough floors of cottages, for back-kitchens, &c.

The fossils gathered from this bed are generally characteristic of the Lincolnshire Limestone; but I must particularly notice a specimen taken by myself from the section nearly thirty years ago, and labelled as a Coral during all that time, but which last July was identified by Professor Phillips, F.R.S., as the spadix or fruit of *Aroides Stutterdi*, Carruth., an Arum-like plant, only previously known, I believe, as occurring in the Stonesfield Slate, and described by Mr. Carruthers in the 'Geological Magazine' for April, 1867.

Fossils from the "Whittering Pendle."

Gervillia, sp. ?
Hinnites abjectus, Phil.
 — *velatus*, Goldf. sp.
Lima cardiiformis, Sow.
 — *impressa*, Mor. & Lyc.
 — *Pontonis*, Lycett.
Pecten aratus (?), Waagen.
 — *lens* (?), Sow. (or new sp. ?).
 — *personatus*, Goldf.
Perna quadrata, Phil.
 — *rugosa*, Goldf.

Pteroperna, sp ?
 —
Lucina Bellona, d' Orb.
 — *Wrightii*, Oppel.
Macrodon Hirsonensis, d' Arch. sp.
Modiola, sp. ?
 —
Belemnites Bessinus, d' Orb.
 —
Aroides Stutterdi, Carruth. (spadix).

WANSFORD, WANSFORD TUNNEL, AND HUNTINGDONSHIRE.

The Lincolnshire Limestone continues to be the surface rock for two miles and a half south of Whittering, and is crossed in that space by two valleys, which deepen down to the Upper Lias. It may be seen in a large quarry at Thornhaugh, and again in the road-cutting on the hill overlooking Wansford and the valley of the Nene; which is here the line of demarcation between Northamptonshire and Huntingdonshire.

The river flows over the Upper Lias (generally covered with alluvium). The famous Haycock Inn and the southern half of the village stand on the Northampton Sand. At a higher level, near Wansford Mill, many years ago, a quarry was opened in the Lincolnshire Limestone, in which, beneath a considerable thickness of marly beds (some blue-hearted), was a zone of very hard crystalline limestone, almost made up of small and comminuted shells: at the

bottom of the section, some feet of the Lower Estuarine Sands were exposed.

Half a mile south of this spot, is a very large quarry, called the "Sheep-pens" quarry. The section here exposes a thickness of about 23 feet of the Lincolnshire Limestone, consisting of some seven beds, which display the usual variable characters; overlying a band of two feet of flaggy limestone, disposed in courses of about 2 inches thickness, and containing numerous plants and *Ferns*. Under this, is a band of about 6 inches of Slate, reposing upon a "reddish sand," Lower Estuarine.

Section at the Sheep-pens Quarry, near Wansford Tunnel.

	ft.	in.
1. Rubble and broken limestone.....	3	6
2. Very hard brown limestone, somewhat crystalline, a capital building-stone, and makes good lime	4	0
3. Soft friable limestone	2	6
4. Compact nearly limestone, having a glistening fracture	1	6
5. Sand	0	6
6. Brownish soft bed	1	6
7. Flaggy blue-hearted limestone	1	0
8. "The hard"—a fossiliferous limestone, having a glistening fracture, and sometimes blue-hearted.....	6	0
9. Hard flaggy limestone, somewhat arenaceous, containing <i>plants</i> and <i>ferns</i> and a few shells, and sometimes blue-hearted, in layers of two inches.....	2	0
10. Slate, in small "potlids"	0	6
11. Ruddy sand—Lower Estuarine.		

At a little more than a quarter of a mile beyond this quarry, is the Wansford Railway Tunnel, which passes for one third of a mile, west and east, through beds of the Lincolnshire Limestone. At its western entrance, are the ancient "Red Stone" or "South Pit" quarries, not now worked. Here the stone is a very red, coarse, shelly oolite: it was formerly much in request for the walls of locks, mill-courses, and other hydraulic work. The fossils obtained from this stone correspond very nearly with those yielded by the Ketton and Casterton freestones, with the addition of *Patella rugosa*, some peculiar *Corals*, and a fine example of *Palæozamia pectinata*.

Fossils from the Old "Red Stone" Quarry, Wansford Tunnel.

Ostrea flabelloides, *Lam.*

— *gregaria*, *Sow.*

Arca Prattii, *Mor. & Lyc.*

Astarte elegans, *Sow.*

Cucullæa oblonga, *Sow.*

Lucina Bellona, *d'Orb.*

— *Wrightii*, *Oppel.*

Homomya crassiuscula, *Mor. & Lyc.*
(*young*).

Myoconcha crassa, *Sow.*

Pholadomya, large sp.?

Trigonia spinulosa (?), *Young & Bird.*

Terebratulæ pervalis, *Sow.*

Natica Leekhamptonensis, *Lycett.*

Nerinea triplicata, *Bronn.*

— *Voltzii*, *Deslong.*

Patella rugosa, *Sow.*

Latimæandra Davidsoni, *Edw. & Haime.*

— *Flemingi*, *Edw. & Haime.*

Thecosmilia gregaria, *M'Coy* (a var.).

Palæozamia pectinata, *Brong.*

Wood.

At the eastern end of the Tunnel, the Lincolnshire Limestone appears in massive beds of hard, compact, marly, blue-hearted limestone. The crown of the Tunnel arch reaches up into the lower beds of the Upper Estuarine Clays, which here, as elsewhere, are characterized by the basal ferruginous band. Many tons of this band (which assumes an almost septarian character in places) were, some years ago, experimentally smelted by the late Marquis of Huntley, and a good quality of pig-iron was produced.

The Great Oolite Limestone surmounts the Upper Estuarine Clays, and forms the surface-rock over the Tunnel; at a short distance south, it is overlain by the Great Oolite Clay; and this, in turn, in less than half a mile in the same direction, by the Cornbrash.

Thus (as I have shown) upon a line between Stamford Field, on the north, and a point half a mile south of the Wansford Tunnel, a distance of about 8 miles, the Lincolnshire Limestone is the prominent surface rock; and at each end of this line, is an escarpment, presenting a perfect sequence of beds from the Cornbrash to the Upper Lias inclusive, in which the distinct individuality of the beds of the Great Oolite Limestone and of the Lincolnshire Limestone is unmistakably demonstrated, and the true place of the latter formation in that sequence as certainly determined.

East of the Wansford Tunnel, on the Huntingdonshire side, and at Castor on the Northamptonshire side, of the river Nene, are quarries in the Lincolnshire Limestone; and from these, I have numerous fossils: those from Castor include *Goniomya literata*, Sow. sp., &c.

Fossils from the Lincolnshire Limestone, South of the Nene, near Wansford, Hunts.

Shelly and Freestone Beds.

Hinnites abjectus, <i>Phil.</i>	Astarte depressa, <i>Mor. & Lyc.</i>
— velatus, <i>Goldf.</i> sp.	— elegans, <i>Sow.</i>
Lima bellula, <i>Mor. & Lyc.</i>	— excavata, <i>Sow.</i> , var. compressiuscula, <i>Mor. & Lyc.</i>
— punctata, <i>Sow.</i>	— minima, <i>Phil.</i>
Ostrea flabelloides, <i>Lam.</i> (young).	— recondita, <i>Phil.</i>
— gregaria, <i>Sow.</i>	— rhomboidalis, <i>Phil.</i> sp. (young).
—, sp.? (young), mimetically marked.	—, sp.?
—, sp.?	Cucullæa oblonga, <i>Sow.</i>
Pecten articulatus, <i>Schloth.</i>	Homomya crassiuscula, <i>Mor. & Lyc.</i> (young).
— demissus, <i>Phil.</i>	Lucina Bellona, <i>d'Orb.</i>
—, small smooth sp.?	— despecta, <i>Phil.</i>
—, new sp.?	— Wrightii, <i>Oppel.</i>
Perna rugosa, <i>Goldf.</i> (young).	Macrodon Hirsonensis, <i>d'Arch.</i> sp.
Pteroperna costatula, <i>Deslong.</i> (young).	Modiola gibbosa, <i>Sow.</i>
— plana, <i>Mor. & Lyc.</i> (young).	Myoconcha crassa, <i>Sow.</i>
Trichites nodosus, <i>Lycett.</i>	Mytilus lunularis, <i>Lycett.</i>
	Opis lunulatus, <i>Sow.</i> sp.
	— similis, <i>Desh.</i> (or new sp.?).
Arca æmula, <i>Phil.</i>	—, smooth new sp.?
— Prattii, <i>Mor. & Lyc.</i>	Pholadomya fidicula, <i>Sow.</i>
— pulchra, <i>Sow.</i>	—, large new sp.?
—, sp. (like Kilverti, <i>Mor. & Lyc.</i>)?	—, sp.?
—, sp.?	

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

277

Trigonia compta, *Lycett*.
— *spinulosa* (?), *Young & Bird*.

Rhynchonella Crossii, *Walker*.

—, new sp.?

Terebratula carinata, *Sow.*, or *emarginata*, *Sow.*? (young).

— *fimbria*, *Sow.* (young).

— *ovoides*, *Sow.*

— *perovalis*, *Sow.*

Actæonina glabra, *Phil.*

—, sp.?

—, sp.?

Alaria trifida, *Phil.*

—, sp. (smooth with spines)?

Cerithium Beanii, *Mor. & Lyc.*

— *gemmatum*, *Mor. & Lyc.*

— *quadrincinctum*, *Goldf.*

Cylindrites bullatus, *Mor. & Lyc.*

— *turriculatus*, *Lycett*.

Delphinula (*Crossostoma*) *Prattii*,
Mor. & Lyc. (a var.).

Monodonta Lyellii (?), *d'Arch.*

—, sp. (like *Labadyei*, *d'Arch.*)?

Natica Leckhamptonensis, *Lycett*.

Nerinea Cotteswoldiæ, *Lycett*.

— *gracilis*, *Lycett*.

— *Jonesii*, *Lycett*.

— *punctata*, *Voltz*.

Nerinea triplicata, *Bronn*.

— *Voltzii*, *Deslong*.

Patella rugosa, *Sow.*

Phasianella conica, *Mor. & Lyc.*

Solarium, sp. (like *varicosum*, *Mor. & Lyc.*)?

Trochus Dunkeri (?), *Mor. & Lyc.*

— *Ibbetsoni*, *Mor. & Lyc.*

— *Leckenbyi*, *Mor. & Lyc.*

— *monilitectus*, *Phil.*

— *spiratus*, *d'Arch.*

— (a var.).

—, new sp.

Turbo bijugatus (?), *Quenst.*

— *gemmatum*, *Lycett*.

— *ornatus* (?), *Quenst.*

— *Phillipsii*, *Mor. & Lyc.*

—, sp.?

Serpula sulcata, *Sow.*

Acrosalenia Lycetti, *Wright*.

Latimæandra Davidsoni, *Edw. & Haime*.

— *Flemingi*, *Edw. & Haime*.

Thecosmilia gregaria, *M'Coy* (a var.).

Acrodus or *Hybodus*? (tooth).

Strophodus subreticulatus, *Ag.* (palate).

Marly Beds.

Avicula Münsteri, *Goldf.*

Lima cardiiformis, *Sow.*

— *duplicata*, *Sow.*

— *Pontonis*, *Lycett*.

— *Rodburgensis*, *Lycett*, *M.S.*

— *semicircularis* (?), *Goldf.*

—, sp.?

Pecten demissus, *Phil.*

—, var. *Gingensis*, *Quenst.*

Pinna cuneata, *Phil.*

Ceromya Bajociana, *d'Orb.*

— *similis*, *Lycett*.

Cucullæa cancellata, *Phil.*

Macrodon Hirsonensis, *d'Arch.* sp.

Modiola Sowerbyana, *d'Orb.*

Rhynchonella plicatella, *Sow.*

Terebratula perovalis, *Sow.*

— *sub-maxillata*, *Sow.*

Natica Leckhamptonensis, *Lycett*.

— (large variety, with beautifully punctated surface).

Annulated annelide tube*.

Serpula intestinalis, *Phil.*

Crustacean claw.

Isastræa limitata, *Edw. & Haime*.

Thamnastræa, sp.?

Hybodus (dorsal spine).

Coniferous wood.

ALWALTON TO PETERBORO'—STILTON.

The southern escarpment of the Nene Valley, between Chesterton and Woodstone (half a mile south-west of Peterboro') exposes beds of Cornbrash and Great Oolite Clay and Limestone. The last is traversed in places by a shelly and crystalline band, the stone of which

* Found also in Great Oolite Limestone of Blisworth, in Ironstone beds of Northampton Sand at Duston, and near Banbury.

takes a good polish, and, having been quarried at Alwalton from ancient times, has been called "Alwalton Marble." The long slender disengaged shafts which decorate the exquisite and unique Early English front of Peterboro' Cathedral (dating from the 13th century) were originally composed of this "marble."

From these beds, and from a small patch of Cornbrash near Stilton, I have obtained the following fossils, those from the Cornbrash being in exceedingly fine preservation :—

Cornbrash Fossils.

Anomia semistriata, *Bean*.
Avicula echinata, *Sow*.
Lima duplicata, *Sow*. sp.
 — *impressa*, *Mor.* & *Lyc*.
 — *læviuscula*, *Sow*. sp.
 — *pectiniformis*, *Schloth.* sp.
 — *rigida*, *Sow*. sp.
 — *rigidula*, *Phil.* sp.
Ostrea (*Exogyra*) *lingulata*, *Walton*,
 M.S.
 — *Marshii*, *Sow*.
 —, large flat smooth sp.
 —, several examples of different sp.
 (mimetically marked by contact with
Trigonia).
Pecten anisopleurus, *Buv*.
 — *articulatus*, *Schloth.*
 — *annulatus*, *Sow*.
 — *demissus*, *Phil.*
 — *inæquicostatus*, *Phil.*
 — *lens*, *Sow*.
 — *Michelensis*, *Buv*.
 — *vagans*, *Sow*.
 —, (a var.).
 —, new sp. ?
 —, new sp. ?

Cardium cognatum, *Phil.*
Cypricardia caudata, *Lycett*.
Goniomya V-scripta, *Sow*. sp.
Homomya crassiuscula, *Mor.* & *Lyc*.
 — *gibbosa*, *Sow*. sp.
Isocardia tenera, *Sow*.
Lucina striatula, *Buv*.
Modiola gibbosa, *Sow*.
 — *imbricata*, *Sow*.
 — *Lonsdalei*, *Mor.* & *Lyc*.
 — *Sowerbyana*, *d'Orb*.
Myacites calceiformis, *Phil.* sp.
 — *decurtata*, *Goldf.* sp.
 — *recurva*, *Phil.* sp.
 — *securiformis*, *Phil.* sp.
 — *sinistra*, *Ag.* sp.
 —, new sp. ?
Pholadomya acuticosta, *Sow*.

Pholadomya deltoidea, *Sow*.
 — *lyrata*, *Sow*.
 — *Phillipsii*, *Morris* *.
Trigonia Scarburgensis, *Lycett*.

Rhynchonella Morieri, *Dav*.
 — *concinna*, *Sow*.
 — *obsoleta*, *Sow*.
 — *varians*, *Schloth*.
Terebratulæ Bentleyi, *Morris*.
 —, a variety.
 — *coarctata*, *Park* †.
 — *intermedia*, *Sow*.
 — *lagenalis*, *Schloth*.
 — *sub-lagenalis*, *Dav*.
 — *maxillata*, *Sow*.
 — *obovata*, *Sow*.
 — *ornithocephala*, *Sow*.
 —, undescribed sp.

Bulla, large new sp.
Chemnitzia simplex, *Mor.* & *Lyc*.
 — *vittata*, *Phil*.
 —, sp. ?
Dentalium, new sp.
Pleurotomaria granulata, *Sow*. sp.

Ammonites Herveyi, *Sow*.
 — *macrocephalus*, *Schloth*.
 — *modiolaris*, *Lhwyd*.
Belemnites, new sp. ?

Serpuia intestinalis, *Phil*.
 — *squamosa*, *Bean*.
 — *tetragona*, *Sow*.

Clypeus Mülleri, *Wright*.
Echinobrissus clunicularis, *Lhwyd*.
 — *orbicularis*, *Phil.* sp.
Holætypus depressus, *Leske*.

Ichthyosaurus—teeth, vertebrae, and
 other bones.
 —, small individual—jaws and
 bones.

* One large example has a brilliant nacre.

† Found in Great Northern Railway cutting north of Peterboro' (Northamptonshire). Presented to me by Mr. Bentley, Feb. 5, 1873.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

279

Plesiosaurus—vertebræ and other bones.
Teleosaurus—jaws and teeth, dorsal scute, vertebræ, &c.
Numerous Crocodilian and other Saurian teeth.

Asteracanthus verrucosus, *Eger*.—large dorsal spine.
Lepidotus—scales.
Pycnodus Bucklandi, *Ag*.—teeth.
Strophodus magnus, *Ag*.—palates.
—— subreticulatus, *Ag*.—palates.
Coniferous wood.

Great Oolite Fossils.

Avicula echinata, *Sow*.
—— Münsteri, *Goldf*.
Lima impressa, *Mor. & Lyc*.
—— semicircularis, *Goldf*.
Ostrea Sowerbyi, *Mor. & Lyc*.
—— sub-rugulosa, *Mor. & Lyc*.
——, sp. ?
Pecten annulatus, *Sow*.
—— demissus, *Phil*.
—— lens, *Sow*.
——, large species ?
Pteroperna costatula, *Deslong*.
Astarte, small sp. ?
Cardium Buckmani, *Mor. & Lyc*.
—— cognatum, *Phil*.
Gresslya peregrina, *Phil*.
Homomya crassiuscula, *Mor. & Lyc*.
—— Vezelayi, *d'Arch*. sp.
Isocardia, sp. ?
Modiola imbricata, *Sow*.
—— Sowerbyana, *d'Orb*.
Myacites decurtata, *Phil*. sp.
Pholadomya deltoidea, *Sow*.
—— lyrata, *Sow*.
—— Phillipsii, *Morris*.
—— socialis, *Mor. & Lyc*.
——, new sp. ? with irregular costæ *.
Trigonia Moretoni, *Mor. & Lyc*.
——, sp. ?
Unicardium, sp. ?
Rynchonella concinna, *Sow*.
—— obsoleta, *Sow*.
Terebratulula intermedia, *Sow*.
—— maxillata, *Sow*.
—— ornithocephala, *Sow*. (or new sp. ?).

Natica, sp. ?
Nerita minuta, *Sow*.
Pleurotomaria, sp. ?
Ammonites bullatus, *d'Orb*.
Belemnites, sp. ?
Nautilus Baberi, *Mor. & Lyc*.
—— hexagonus (?), *Sow*.
—— subtruncatus, *Mor. & Lyc*.
Serpula oblique-striata, *Lycett*.
——, sp. ?
Clypeus Mülleri, *Wright*.
Echinobrissus clunicularis, *Lhwjd*.
—— Griesbachii, *Wright*.
—— Woodwardi, *Wright*.
Holectypus depressus, *Leske*.
Glyphea rostrata, *Phil*.
Crocodilian teeth.
Ichthyosaurus—vertebræ and teeth.
Plesiosaurus—vertebræ and other bones.
Hybodus dorsalis, *Ag*.—large spine.
——, sp. ?—dorsal spine.
Lepidotus, sp. ?—jaw.
——, sp. ?—large scales.
Pycnodus Bucklandi, *Ag*.—palates and teeth.
Strophodus magnus, *Ag*.—palates.
—— subreticulatus, *Ag*.
Wood.

A little south of Castor, the Lincolnshire Limestone is only thinly shown; and a little further south, the Upper Estuarine and the Lower Estuarine have once more come together, without the intervention of the Lincolnshire Limestone, which has thinned away and disappeared: the same phenomena are repeated at the opposite, the Huntingdonshire side of the river, at Water Newton; and here, therefore, we may place the limit of the Eastern extension of the formation.

* Found also at Essendine and Danes' Hill, and near Northampton—all in Great Oolite Limestone.

SOUTHERN ESCARPMENT OF THE NENE VALLEY.

West of Wansford Tunnel, near Elton, the Lincolnshire Limestone, in like manner, has disappeared from the southern escarpment of the Nene Valley; but this has received the addition of the Oxford Clay (based by the Kelloway Rock) as the summit bed, and continues unaltered as to its constituent beds (represented in the diagram of vertical sections *) as far as Higham Ferrers (18 miles); near to which town, its upper portion, consisting of the Oxford Clay, Kelloway Rock, and Cornbrash, trends southward, by Rushden and Wymington, into Bedfordshire, and, crossing the valley of the Ouse, passes on into Buckinghamshire, and away.

OUNDLE.

At Oundle (the eastern extremity of my horizontal section †), the escarpment upon either side of the Nene valley is precisely the same. No trace of the Lincolnshire Limestone is found, although at Weldon, only six miles distant due west, it has a thickness exceeding thirty feet; so that the thinning away of that formation in the short intervening space must have been very rapid.

The Oxford Clay caps the high ground of that intervening space, and extends over a considerable tract west and south-west of the same; while at Benefield, three miles west of Oundle, and also on the line of my horizontal section, the Cornbrash and great Oolite beds are also accessible.

Fossils from Benefield.

CORNBRASH.

Rhynchonella Morieri, *Dav.*

Terebratula lagenalis, *Schloth.*
— *obovata*, *Sow.*

GREAT OOLITE.

Cyprina Loweana, *Mor. & Lyc.*
Isocardia tenera, *Sow.*
Modiola imbricata, *Sow.*
Pinna ampla, *Sow.*

Terebratula intermedia, *Sow.*
— *maxillata*, *Sow.*
Clypeus Müllerii, *Wright.*

In a quarry in the Great Oolite Limestone near Oundle, long since closed, were formerly obtained specimens of a beautiful little *Ophiurella*, named by Dr. Wright, *O. Griesbachii*, in compliment to the late well-known veteran Northamptonshire geologist of that name.

Section in Great Oolite Limestone, Oundle.

	ft.	in.	ft.	in.
1. "Lime Earth"—a cream-coloured calcareous argillaceous material, sometimes used for mortar without burning...			2	0
2. "Pendle"—very hard and glistening, splits into thin flags or slates (<i>Modiola imbricata</i> , <i>Sow.</i>)			0	6
3. Blue Clay	0	9 to 1	1	0
4. Calcareo-argillaceous bed, full of <i>Ostrea Sowerbyi</i> , <i>Mor.</i> & <i>Lyc.</i> , burnt for lime	1	0 to 1	1	3
5. Very shelly hard limestone, in thin layers			0	9

* See fig. 4, II., page 283.

† Plate X. fig. 2.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

281

	ft.	in.
6. "Best building-stone"—hard crystalline blue-hearted limestone, full of comminuted shells, worked into lintels and sills of windows, quoins, &c.	3	0
7. Marly limestone, full of shells	3	0
8. Very hard blue-hearted crystalline limestone, full of comminuted shells (small <i>Pteroperna</i>), used for road-metal.....	2	6
9. Upper Estuarine Clays.		

Fossils.

<i>Lima cardiiformis</i> , <i>Sow.</i>	<i>Modiola imbricata</i> , <i>Sow.</i>
—, sp. (like, but not, duplicata, <i>Sow.</i>)?	<i>Trigonia Moretoni</i> , <i>Mor. & Lyc.</i>
<i>Ostrea Sowerbyi</i> , <i>Mor. & Lyc.</i>	<i>Rhynchonella concinna</i> , <i>Sow.</i>
— sub-rugulosa, <i>Mor. & Lyc.</i>	<i>Belemnites</i> (small).
<i>Pecten lens</i> , <i>Sow.</i>	<i>Strophodus magnus</i> , <i>Ag. (palate).</i>
<i>Pteroperna</i> , sp.?	<i>Ophiurella Griesbachii</i> , <i>Wright.</i>

THRAPSTONE TO NORTHAMPTON.

North of Thrapstone and Islip, the high ground has a thin capping of Cornbrash; beneath which, upon a western escarpment, crops out the same sequence of beds as that with which I commenced my descriptions, viz.—

Great Oolite Limestone.	
Upper Estuarine Series.	
Lower Estuarine Series,	} Northampton Sand.
Ferruginous Beds,	
Upper Lias Clay	

And the same sequence again and again occurs throughout the district intervening between Islip and Northampton.

Great Oolite Fossils from Thrapstone.

<i>Cypricardia nuculiformis</i> , <i>Römer.</i>	<i>Terebratula intermedia</i> , <i>Sow.</i>
<i>Modiola imbricata</i> , <i>Sow.</i>	— maxillata, <i>Sow.</i>
<i>Ostrea Sowerbyi</i> , <i>Mor. & Lyc.</i>	<i>Echinobrissus clunicularis</i> , <i>Lhwyd.</i>
<i>Rhynchonella concinna</i> , <i>Sow.</i>	

Cornbrash Fossils from Islip.

<i>Cypricardia caudata</i> , <i>Lycett.</i>	<i>Terebratula lagenalis</i> , <i>Schloth.</i>
<i>Modiola Lonsdalei</i> , <i>Mor. & Lyc.</i>	— obovata, <i>Sow.</i>
<i>Myacites decurtata</i> , <i>Goldf. sp.</i>	—, new sp.
— securiformis, <i>Phil. sp.</i>	<i>Echinobrissus orbicularis</i> , <i>Phil. sp.</i>
—, sp.?	

The Ferruginous beds become thicker and richer at Woodford, Cranford, Finedon, Wellingboro', and other neighbouring places. The quarrying of ironstone has greatly increased in Northamptonshire during 1872; a quantity exceeding 20,000 tons being at this time (November, 1872) weekly raised in the county.

CONCLUSION.

The geological phenomena of the numerous localities described have served, I would submit, to establish the following Propositions:—

- I. That the Upper Estuarine Series of Sands and Clays (subject to local denudation) ranges over the whole of the wide reach of country described, and is everywhere immediately subjacent to the Limestone beds of the Great Oolite, when these are present.
- II. That the Lower Estuarine Series has an equally wide spread, and as persistently overlies the more or less Ferruginous beds of the Northampton Sand.
- III. That, throughout the wide range of the Lincolnshire Limestone, the place of that Limestone is immediately below the Upper Estuarine Series (where not denuded), and immediately above the Lower Estuarine Series.
- IV. That, therefore, necessarily, the Limestone of Kingsthorpe and other high localities about Northampton, and the extensive Limestone now known as the Lincolnshire Limestone, are distinct, and belong to different geological periods.

The *palæontological* evidence also necessitates the last Proposition, and determines further the geological period to which each of these distinct formations belongs—and this, not upon the presence of one or a few fossils only, but upon the *general facies* which the organic contents of these formations severally present:—

- I. Organic forms obtained from the Limestone overlying the Upper Estuarine Series in the neighbourhood of Northampton are as a group so identical with those obtained from beds in the same position in the neighbourhood of Stamford, that, if a number of fossils from one locality were asserted to have come from the other locality, and *vice versâ*, the assertion could not be disputed on palæontological grounds; and the general character of these forms would indicate that they, and therefore the beds in which they occur, belong to the Great-Oolite period.
- II. The organic forms obtained from the Lincolnshire Limestone comprise an altogether different group. Few of them are identical with fossils of the Great Oolite Limestone, while many are identical with fossils which occur in the Ferruginous beds of the Northampton Sand; and their general character indicates that they, and therefore the Lincolnshire Limestone, are of the period of the Inferior Oolite.

These Propositions proven, I contend that Physical Geology and Palæontological Geology combine to afford indisputable evidence of the truth of the conclusions which I pre-stated in my Introduction.

Therefore—the General Section of the north-eastern portion of the Northern Division of Northamptonshire, will be represented in the Diagram I., which *includes* the Lincolnshire Limestone; while the General Section of the south-western portion of that Division, will be represented in the Diagram II., which *excludes* that formation*.

* See fig. 4—opposite page.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

283

Fig. 4.—Sections illustrating the Relations of the Lincolnshire Limestone to the Northamptonshire Oolite.

Maximum Thicknesses of Beds within the Area described.

	feet.
Oxford Clay . . . . . uncertain.	
Cornbrash	15
Great Oolite—Clay	20 feet.
„ Limestone	25 „
„ Upper Estuarine	33 „
	78
Inferior Oolite—Lincolnshire Limestone	75 feet.
„ Northampton Sand—	
Lower Estuarine	18 feet.
Ferruginous beds	60 „
	78 „
	153
	246

A few words as to the extent of the area occupied by the Lincolnshire Limestone. It ranges through the whole of the county of Lincoln, stretching into South Yorkshire on the north, and through Rutland into Northamptonshire on the south. The length of this outcrop (three fourths of which is in Lincolnshire) is about 110 miles. In Northamptonshire, its greatest apparent width across the strike is about 16 miles. It attains to its greatest thickness in Lincolnshire, exceeding near Sleaford 200 feet. Mr. Judd (who has “beaten its bounds”) considers that it had originally an elongated lenticular form, and probably thinned away irregularly in every direction from the point of its greatest thickness. In Lincolnshire, its eastern boundary is hidden under superincumbent beds of other formations; and its western boundary, to a very great extent, has been pared away by the denudation which scooped out the great valley of the Witham and the Trent.

It is probable that the Lincolnshire Limestone and the Northampton Sand have no exact analogues elsewhere; but, as the series of these Northamptonshire beds has at its upper limit and lower limit respectively such definite formations as the Oxford Clay and Cornbrash, and the Upper Lias, it may not be impossible approximately to correlate its several members with formations in other parts of the country. I have accordingly drawn up the following Table.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

285

Table showing the Probable Synchronous Relation of the Northamptonshire Beds to Formations in Yorkshire and the South-West of England.

Yorkshire.	Northamptonshire, &c.	S.W. of England.
OXFORD CLAY.	OXFORD CLAY.	OXFORD CLAY.
CORNBRASH.	CORNBRASH.	CORNBRASH.
GREAT OOLITE :— ——— ———	GREAT OOLITE :— Great Oolite Clay. Great Oolite Limestone.	GREAT OOLITE :— Forest Marble. Bradford Clay. Limestones of Great Oolite of Bath, the Cotteswolds, and the Upper Beds at Minchinhampton. Stonesfield Slate.
Upper Plant Shale ?	Upper Estuarine.	
INFERIOR OOLITE :— Grey Limestone of Scarborough.	INFERIOR OOLITE :— Lincolnshire Limestone.	INFERIOR OOLITE :— Zone of <i>Am. Parkinsoni</i> . Zone of <i>Am. Humphreysianus</i> .
Lower Plant Shale. Ferruginous Beds of Glaizedale. The Dogger.	Lower Estuarine. } Ferruginous Beds, Upper Portion. } Ferruginous Beds, Lower Portion. }	— Zone of <i>Am. Murchisonæ</i> *. Zone of <i>Am. opalinus</i> , and the Midford Sand.
UPPER LIAS.	UPPER LIAS.	UPPER LIAS.

The *Great Oolite Clay* is not represented in Yorkshire, but doubtless answers to the Forest Marble and Bradford Clay of the south-west.

The *Great Oolite Limestone*, also, has no representative in Yorkshire, but is probably equivalent to the Great Oolite of the south-west and to the upper beds of Minchinhampton.

The *Upper Estuarine Series*, perhaps, answers to the Upper Plant Shale of Yorkshire, but is more certainly the equivalent of the Stonesfield Slate.

With the *Lincolnshire Limestone*, the series of Inferior Oolite formations comes in. As indicated by the presence of *Am. terebratus*, Phil., it probably is nearly synchronous with the Grey Limestone of

* Dr. Wright, F.R.S.E., F.G.S., in his "Correlation of the Jurassic Rocks," 1869, in a complimentary allusion to my *Stellaster Sharpi*, Wright, describes that fossil as having been discovered in the *Ammonites-Murchisonæ* zone, near Northampton. It was found in the upper portion of the Ironstone section at Duston.

Scarboro', and embraces (as indicated by the occurrence of *Ammonites Murchisonæ*, Sow.) lower beds of the south-west.

The *Lower Estuarine Series* probably tallies with the Lower Plant Shale of Yorkshire, but is not represented in the south-west.

The upper portion of the *Ferruginous Beds* has representatives in the Glazedale beds and the Dogger of Yorkshire, and in the *Ammonites-Murchisonæ* zone of Cleve Hill, &c.

The lower portion of the *Ferruginous Beds* (as indicated respectively by the presence of *Am. opalinus*, Rein., and of *Am. insignis*, Schubler) is probably nearly equivalent to the *Am.-opalinus* zone of Dr. Wright, and to the Midford sand.

I must not omit to acknowledge the information I have so largely derived from, and which on all occasions has been so freely given by, Mr. Judd, and the great aid which his remarkable Map, sheet 64, has afforded me in revising the rough draft of this Memoir. I must also warmly thank him for the assistance he has rendered me in the preparation of my Map and Diagrams.

My warm thanks are also due to Mr. Etheridge, F.R.S., for the time, great labour, and patience he has bestowed in examining my multitude of fossils, in revising and in many cases correcting my identifications, and in determining those of fossils which were beyond my reach; enabling me to construct the appended Comparative Tables of the organic contents of the several formations.

The preparation of this Memoir has not been unattended with labour upon my part; and the data upon which it is founded have been gathered during many years' acquaintance with, and observations in, the districts to which it refers. If I shall have contributed my quota to a knowledge of the beds I have described and of the geology of the county I have traversed, I shall be satisfied with the result.

The whole of the fossils from the neighbourhood of Northampton which I exhibited on the last occasion were collected by me. A considerable proportion of those which illustrate this my Second Part are also of my collection; some were jointly collected by Mr. Bentley and myself, some by Mr. Bentley alone (notably those from Morcot and Denton), and a few by the late Dr. Porter of Peterboro'.

The ear probably may have wearied in listening to my lengthy and somewhat dry exposition; but the eye, meanwhile, may have rested satisfactorily on the fossils which, fortunately, I have been enabled to exhibit—fossils which are the evidence of the richness of the *fauna* of the periods and of the areas to which I have directed attention.

Exalted personages attend at Court on ceremonial occasions, decorated with the respective stars of their several orders. The distinguished geological entities—the Lincolnshire Limestone and the Northampton Sand—have appeared at the Court of this Society, in like manner decorated with the *Astropecten* and the *Stellaster*, the respective and characteristic "stars"—the *Astrum* and the *Stella*—of those formations.

Fossils from the GREAT OOLITE Beds of the Northampton and Stamford Districts*.—Table showing which Forms are contained also in the Beds of the LINCOLNSHIRE LIMESTONE and the NORTHAMPTON SAND.

Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Line. Limestone.	Northampton Sand.	Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Line. Limestone.	Northampton Sand.
LAMELLIBRANCHIATA (<i>Monomyaria</i>).					Pecten				
<i>Avicula</i>					Wollastonensis, <i>Lycett.</i>	*			
<i>echinata, Sow.</i>	*	*	*		Perna				
<i>Münsteri, Goldf.</i>	*	*	*	*	<i>foliacea, Lycett.</i>	*			
<i>Exogyra</i>					(like) <i>mytiloides, Lam.</i>	*			
<i>auriformis, Goldf.</i>	*				<i>quadrata, Sow.</i>	*	*	*	
<i>Gervillia</i>					<i>rugosa, Goldf.</i>	*	*	*	*
<i>crassicosta, Mor. & Lyc. (a var.)</i>	*				<i>new sp. ?</i>	*			
<i>Islipensis, Lycett.</i>	*	*			Pinna				
<i>monotis, Deslong.</i>	*				<i>ampla, Sow.</i>	*			
<i>Lima</i>					<i>cuneata, Phil.</i>	*	*	*	*
<i>cardiiformis, Sow. sp.</i>	*	*	*	*	<i>new sp. ?</i>	*			
<i>duplicata, Phil. sp.</i>	*	*	*	*	Placunopsis				
<i>impressa, Mor. & Lyc.</i>	*	*	*	*	<i>socialis, Mor. & Lyc.</i>	*		*	
<i>semicircularis, Goldf.</i>	*	?			Pteroperna				
<i>Ostrea</i>					<i>costatula, Deslong.</i>	*	*	*	*
<i>acuminata, Sow.</i>	*				<i>emarginata, Mor. & Lyc.</i>	*			
<i>costata, Sow.</i>	*			*	<i>plana, Mor. & Lyc.</i>	*	*	*	*
<i>flabelloides, Lam.</i>	*		*	*	<i>new sp. (finely costated) ?</i> ...	*			
<i>Sowerbyi, Mor. & Lyc.</i>	*	*			LAMELLIBRANCHIATA				
<i>sub-rugulosa, Mor. & Lyc.</i>	*	*	?		(<i>Dmyaria</i>).				
<i>large flat sp. ?</i>	*	*	*	*	Anatina				
Pecten					<i>plicatella, Mor. & Lyc.</i>	*			
<i>annulatus, Sow.</i>	*	*			<i>undulata, Sow.</i>	*			
<i>articulatus, Schloth.</i>	*	*	*	*	Arca				
<i>demissus, Phil.</i>	*	*	*	*	<i>Eudesii, Mor. & Lyc.</i>	*			
<i>Griesbachii, Lycett.</i>	*				<i>rugosa, Mor. & Lyc.</i>	*			
<i>lens, Sow.</i>	*	*	*	*	Astarte				
— (or large new sp.) ? ...	*	*	*	*	<i>angulata, Mor. & Lyc.</i>	*			
<i>retiferus (?)</i> , <i>Mor. & Lyc.</i>	*				<i>depressa, Goldf.</i>	*	*	*	*
— <i>rigidus, Sow.</i>	*				<i>sp. (like elegans, Sow.) ?</i> ...	*			

* *Errata* in the Table published with the First Part of this Memoir have been corrected in this Table, and the names of some Fossils since obtained have been added.

Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.	Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.
Cardium					Lucina				
Buckmani, <i>Mor. & Lyc.</i> ...	*	*	*	*	crassa, <i>Sow.</i>	*			
cognatum, <i>Phil.</i>	*	*	*	*	Macrodon				
incertum, <i>Phil.</i>	*	*	*	*	Hirsonensis, <i>d'Arch. sp.</i> ...	*	*	*	*
lingulatum, <i>Lycett</i>	*	*			— var. rugosa, <i>Lycett</i> ...	*			
Stricklandi, <i>Mor. & Lyc.</i> ...	*	*			Modiola				
subtrigonum, <i>Mor. & Lyc.</i> ...	*	*	*	*	cuneata, <i>Sow.</i>	*	*	*	*
Ceromya					gibbosa, <i>Sow.</i>	*	*	*	*
concentrica, <i>Sow.</i>	*	*	*	*	imbricata, <i>Sow.</i>	*	*	*	*
plicata, <i>Ag. sp.</i>	*				Lonsdalei, <i>Mor. & Lyc.</i>	*	*	?	?
Symondsii, <i>Mor. & Lyc.</i>	*	*			Sowerbyana, <i>d'Orb.</i>	*	*	*	*
undulata, <i>Mor. & Lyc.</i>	*				subreniformis, <i>Mor. & Lyc.</i> ..	*	*	?	?
Corbis (Corbicella)					tenuistriata, <i>Münst. sp.</i>	*			
Bathonica, <i>Mor. & Lyc.</i>	*	*			Myacites				
Cucullæa					Beanii, <i>Mor. & Lyc.</i>	*			
concinna, <i>Phil.</i>	*	*			calceiformis, <i>Phil. sp.</i>	*	*	*	
cucullata, <i>Goldf.</i>	*	*	*	*	compressus, <i>Mor. & Lyc.</i> ...	*			
triangularis, <i>Phil.</i>	*	*			decurtata, <i>Phil. sp.</i>	*	*	*	
large sp. ?	*	*			securiformis, <i>Phil. sp.</i>	*	*	*	
Cypicardia					Terquemea, <i>Buv. sp.</i>	*	*	*	
Bathonica, <i>d'Orb.</i>	*	*	*	*	tumida, <i>Mor. & Lyc.</i>	*			
caudata, <i>Lycett.</i>	*	*			sp. ?	*	*		
nuculiformis, <i>Römer</i>	*	*	*	*	Mytilus				
rostrata, <i>Sow.</i>	*				asper, <i>Sow.</i>	*			
Cyprina					tumidus, <i>Mor. & Lyc.</i>	*			
Davidsoni, <i>Lycett.</i>	*	*			Nerera				
depressiuscula, <i>Mor. & Lyc.</i> ...	*				Ibbetsoni, <i>Morris</i>	*	*		
Islipensis, <i>Lycett.</i>	*				Nucula				
Loweana, <i>Mor. & Lyc.</i>	*	*	*	*	Menkei, <i>Römer</i>	*			
— var. elongata, <i>Mor. & Lyc.</i> ..	*	*			Waltoni, <i>Mor. & Lyc.</i>	*			
trapeziformis, <i>Römer</i>	*	*	*	*	Opis				
Cyrena, sp. ?	*				new sp. ?	*	*		
Goniomya					Pholadomya				
hemicosta, <i>Mor. & Lyc.</i>	*				acuticosta, <i>Sow.</i>	*	*		
Gresslya					deltoides, <i>Sow.</i>	*	*		
peregrina, <i>Phil.</i>	*	*			Heraulti, <i>Ag.</i>	*	*	*	*
Homomya (Myacites)					lyrata, <i>Sow.</i>	*	*	*	*
crassiuscula, <i>Mor. & Lyc.</i> ...	*	*	*	?	oblita, <i>Mor. & Lyc.</i>	*	*		
unioniformis, <i>Mor. & Lyc.</i> ...	*	*	*	*	Samanni, <i>Mor. & Lyc.</i>	*			
Vezelayi, <i>d'Arch. sp.</i>	*	*	*	?	socialis, <i>Mor. & Lyc.</i>	*	*		
Isocardia					solitaria, <i>Mor. & Lyc.</i>	*	*		
nitida, <i>Phil.</i>	*	*			new large sp. with coarse <i>costæ</i>	*	*		
tenera, <i>Sow.</i>	*	*			new round sp. with irregular <i>costæ</i>	*	*		
new sp. ?	*	*			several undetermined sp. ...	*	*		
Limopsis					Pholas				
oolitica, <i>d'Arch.</i>	*	*			oolitica, <i>Mor. & Lyc.</i>	*	?		
Lithodomus					pulchralis, <i>Bean.</i>	?			
inclusus, <i>Phil.</i> (very small)	*	*	*	*	Quenstedtia				
Lucina					lavigata, <i>Mor. & Lyc.</i>	*		*	
Bellona, <i>d'Orb.</i>	*	*	*	*	oblita, <i>Phil., sp.</i>	?		*	*

Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.	Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.
Sphæra					Dentalium				
<i>Madridi, d'Arch. sp.</i>	*	*			<i>sp. ?</i>	*			
Tancredia					Natica				
<i>angulata, Lycett</i>	*	*	*	*	<i>formosa, Mor. & Lyc.</i>	*	*	*	
<i>axiniformis, Phil. sp.</i>	*	*	*	*	<i>globosa, Römer</i>	*	*	*	
<i>planata, Mor. & Lyc.</i>	*	*	*	*	<i>grandis, Goldf.</i>	*	*	*	
<i>truncata, Lycett</i>	*	*			<i>intermedia, Mor. & Lyc.</i>	*	*	*	
Tellina, sp. ?	*	*			<i>Michelini, d'Arch.</i>	*	*	*	
Thracia					<i>neritoides, Mor. & Lyc.</i>	*			?
<i>amygdaloides, Lycett</i>	*	*			<i>texata (?), Lycett</i>	*			
<i>curtansata, Mor. & Lyc.</i>	*	*			<i>Verneuilii, d'Arch.</i>	*			?
Trigonia					<i>(Euspira) canaliculata, Mor.</i>				
<i>costata, Park.</i>	*	*	*	*	<i>& Lyc.</i>	*	*		
<i>—, var. pullus, Mor. & Lyc.</i>	*	*			<i>— coronata, Mor. & Lyc.</i>	*			
<i>—, var. elongata, Mor. & Lyc.</i>	*	*			<i>— pyramidata, Mor. & Lyc.</i>	*	*		
<i>Lyc.</i>	*				<i>— Sharpei, Mor. & Lyc.</i>	*	*		
<i>Crucis, Lycett.</i>	*				Nerinæa				
<i>Goldfussi, Ag.</i>	?				<i>funiculus, Deslong.</i>	*			
<i>Moretoni, Mor. & Lyc.</i>	*	*			<i>punctata, Voltz</i>	*		*	
Unicardium					<i>Stricklandi, Mor. & Lyc.</i>	*		*	
<i>impressum, Mor. & Lyc.</i>	*	*	*	*	<i>Voltzii, Deslong.</i>	*		*	
<i>parvulum, Mor. & Lyc.</i>	*	*	*	*	<i>sp. ?</i>	*	*		
<i>varicosum, Sow. sp.</i>	*	*			Nerita				
BRACHIOPODA.					<i>hemisphærica, Römer</i>	*			?
Rhynchonella					<i>minuta, Sow.</i>	*			
<i>concinna, Sow. sp.</i>	*	*			Phasianella				
<i>Hopkinsi (?), M' Coy, sp.</i>	*				<i>acutiuscula, Mor. & Lyc.</i>	*		*	
<i>obsoleta, Sow. sp.</i>	*	*			<i>elegans, Mor. & Lyc.</i>	*		*	
Terebratula					Pleurotomaria				
<i>digona, Sow.</i>	*	*			<i>armata, Münst.</i>	*		*	*
<i>globata (?), Sow. (a var.) ?</i>	*				<i>scalaris, Deslong.</i>	*		*	
<i>intermedia, Sow.</i>	*	*			<i>sp. ?</i>	*		*	
<i>maxillata, Sow.</i>	*	*			Trochotoma				
<i>obovata, Sow.</i>	*	*			<i>tabulata, Mor. & Lyc.</i>	*	*	*	*
<i>ornithocephala, Sow. (or new sp. ?)</i>	*	*			Trochus ?	*			
GASTEROPODA.					CEPHALOPODA.				
Alaria					Ammonites				
<i>armata, Mor. & Lyc.</i>	*	*	*	*	<i>bullatus, d'Orb.</i>	*		*	
<i>trifida, Phil. sp.</i>	*	*	*	*	<i>gracilis, Buckm. (large)</i>	*		*	
Amberleya					<i>macrocephalus, Schloth.</i>	*		*	
<i>nodosa, Mor. & Lyc.</i>	*	*			<i>sp. ?</i>	*		*	
Bulla					<i>small sp. ?</i>	*		*	
<i>undulata, Bean</i>	*				<i>large sp.</i>	*		*	
<i>Cerithium, sp. ?</i>	*				Belemnites				
<i>Chemnitzia</i>					<i>Bessinus, d'Orb.</i>	*	*	*	*
<i>Hamptonensis (?), Mor. & Lyc.</i>	*				<i>sp. (ungrooved ?)</i>	*	*		
<i>Delphinula, sp. ?</i>	*				Nautilus				
					<i>Baberi, Mor. & Lyc.</i>	*	*		

Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.	Great-Oolite Fossils.	Gt. O., Northampton.	Gt. O., Stamford.	Linc. Limestone.	Northampton Sand.
<i>Nautilus</i>					<i>Isastræa</i>				
<i>hexagonus</i> (?), <i>Sow.</i>	*				<i>gibbosa</i> (?), <i>Duncan</i>	*			
<i>latidorsatus</i> , <i>d'Orb.</i>	*				POLYZOA.				
<i>subtruncatus</i> , <i>Mor. & Lyc.</i> ...	*				<i>sp.?</i>	*	*		
ARTICULATA.					CRUSTACEA.				
<i>Serpula</i>					<i>Eryma</i>				
<i>oblique-striata</i> , <i>Lycett</i>	*	*			<i>sp.</i> (allied to <i>elegans</i> ,				
<i>socialis</i> , <i>Goldf.</i>	*	*	*	*	<i>Oppel</i>)?	*	*		
<i>Annulated Annelide tubes</i> † ..	*	*	*	*	<i>Glyphea</i>				
ECHINODERMATA.					<i>rostrata</i> , <i>Phil. sp.</i>	*			
<i>Aerosalenia</i>					PISCES.				
<i>hemicidaroides</i> , <i>Wright</i>	*	*			<i>Ilybodius</i>				
<i>pustulata</i> , <i>Forbes</i>	*	*			<i>dorsalis</i> , <i>Ag.</i> —spines	*	*	*	
<i>spinosa</i> , <i>Ag.</i>	*			?	—, or <i>sp.?</i> —large spines..	*	*	*	
<i>Wiltoni</i> , <i>Wright</i>	*	*			<i>Lepidotus</i>				
<i>Clypeus</i>					<i>sp.?</i> —jaw, teeth, and scales	*	*		
<i>Mülleri</i> , <i>Wright</i>	*	*			<i>Pholidophorus</i>				
<i>Plottii</i> , <i>Klein</i>	*				<i>Flesheri</i> , <i>Ag.</i>	*			
<i>Echinobrissus</i>					<i>Pycnodus</i>				
<i>clunicularis</i> , <i>Llwyd</i>	*	*	*	*	<i>Bucklandi</i> , <i>Ag.</i> —palate and				
<i>Griesbachii</i> , <i>Wright</i>	*	*			teeth	*	*	*	
<i>sp.</i> (like <i>orbicularis</i> , <i>Phil.</i>					<i>Strophodus</i>				
<i>sp.</i>)?	*				<i>magnus</i> , <i>Ag.</i> —palates	*	*	*	
<i>Woodwardii</i> , <i>Wright</i>	*	*			<i>subreticulatus</i> , <i>Ag.</i> —palates.	*	*	*	
<i>Hemicidaris</i>					REPTILIA.				
<i>minor</i> (?), <i>Ag.</i>	*				<i>Crocodylia</i>				
<i>Holactypus</i>					<i>numerous perfect teeth</i>	*	*	*	
<i>depressus</i> , <i>Lesle</i>	*	*			<i>Ichthyosaurus</i>				
<i>Ophiurella</i>					<i>sp.?</i> —vertebræ and teeth	*	*		
<i>Griesbachii</i> , <i>Wright</i> (Oundle)...	*				<i>Plesiosaurus</i>				
<i>Pentacrinus</i>					<i>sp.?</i> —vertebræ	*			
<i>stem and branch joints</i>	*				<i>Telcosaurus</i>				
ZOOPHYTA.					<i>sp.?</i> (large)—bones, verte-				
<i>Anabacia</i>					bræ, teeth, and dorsalscutes	...	*		
<i>orbulites</i> , <i>Lamar. sp.</i>	*	...	*		<i>sp.?</i> (small)—jaws, scute,				
<i>Calamophyllia</i>					teeth, vertebræ, atlas	*	*	*	
<i>radiata</i> , <i>Lamar. sp.</i>	*	...	*		PLANTÆ.				
<i>Cladophyllia</i>					<i>Carpolithes</i> (fruit)	*	...	...	*
<i>Babeana</i> , <i>Edw. & Haime</i> ...	*	...	*		<i>Kaidacarpum</i>				
<i>Cyathophora</i>					<i>ooliticum</i> , <i>Carruth</i>	*			
<i>tuberosa</i> , <i>Duncan</i>	*				<i>Plants in vertical position</i> ...	*	...	...	*
<i>Isastræa</i>					<i>Wood</i>	*	*	*	*
<i>limitata</i> , <i>Lamar.</i>	*	...	*						

† Found also near Banbury.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

291

Fossils from the LINCOLNSHIRE LIMESTONE.—Table showing which Forms are contained also in the Beds of the NORTHAMPTON SAND and of the GREAT OOLITE of the Northampton and Stamford Districts.

Lincolnshire Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.	Lincolnshire Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.
LAMELLIBRANCHIATA (<i>Monomyaria</i>).			<i>Ostrea</i>		
<i>Anomia</i>			<i>sulcifera</i> , <i>Phil.</i>	*	
smooth sp. ?			large flat sp. ?	*	*
(<i>Placunopsis</i>) <i>semi-striata</i> , <i>Bean.</i>			<i>Pecten</i>		
<i>Avicula</i>			<i>aratus</i> , <i>Waagen.</i>		
<i>Braamburiensis</i> , <i>Sow.</i>	*		<i>arcuatus</i> , <i>Sow.</i>	*	
<i>clathrata</i> (?), <i>Lycett.</i>			<i>articulatus</i> , <i>Schloth.</i> (a var. ?)	*	*
<i>echinata</i> , <i>Sow.</i>	*	*	<i>clathratus</i> , <i>Römer</i>	*	*
<i>Münsteri</i> , <i>Goldf.</i>	*	*	<i>demissus</i> , <i>Phil.</i>	*	*
<i>sub-costata</i> (?), <i>Römer.</i>			—, var. <i>Gingensis</i> , <i>Quenst.</i>		
<i>Gervillia</i>			<i>lens</i> , <i>Sow.</i>	*	*
<i>acuta</i> , <i>Sow.</i>	*		—, or new large sp. ?	*	*
<i>lata</i> , <i>Phil.</i>	*		<i>personatus</i> , <i>Goldf.</i>	*	
<i>ornata</i> (?), <i>Lycett.</i>			<i>vagans</i> (?), <i>Sow.</i>		
<i>radians</i> , <i>Mor. & Lyc.</i>			—, var. <i>peregrinus</i> , <i>Mor. &</i> <i>Lyc.</i>		
(nearly allied to <i>G. Hart-</i> <i>manni</i> , <i>Goldf.</i>)			<i>Perna</i>		
<i>Gryphæa</i>			<i>quadrata</i> , <i>Sow.</i> sp.	*	*
<i>mima</i> , <i>Phil.</i>	*		<i>rugosa</i> , <i>Goldf.</i>	*	*
<i>Hinnites</i>			<i>Pinna</i>		
<i>abjectus</i> , <i>Phil.</i> sp.	*		<i>cancellata</i> , <i>Bean.</i>		
<i>tegulatus</i> , <i>Mor. & Lyc.</i>			<i>cuneata</i> , <i>Phil.</i>	*	*
<i>velatus</i> , <i>Goldf.</i> sp.	*		<i>Placunopsis</i>		
<i>Inoceramus</i>			<i>ornatus</i> , <i>Mor. & Lyc.</i>		
<i>obliquus</i> , <i>Mor. & Lyc.</i>	*		<i>socialis</i> , <i>Mor. & Lyc.</i>	*	*
<i>Lima</i>			<i>Plicatula</i>		
<i>bellula</i> , <i>Mor. & Lyc.</i>	*		<i>tuberculosa</i> , <i>Mor. & Lyc.</i>	*	
<i>cardiiformis</i> , <i>Sow.</i> sp.	*	*	<i>Pteroperna</i>		
<i>duplicata</i> , <i>Sow.</i> sp.	*	*	<i>costatula</i> , <i>Deslong.</i>	*	*
<i>Etheridgii</i> , <i>Wright.</i>	*	*	<i>plana</i> , <i>Mor. & Lyc.</i>	*	*
<i>gibbosa</i> , <i>Sow.</i> sp.			<i>pygmæa</i> , <i>Mor. & Lyc.</i>		
<i>impressa</i> , <i>Mor. & Lyc.</i>	*	*	<i>Trichites</i>		
<i>Luciensis</i> (?), <i>d'Orb.</i>	*	*	<i>nodosus</i> , <i>Lycett.</i>		
<i>Pontonis</i> , <i>Lycett</i>	*		LAMELLIBRANCHIATA (<i>Dimyaria</i>).		
<i>proboscidea</i> , <i>Sow.</i> sp.	*		<i>Arca</i>		
<i>punctata</i> , <i>Sow.</i> sp.	*		<i>æmula</i> , <i>Phil.</i>	*	
<i>Rodbургensis</i> , <i>Lycett.</i> <i>MS.</i>	*		<i>Prattii</i> , <i>Mor. & Lyc.</i>	*	
<i>semicircularis</i> (?), <i>Goldf.</i>	*	*	<i>pulchra</i> , <i>Sow.</i>	*	
large sp. (allied to <i>L. grandis</i> , <i>Römer</i>) ?	*		<i>Astarte</i>		
large sp. ?	*		<i>depressa</i> , <i>Goldf.</i>	*	*
<i>Ostrea</i>			<i>elegans</i> , <i>Sow.</i>	*	
<i>flabelloides</i> , <i>Lam.</i>	*	*	<i>excavata</i> , <i>Sow.</i>	*	
<i>gregaria</i> , <i>Sow.</i>	*		—, var. <i>compressiuscula</i> , <i>Mor. & Lyc.</i>		
			<i>excentrica</i> , <i>Mor. & Lyc.</i>		

Lincolnshire-Limestone Fossils.	Northampton Sand. Great Oolite, N. & S.	Lincolnshire-Limestone Fossils.	Northampton Sand. Great Oolite, N. & S.
Astarte		Lucina	
<i>finbriata</i> , Walton, M.S.		<i>Bellona</i> , d'Orb.	* *
<i>minima</i> , Phil.	*	<i>despecta</i> , Phil.	* *
<i>Pontonis</i> , Lycett.		—, var. <i>cardioides</i> , d'Arch.	*
<i>recondita</i> , Phil. sp.		<i>rotundata</i> , Römer	*
<i>rhomboidalis</i> , Phil. sp. (Hippopodium Lucienae, d'Orb.)	*	<i>Wrightii</i> , Oppel.	
<i>Wiltoni</i> (?), Mor. & Lyc.		—, sp. ?	
sp. ?		Macrodon	
Cardium		<i>Hirsonensis</i> , d'Arch.	* *
<i>Buckmani</i> , Mor. & Lyc.	* *	Modiola	
<i>cognatum</i> , Phil.	* *	<i>Binfieldi</i> , Mor. & Lyc.	* *
<i>incertum</i> , Phil.	*	<i>cuneata</i> , Sow. sp.	* *
sp. (near to <i>Stricklandi</i> , Mor. & Lyc.) ?	*	— (a <i>Perna</i> -like variety).	
<i>subtrigonum</i> , Mor. & Lyc.	*	<i>gibbosa</i> , Sow.	* *
Ceromya		<i>Sowerbyana</i> , d'Orb.	* *
<i>Bajociana</i> , d'Orb.	*	<i>sublevis</i> (?), Sow.	
<i>concentrica</i> , Sow.	*	Myacites	
<i>similis</i> , Lycett.		<i>calceiformis</i> , Phil. sp.	*
Cucullæa		<i>decurtata</i> , Phil. sp.	*
<i>cancellata</i> , Phil.	*	<i>Scarburgensis</i> , Phil. sp.	
<i>cucullata</i> , Goldf.	* *	<i>securiformis</i> , Phil. sp.	*
<i>elongata</i> , Sow.	*	Myoconcha	
<i>Goldfussii</i> , Römer.		<i>crassa</i> , Sow.	*
<i>imperialis</i> , Phil.	*	Mytilus	
<i>oblonga</i> , Sow.	*	<i>furcatus</i> , Goldf.	*
Cypricardia		<i>lunularis</i> , Lycett.	
<i>Bathonica</i> , d'Orb.	* *	Opis	
<i>nuculiformis</i> , Römer.	* *	<i>gibbosus</i> , Lycett.	
Cyprina		<i>lunulatus</i> , Sow. sp.	*
<i>Jurensis</i> , Goldf. sp.		<i>similis</i> (?), Deslong. (or allied sp. ?)	
<i>Loweana</i> , Mor. & Lyc.	*	smooth new sp. ?	
<i>nuciformis</i> , Lycett.		Pholadomya	
<i>trapeziformis</i> , Römer, sp.	* *	<i>Dawalquea</i> , Lycett.	
Cytherea (Cyprina)		<i>fidicula</i> , Sow.	*
<i>dolabra</i> , Phil.		<i>Heraulti</i> , Ag.	* *
Goniomya		<i>lirata</i> , Sow.	*
<i>literata</i> , Sow. sp.		<i>ovalis</i> , Sow.	*
<i>V-scripta</i> , Sow. sp.		<i>ovulum</i> , Ag.	
Gresslya , sp. ?		<i>Zietenii</i> , Ag.	*
Homomya (Myacites)		large new sp. ?	
<i>crassiuscula</i> , Mor. & Lyc.	? *	Pholas ? sp. ? (oolitica, Mor. & Lyc. ?)	*
<i>unioniformis</i> , Mor. & Lyc.	? *	Tancredia	
<i>Vezelayi</i> (?), d'Arch.	? *	<i>angulata</i> , Lycett	* *
Isocardia		<i>axiniformis</i> , Phil.	* *
<i>cordata</i> , Buckm.	*	<i>Tellina</i> , sp. ?	
Lithodomus		Trigonia	
<i>inclusus</i> , Phil.	* *	<i>compta</i> , Lycett	*

Lincolnshire-Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.	Lincolnshire-Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.
Trigonia			Ceritella		
<i>costata, Park.</i>	*	*	<i>acuta, Mor. & Lyc.</i>		
—, a minute var.	*		<i>parvula, Römer, sp.</i>		
<i>hemispharica, Lycett</i>	?		sp. (like <i>Sowerbyi, Mor. & Lyc.</i>)?		
<i>Phillipsii, Mor. & Lyc.</i>	*		Cerithium		
<i>sculpta, Lycett.</i>			<i>Beanii, Mor. & Lyc.</i>		
<i>Sharpiana (?) , Lycett</i>	*		sp. (like <i>costigerum, Piette</i>) ?		
<i>spinulosa (?) , Young & Bird.</i>			<i>gemmaum, Mor. & Lyc.</i>	*	
<i>striata, Miller.</i>			<i>limæforme (?) , Römer.</i>		
<i>V-costata, Lycett</i>	*		<i>quadriceinctum, Goldf.</i>		
Unicardium			<i>strangulatum, d'Arch.</i>		
<i>depressum, Phil. sp.</i>	*		Chemnitzia		
<i>impressum, Mor. & Lyc.</i>	*	*	<i>vetusta, Mor. & Lyc.</i>		
<i>parvulum, Mor. & Lyc.</i>	*	*	<i>Wetherellii, Mor. & Lyc.</i>		
BRACHIOPODA.			Cylindrites		
Rhynchonella			<i>acutus, Sow. sp.</i>		
<i>angulata, Sow. sp.</i>	*		<i>brevis, Mor. & Lyc.</i>		
<i>Crossii, Walker.</i>			<i>bullatus, Mor. & Lyc.</i>		
<i>plicatella, Sow.</i>			<i>cylindricus, Lycett.</i>		
<i>quadruplicata, Ziet.</i>	*		<i>turriculatus, Lycett.</i>		
<i>spinosa, Schloth. sp.</i>			Delphinula		
<i>sub-decorata, Dav.</i>	*		<i>alta (?) , Mor. & Lyc.</i>		
<i>sub-tetraëdra, Dav.</i>	*		sp. ? (see 'Phil. G. Y.' i.		
<i>varians (?) , Schloth. sp.</i>			tab. ix. f. 32).		
Terebratula			(<i>Crossostoma</i>) <i>Prattii, Mor. & Lyc.</i>		
<i>Buckmani, Dav.</i>	*		Dentalium (?) , new sp. ?		
<i>fimbria, Sow.</i>			Monodonta		
<i>globata, Sow.</i>	*		<i>Lyellii (?) , d'Arch.</i>		
<i>ovoides, Sow.</i>	*		sp. (like <i>Labadyei, d'Arch.</i>) ?		
<i>perovalis, Sow.</i>	*		Natica		
<i>simplex, Buckm.</i>			<i>adducta, Phil.</i>	*	
<i>sphaeroidalis, Sow.</i>			(<i>Uaspira</i>) <i>canaliculata, Mor. & Lyc.</i>		*
<i>sub-maxillata, Sow.</i>	*		<i>formosa, Mor. & Lyc.</i>		*
GASTEROPODA.			<i>grandis, Goldf.</i>		*
Actæonina			<i>Leckhamptonensis, Lycett.</i>		
<i>glabra, Phil. sp.</i>			—, large var. with punctated surface.		
<i>parvula, Römer, sp.</i>			<i>Michelini, d'Arch.</i>		*
large sp. ?			<i>punctura, Bean, sp.</i>		
Alaria			Nerinaa		
<i>armata, Mor. & Lyc.</i>	*	*	<i>cingenda, Bronn</i>	*	
<i>hamulus, Deslong. sp.</i>	?		<i>Cotteswoldia, Lycett.</i>		
<i>hamus, Deslong. sp.</i>	*		sp. (like small <i>Eudesii, Mor. & Lyc.</i>) ?		
<i>Phillipsii, d'Orb. sp.</i>			sp. (like <i>funiculus, Deslong.</i>) ?		
<i>sub-punctata, Goldf. sp.</i>			<i>gracilis, Lycett.</i>		
<i>trifida, Phil. sp.</i>	*	*			

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

295

Lincolnshire-Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.	Lincolnshire-Limestone Fossils.	Northampton Sand.	Great Oolite, N. & S.
Cidaris Fowleri, <i>Wright</i>	*		Thecosmilia gregaria, <i>M Coy</i>	*	
Wrightii, <i>Desor</i>	*		—, a var. Genus?, sp.?		
Clypeus Michelini, <i>Wright</i> .			CRUSTACEA.		
Echinobrissus clunicularis, <i>Lhwyd</i>	*	*	Pseudophyllia, sp.?		
Galeropygus (Hyboelypus) agariciformis, <i>Forbes</i>	*		Claws—genus? sp.?	*	
Holcotypus hemisphaericus, <i>Ag.</i>			PISCES.		
Pseudodiadema depressum, <i>Ag.</i>			Acrodus or Hybodus? tooth.		
Pygaster semisulcatus, <i>Phil.</i>	*		Hybodus dorsalis (?), <i>Ag.</i> , large spine		*
Stomechinus germinans, <i>Phil.</i>	*		— (?), small spine		*
Astropecten Cotteswoldiae, var. <i>Stamford-</i> <i>ensis, Wright.</i>			Pycnodus Bucklandi, <i>Ag.</i> —teeth		*
Pentacrinus sub-sulcatus, <i>Goldf.</i> (joints).			Strophodus magnus, <i>Ag.</i> —palates		*
			sub-reticulatus, <i>Ag.</i> —palates		*
ZOOPLHYTA.			REPTILIA.		
Anabacia orbulites, <i>Lamx.</i> sp.	*		Teleosaurus sp.?		
Calamophyllia radiata, <i>Lamx.</i> sp.	*		sp.?		
sp. (like <i>Stokesii</i> , <i>Edw. &</i> <i>Haime</i>)?			} teeth.		*
Cladophyllia Babeana, <i>Edw. & Haime</i>	*		POLYZOA.		
Isastræa limitata, <i>Lamx.</i> sp.	*		Several sp?	*	
Latimæandra Davidsoni, <i>Edw. & Haime</i> ...	*		PLANTÆ.		
Flemingii, <i>Edw. & Haime.</i>			Aroides Stutterdi, <i>Carruth.</i>		
Montlivaltia Delabechii, <i>Edw. & Haime.</i>			Ferns, sp.		
tenuilamellosa, <i>Edw. & Haime.</i>			Pecopteris polypodioides, <i>Lindley.</i>		
Thamnastrea concinna, <i>Goldf.</i>	*		Palæozamia pectinata, <i>Brong.</i>		
—, a very fine variety.			Zamia sp. ? (small perfect frond).		
			Wood	*	*

Fossils from the Ferruginous Beds of the NORTHAMPTON SAND†.—Table showing which Forms are contained also in the LINCOLNSHIRE LIMESTONE and GREAT OOLITE Beds of the Northampton and Stamford Districts.

Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.	Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.
LAMELLIBRANCHIATA (<i>Monomyaria</i>).					
<i>Avicula</i>			<i>Lima</i>		
<i>Braamburiensis</i> , Sow.	*		<i>rudis</i> , Sow. sp.		
sp. (allied to <i>complicata</i> , Buckm.)?			<i>Sharpiana</i> , Ether. M.S.		
<i>univalvis</i> , Sow.			large sp. (allied to <i>L. grandis</i> , Römer)?	*	
<i>eri</i> , Goldf.	*	*	large sp.?	*	
			several new or unidentified forms.		
<i>acuta</i> , Sow.	*		<i>Ostrea</i>		
<i>Hartmanni</i> , Goldf. (allied to <i>radiana</i> , Mor. & Lyc.)	*		<i>costata</i> , Sow.	*	
sp. (like <i>Islipensis</i> , Lycett)?			<i>cristagalli</i> , Quenst.		
<i>lata</i> , Phil.	*		<i>flabelloides</i> , Lam.	*	*
sp. (like <i>ovata</i> , Sow. sp.)?			<i>gregaria</i> , Sow.	*	
<i>pernoides</i> , Deslong.			<i>sub-rugulosa</i> (?), Mor. & Lyc.	*	*
<i>prælonga</i> (?), Lycett.			<i>sulcifera</i> , Phil.	*	
(<i>Gastrochæna</i>) <i>tortuosa</i> , Phil. sp.			large flat sp.?	*	*
<i>Gryphæa</i>			<i>Pecten</i>		
<i>mima</i> , Phil.	*		<i>arcuatus</i> , Sow.	*	
<i>subloba</i> , Desh.			<i>articulatus</i> , Schloth. (a var.)...	*	*
<i>Hinnites</i>			<i>clathratus</i> , Römer	*	
<i>abjectus</i> , Phil. sp.	*		<i>demissus</i> , Phil.	*	*
<i>velatus</i> , Goldf. sp.	*		<i>lens</i> , Sow.	*	*
<i>Inoceramus</i>			<i>personatus</i> , Goldf.	*	
<i>Fittoni</i> , Mor. & Lyc.			<i>Perna</i>		
<i>obliquus</i> , Mor. & Lyc.	*		(<i>Inoceramus</i>) <i>quadrata</i> , Phil. sp.		
<i>Lima</i>			<i>rugosa</i> , Goldf.	*	*
sp. (closely allied to <i>antiquata</i> , Sow.)?			<i>Pinna</i>		
<i>bellula</i> , Mor. & Lyc.	*		<i>ampla</i> , Sow. (a var.)		
<i>cardiiformis</i> , Sow. sp.	*	*	<i>cuneata</i> , Phil.	*	*
<i>deltoides</i> , Ether. M.S.			<i>Placunopsis</i>		
<i>duplicata</i> , Sow. sp.	*	*	<i>Jurensis</i> , Römer.		
<i>Dustonensis</i> , Ether. M.S.			<i>Plicatula</i>		
<i>electra</i> , d'Orb.			<i>tuberculosa</i> , Mor. & Lyc.	*	
<i>etheridgii</i> , Wright	*	*	<i>Pteroperna</i>		
<i>impressa</i> , Mor. & Lyc.	*	*	<i>costatula</i> , Deslong.	*	*
<i>interincta</i> , Phil.			<i>gibbosa</i> , Lycett.		
<i>Luciensis</i> (?), d'Orb.	*		<i>plana</i> , Mor. & Lyc.	*	*
<i>pectinoides</i> , Sow. sp.			LAMELLIBRANCHIATA (<i>Dimyaria</i>).		
<i>Pontonis</i> , Lycett	*		<i>Arca</i>		
<i>proboscidea</i> , Sow. sp.	*		<i>æmula</i> , Phil.	*	
<i>punctata</i> , Sow. sp.	*		<i>minuta</i> , Sow.		
<i>Rodburgensis</i> , Lycett M.S.	*		<i>Prattii</i> , Mor. & Lyc.	*	

† *Errata* in the Table published with the First Part of this Memoir have been corrected in this Table, and the names of some Fossils since obtained have been added.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

297

Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.	Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.
Astarte			Lucina		
depressa, <i>Goldf.</i>	*	*	Bellona, <i>d' Orb.</i>	*	
elegans, <i>Sow.</i>	*		despecta, <i>Phil.</i>	*	
excavata, <i>Sow.</i> (a var.)	*		— var. cardioides, <i>d' Arch.</i>	*	
minima, <i>Phil.</i>	*		rotundata, <i>Römer, sp.</i>	*	
rhomboidalis, <i>Phil.</i> , sp. (<i>Hip-</i>			Macrodon		
<i>popodium Luciense, d' Orb.</i>)	*		Hirsonensis, <i>d' Arch. sp.</i>	*	*
Cardium			Modiola		
Buckmani, <i>Mor. & Lyc.</i>	*	*	Binfieldi (?), <i>Mor. & Lyc.</i>	*	
cognatum, <i>Phil.</i>	*	*	cuneata, <i>Mor. & Lyc.</i>	*	*
semicostatum, <i>Lycett.</i>			explanata, <i>Morris.</i>		
sp. (near to Stricklandi, <i>Mor.</i>			gibbosa, <i>Sow.</i>	*	*
& <i>Lyc.</i>)?	*		Leckenbyi, <i>Mor. & Lyc.</i>		
Ceromya			Lonsdalei (?), <i>Mor. & Lyc.</i>	*	
Bajociana, <i>d' Orb.</i>	*		solenoides, <i>Mor. & Lyc.</i> (a		
concentrica, <i>Sow.</i>	*	*	new var.)		
new sp.?			Sowerbyana (?), <i>d' Orb.</i>	*	*
Corbis (Corbicella)			subreniformis, <i>Mor. & Lyc.</i>	*	*
sp. (allied to Bathonica, <i>Mor.</i>			Myacites		
& <i>Lyc.</i>)?			dilatatus, <i>Phil. sp.</i>		
sp. (allied to Lajoyei,			sp.?		
<i>d' Arch.</i>)?			Myoconcha		
Cucullæa			crassa, <i>Sow.</i>	*	
cancellata, <i>Phil.</i>	*		Mytilus		
cucullata, <i>Goldf.</i>	*	*	furcatus, <i>Goldf.</i>	*	
elongata, <i>Sow.</i>	*		Opis		
imperialis, <i>Phil.</i>	*		lunulatus, <i>Sow. sp.</i>	*	
oblonga, <i>Sow.</i>	*		Pholadomya		
ornata, <i>Phil.</i>			ambigua, <i>Sow.</i>		
large new sp., beautifully can-			fidicula, <i>Sow.</i>	*	
cellated.			fidicula? or new sp. (semi-		
several sps. undetermined.			acuticosta?).		
Cypricardia			Heraulti, <i>Ag.</i>	*	*
acutangula, <i>d' Orb.</i>			ovalis (?), <i>Sow.</i> (a very large		
Bathonica, <i>d' Orb.</i>	*	*	example)		
nuculiformis, <i>Römer, sp.</i>	*	*	Zietenii, <i>Ag.</i>	*	
Cyprina			large new sp.?		
trapeziformis, <i>Römer</i>	*	*	Quenstedtia		
Gressiia			lavigata, <i>Mor. & Lyc.</i>	*	
abducta, <i>Phil.</i>			oblita, <i>Phil. sp.</i>	*	
latirostris, <i>Ag.</i>			Tancredia		
rostrata, <i>Ag.</i>			angulata, <i>Lycett</i>	*	*
Homomya (Myacites)			axiniformis, <i>Phil. sp.</i>	*	*
crassiuscula (?), <i>Mor. & Lyc.</i>	*	*	planata, <i>Mor. & Lyc.</i>	*	*
Vezelayi (?), <i>d' Arch. sp.</i>	?	*	Trigonia		
sp. (new, allied to gibbosa,			angulata, <i>Sow.</i>		
<i>Sow. sp.</i>)?			compta, <i>Lycett.</i>	*	
Isocardia			— (a large var.)		
cordata, <i>Buckm.</i>	*		costata, <i>Park.</i>	*	*
large "new sp." (<i>Dr. Lycett.</i>)			— (a minute var.)?	*	
Lithodomus			duplicata, <i>Sow.</i>		
inclusus, <i>Phil.</i> (very large)....	*	*	hemispherica (?), <i>Lycett</i>	*	

Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.	Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.
Trigonia			Natica		
Phillipsii, <i>Mor. & Lyc.</i>	*		adducta, <i>Phil.</i>	*	
Sharpiana, <i>Lycett</i> , M.S.	?		neritoidea (?), <i>Mor. & Lyc.</i>		*
tuberculosa (?), <i>Lycett</i>	*		Verneuilii (?), <i>d'Arch.</i>		*
V-costata, <i>Lycett</i>	*		large spired sp. ?		
new sp. ?			several undetermined sp.		
Unicardium			Nerinaea		
depressum, <i>Phil.</i> sp.	*		cingenda, <i>Bronn</i>	*	
gibbosum, <i>Mor. & Lyc.</i>	*	*	triplicata, <i>Bronn</i>	*	
impressum, <i>Mor. & Lyc.</i>	*	*	several small sps. ?		
parvulum, <i>Mor. & Lyc.</i>	*	*	Nerita		
BRACHIOPODA.			costulata, <i>Desh.</i>		
Rhynchonella			hemisphaerica (?), <i>Römer</i>		*
angulata, <i>Sow.</i> sp.	*		Onustus ?		
cyncephala, <i>Rich.</i>			sp. (like <i>Burtonensis</i> , <i>Lyc.</i>) ?		?
quadriplicata, <i>Ziet.</i>	*		Patella		
sub-decorata, <i>Dav.</i>	*		inornata, <i>Mor. & Lyc.</i>		
tetraëdra, <i>Sow.</i> †			nana, <i>Sow.</i>		
sub-tetraëdra, <i>Dav.</i>	*		rugosa, <i>Sow.</i>	*	
variabilis, <i>Schloth.</i>			Pleurotomaria		
—, var. <i>bidens</i> , <i>Phil.</i> sp.			Aglaiia, <i>d'Orb.</i> (a var.)		
—, var. <i>triplicata</i> , <i>Phil.</i> sp.			armata, <i>Münst.</i>	*	*
Terebratula			clathrata, <i>Goldf.</i>		
Buckmani, <i>Dav.</i>	*		sp. (allied to <i>Marcousana</i> ,		
globata, <i>Sow.</i>	*		<i>d'Orb.</i>) ?		*
impressa, <i>Von Buch.</i>			ornata, <i>DeFrance</i>	*	
ovoides, <i>Sow.</i>	*		pyramidata, <i>Phil.</i> sp.		
perovalis, <i>Sow.</i>	*		Rimula		
sub-maxillata, <i>Sow.</i>	*		Blottii, <i>Deslong.</i>	*	
GASTEROPODA.			Trochotoma		
Actæon			calix, <i>Phil.</i>		
Sedgwicki, <i>Phil.</i> sp.			obtusa, <i>Mor. & Lyc.</i>	*	
Actæonina			tabulata, <i>Mor. & Lyc.</i>	*	*
large new sp. ?			Trochus		
Alaria			ornatissimus, <i>d'Orb.</i> , var.		
armata, <i>Mor. & Lyc.</i>	*	*	Pontonis, <i>Morris</i> (?)	*	
large sp. (like <i>hamulus</i> , <i>Mor.</i>			CEPHALOPODA.		
& <i>Lyc.</i>) ?			Ammonites		
hamus, <i>Deslong.</i>	*		bifrons, <i>Phil.</i>		
trifida, <i>Phil.</i>	*	*	insignis (junior), <i>Schubler.</i>		
Cerithium			Murchisonae, <i>Sow.</i>	*	
gemmatum, <i>Mor. & Lyc.</i>	*		—, var. <i>corrugatus</i> , <i>Sow.</i>		
Chemnitzia			Niortensis, <i>d'Orb.</i>		
Scarburgensis, <i>Mor. & Lyc.</i>			opalinus, <i>Rein.</i>		
Cirrus			sp. (like <i>solaris</i> , <i>Phil.</i>) ?		
acutus, <i>Sow.</i>			sp. (unnamed, also found in		
nodosus, <i>Sow.</i>			Inf. Oolite at Yeovil).		
Littorina			large smooth sp. ?	*	
punctura, <i>Bean.</i>			Belemnites		
			sp. (like <i>acutus</i> , <i>Miller</i>) ?	*	

† Vide Davidson's Ool. Brach., pl. xviii. fig. 10. It is commonly found in the "Ironstone" of Northamptonshire.

1873.]

SHARP—OOLITES OF NORTHAMPTONSHIRE.

299

Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.	Northampton-Sand Fossils.	L. Limestone.	Gt. Oolite, N. & S.
Belemnites			ZOOPHYTA.		
Bessinus (?), <i>d' Orb.</i>	*	*	Isastræa		
Blainvillii, <i>Voltz</i>	*		explanulata, <i>M'Coy.</i>		
canaliculatus, <i>Schloth.</i>	*		Richardsoni, <i>Edw. & Haime.</i>	*	
elongatus, <i>Miller.</i>			Latimæandra		
phragmacones of large sp.			Davidsoni, <i>Edw. & Haime.</i>		
Nautilus			Montlivaltia		
clausus, <i>d' Orb.</i>			Wrightii, <i>Edw. & Haime.</i>		
obesus, <i>Sow.</i>	*		Stylina		
polygonalis, <i>Sow.</i>	*		solida (?), <i>M'Coy.</i>		
sinuatus, <i>Sow.</i>			Thamnastræa		
ARTICULATA.			concinna, <i>Goldf.</i>	*	
Serpula			Thecosmilia		
convoluta, <i>Goldf.</i>	*		gregaria, <i>M'Coy</i>	*	
plicatilis (?), <i>Goldf.</i>	*		Wrightii, <i>Duncan.</i>		
socialis, <i>Goldf.</i>	*	*	Several undetermined genera		
Vermicularia			and sp.?		
compressa, <i>Phil.</i>			AMORPHOZOA.		
Annulated Annelide tubes (found			Spongiæ, sp.?		
also near Banbury).....	*	*	POLYZOA.		
ECHINODERMATA.			several sp.?	*	
Acrosalenia			CRUSTACEA.		
Lycettii, <i>Wright</i>	*		claws, &c., sp.?	*	
spinosa (?), <i>Ag.</i>	*	*	REPTILIA.		
Cidaris			Megalosaurus—tooth.		
Fowleri, <i>Wright</i>	*		Teleosaurus, un-		
Wrightii, <i>Desor</i>	*		described sp.? } dorsal scute.		
Clypeus			PLANTE.		
Hugii, <i>Ag.</i>			Impressions—		
Echinobrissus			bracts of cones?		
clunicularis, <i>Lhwyd</i>	*	*	flag-like leaves, with seeds.		
Galeropygus (Hyboclypus)			leaf-shape pinnule of fern-		
agariciformis, <i>Forbes</i>	*		frond, showing venation		
Hyboclypus			and sori.		
ovalis, <i>Wright.</i>			Carpolithes (fruit)	*	
Pygaster			Seeds.		
semisulcatus, <i>Phil.</i>	*		Wood.....	*	*
Stomechinus					
germinans, <i>Phil.</i>	*				
Stellaster					
Sharpii, <i>Wright.</i>					
Pentacrinus					
stem and branch joints.					

Approximate Analysis of the Synchronous Distribution of Fossils.

The foregoing Lists give the following results :—

Number of Great-Oolite forms	230
„ „ Lincolnshire-Limestone forms	315
„ „ Northampton-Sand forms	250
Total	795

Of these—

- 135 are common to the Northampton Sand and the Lincolnshire Limestone.
- 70 to the Lincolnshire Limestone and Great Oolite.
- 46 to the Great Oolite and Northampton Sand.
- 43 to all three formations.

If, from the number of forms common to the Northampton Sand and the Lincolnshire Limestone, and from the number common to the Lincolnshire Limestone and the Great Oolite, respectively, be deducted the number of forms common to all three formations (as having no significance in determining their comparative affinity), there will remain :—

- 92 forms common only to the Northampton Sand and the Lincolnshire Limestone.
- 27 forms common only to the Lincolnshire Limestone and the Great Oolite.

Many fossils of the Lincolnshire Limestone and Northampton Sand have yet to be identified and described, which ultimately will considerably swell the foregoing numbers as regards those formations.

EXPLANATION OF PLATES IX. AND X.

PLATE IX.

Map of the author's route and of the southern portion of the Lincolnshire Limestone. * * Points of thinning-out of the Lincolnshire Limestone. For section see Pl. X. fig. 2.

PLATE X.

Fig. 1. Diagrammatic section across the Welland Valley at Stamford.
Fig. 2. Generalized section from the valley of the Welland to that of the Nene, illustrating the thinning-out of the Lincolnshire Limestone.

DISCUSSION.

Mr. ETHERIDGE expressed his obligation to the author for his paper, and for the remarkable collection of fossils he had exhibited. The ground over which he had worked was one the features of which had required a great amount of well-directed labour to decipher. He considered, however, that Mr. Sharp and Mr. Judd had settled the question of the sequence of these rocks, and their relation to the Oolitic beds of Yorkshire to the north, and Gloucestershire to the west. The importance of the determination of the position of a bed of such commercial value as the Northampton Sand could hardly be overestimated; and it was to Mr. Judd and to the author that this

MAP OF ROUTE AND OF SOUTHERN PORTION OF AREA OF THE LINCOLNSHIRE LIMESTONE.

Fig:
DIAGRAM OF SECTION ACROSS THE WELLAND VALLEY
AT STAMFORD

Fig:2.
GENERALIZED SECTION FROM THE VALLEY OF THE NENE TO THAT OF THE NENE, ILLUSTRATING THE THINNING OUT OF THE
LINCOLNSHIRE LIMESTONE

determination was due. It had, moreover, been attained under very adverse circumstances; for at the time when the survey of the district had been undertaken, the Lincolnshire Limestone had not been recognized; and even Prof. Morris had at first failed to see that this bed intervened between the Northampton Sand and the Great Oolite. It was mainly due to the extensive collection formed by Mr. Sharp that the key to the existence of this important bed and to the geological history of the whole district during the Mesozoic period was discovered. The mapping of the country was not to be effected by studying merely its lithological characters, but was mainly dependent on a knowledge of the palæontological features of each of the successive beds. Mr. Etheridge pointed out the close correspondence between the position of the Northampton beds and those of Yorkshire, with the exception of the absence of the Great Oolite in the latter area. Though the Stonesfield slates and the Collyweston beds were so similar in lithological character that even the most experienced might take the one for the other; yet, when the organic contents came to be examined, the difference became evident; and in Mr. Sharp's paper stratigraphical evidence had been brought to corroborate the palæontological, and to show conclusively the difference in the horizon of the two beds.

Mr. JUDG could not overestimate the value of Mr. Sharp's labours, extended, as they had been, over nearly a whole lifetime. They afforded another instance of the great value of local inquiries in geology. It was becoming more and more evident that the sequence of beds which held good for one place required some modification in another, and that in each case there was a more or less distinct local series—showing that in no one locality was the sequence absolutely perfect, as indeed had been already pointed out by Mr. Darwin.

Mr. CHARLESWORTH remarked on a specimen of the teeth of the genus *Lepidotus* exhibited, consisting of cylindrical columns surmounted by a conical crown, which struck him as one of unusual interest in the magnificent collection displayed. He commented on the value of such local collections for palæontological purposes, and on the necessity of their being formed if the progress of geology was to be furthered.

Prof. DUNCAN also was highly impressed with the value of Mr. Sharp's collection. He remarked on two specimens of *Madreporearia*, one of which, *Thamnastræa concinna*, presented a series of ridges significant of intermittent growth. The other form presented a strange relic of palæozoic coral forms, being intersected by tabulæ like those of ancient times. It afforded an instance of a tabulate Actinozoon, in opposition to the opinion of Agassiz that the Tabulata belonged to the Hydrozoa. Both species grew on narrow bases; and though not reef-corals, were forms such as were to be found in the neighbourhood of reefs. Taken in conjunction with the Saurian remains, he thought they were symptomatic of shallow sea conditions, such as those existing between the continent of America and the West-Indian Islands.

Mr. H. WOODWARD contrasted the collections formerly exhibited

to the Society with that now shown, inasmuch as in former times the same amount of regard had not been paid to the stratigraphical position of the specimens, and their value was in consequence almost destroyed.

Mr. SHARP, in reply, mentioned that some portion of his collection had been formed by Mr. Bentley, and not by himself, and that therefore he could not claim the credit of the whole. He was gratified to find that there was so little disagreement with his conclusions.

The CHAIRMAN (Mr. Warrington W. Smyth), in conclusion, expressed his satisfaction at the fact that, notwithstanding the existence of the Geological Survey, there was still room left for individual discovery and research, which in this instance had served to rectify what might have been erroneous views on the part of the Survey.