PS. Zur Orientirung über den weiteren Lauf des Cometen — derselbe wird gegen Ende Juli für uns unsichtbar — habe ich noch die nebenstehenden Oerter gerechnet.

Herr Dr. Schorr machte mich auf die Aehnlichkeit der Bahn mit derjenigen des Pons-Brooks'schen Cometen 1884 I aufmerksam. Es wäre von Interesse, wenn auch die Bahn des Cometen Perrine sich als Ellipse herausstellen sollte.

12^h M. Z. Berlin.

1898	α	δ	$\log r$	$\log A$	H
Aug. 14	8 ^h 9 ^m 0	+11° 56'	9.8039	0.1626	8.4
30	9 17.2	— 9 14	9.8475	0.1577	7.0
Sept. 15	10 36.0	— 24 43	9.9513	0.1920	3.7
Oct. 1	11 57.2	— 37 44	0.0524	0.2523	1.8
17	13 11.7	— 43 29	0.1374	0.3173	0.9

A. B.

Ephemeride des Cometen 1898... (Coddington-Pauly).

Fortsetzung zu A. N. 3500. 12^h M. Z. Berlin.

1898	α app.	δ app.	$\log r$	$\log A$	1898	α app.	δ app.	$\log r$	$\log A$
Juli 25	14 ^h 9 ^m 31 ^s	—45° 38' 5	0.2600	0.1191	Aug. 20	13 ^h 38 ^m 4 ^s	—52° 39' 2		
26	7 35	45 56.0			21	37 33	52 55.5		
27	5 42	46 13.2			22	37 4	53 11.8	0.2352	0.2054
28	3 53	46 30.2			23	36 38	53 28.3		
29	2 8	46 47.0	0.2556	0.1327	24	36 14	53 44.8		
30	14 0 26	47 3.6			25	35 53	54 1.5		
31	13 58 49	47 20.1			26	35 34	54 18.3	0.2330	0.2154
Aug. 1	57 15	47 36.4			27	35 18	54 35.2		
2	55 45	47 52.6	0.2514	0.1461	28	35 4	54 52.2		
3	54 19	48 8.7			29	34 52	55 9.3		
4	52 56	48 24.8			30	34 42	55 26.6	0.2312	0.2246
5	51 36	48 40.8			31	34 35	55 44.0		
6	50 20	48 56.7	0.2475	0.1591	Sept. 1	34 30	56 1.6		
7	49 7	49 12.6			2	34 27	56 19.3		
8	47 58	49 28.4			3	34 26	56 37.2	0.2298	0.2332
9	46 52	49 44.2			4	34 27	56 55.2		
10	45 48	49 59.9	0.2439	0.1716	5	34 31	57 13.3		
11	44 48	50 15.7			6	34 37	57 31.6		
12	43 51	50 31.5			7	34 44	57 50.1	0.2289	0.2411
13	42 57	50 47.3			8	34 54	58 8.7		
14	42 7	51 3.1	0.2406	0.1835	9	35 7	58 27.5		
15	41 20	51 19.0			10	35 22	58 46.5		
16	40 35	51 34.9			11	35 38	59 5.6	0.2283	0.2483
17	39 53	51 50.9			12	35 56	59 24.9		
18	39 14	52 6.9	0.2377	0.1948	13	36 17	59 44.4		
19	38 38	52 23.0			14	36 40	60 4.1		
20	13 38 4	—52 39.2			15	13 37 5	—60 24.0	0.2282	0.2550

Eine Strassburger Beobachtung vom 21. Juni giebt die Correction $-0.3 + 0.15$.

Berlin 1898 Juni 26.

A. Berberich.

On the Constant of Aberration.

From observations made here according to Talcott's method the value of the constant of aberration is

$$20.477 \pm 0.040.$$

The constant obtained from stars of average magnitude 4.4 is

$$20.639 \pm 0.075.$$

From those of average magnitude 5.4 it is

$$20.430 \pm 0.063,$$

Hongkong Observatory, 10th May 1898.

and from those of average magnitude 6.4 it is

$$20.385 \pm 0.066.$$

The fact that the constant of aberration thus determined is found to depend upon the magnitude of the stars observed may possibly serve to explain differences in the values obtained at different observatories, which differences far exceed their probable errors.

W. Doberck.