Correspondence.

"Audi alteram partem."

THE JUBILEE AND THE HOSPITALS.

To the Editors of THE LANCET.

SIRS,—As the Queen's Jubilee requires commemoration in some manner worthy of her great reign, I beg to suggest that no better method could be found than to raise a fund for the permanent endowment of our great London hospitals. I think a fund such as you suggested last week, to free them from debt, would not be adequate to the occasion. The Imperial Institute and Church House will be supported by Loyalists and Churchmen all over the world, but unless some great scheme is started in London, to which all will feel bound to contribute, the money of the charitable will be frittered away in a thousand hobbies of doubtful utility. A sum of this magnitude would never be raised in the ordinary method of appeals to the benevolent few; but if the scheme were to receive the assent of the great body of the people, a voluntary assessment of a fiftieth part of the income of all those who now pay income-tax would receive a response sufficient for even so great a purpose.—I am, Sirs, your obedient servant,

Clifford-st., Bond-st., W., March 15th, 1887. G. SLIGHT, M.D.

* * We shall be only too happy if Dr. Slight's suggestion should find favour, and prove successful. Could not the heads of the medical colleges and the great hospitals organise a meeting to give form to the extensive wish to signalise the Jubilee by help to the hospitals?—Ed. L.

PERFLATION IN EMPYEMA.

To the Editors of THE LANCET.

SIRS,—The method of perflation described by me in your issue of July 31st, 1886, was capable of improvement, and has been improved in two essential particulars. 1. The idea suggested by Dr. Sansom at the last meeting of the Royal Medical and Chirurgical Society (which the lateness of the hour prevented my acknowledging), as to the advantages of an antiseptic air-filter instead of a column of antiseptic fluid, occurred to me several months ago, and was put into practice. Instead of containing a short column of carbolic acid solution, the Woolff's bottle was filled with tow saturated with liquefied carbolic acid or with strong solution of thymol, with a view to securing the following advantages: (a) absence of any fluid which, through inadvertence, might be allowed to run into the chest; (b) considerable subdivision of the air during its passage through the tow, and thorough mechanical cleansing of the same; (c) increased facility for warming the air in the bottle when necessary. At an early date, however, I discarded the Woolff's bottle, preferring to use a broad tube (one inch and a half in diameter and about ten inches long) fitted at either end with an indiarubber plug and short glass tube, and packed with medicated tow. This apparatus can lie flat on the table or on the bed, and is simply a broader segment inserted in the length of the tubing. It can readily be warmed by placing it in an appropriate hot-water trough, or by keeping it surrounded with hot flannel. 2. For the other improvement I am indebted to a suggestion from Mr. Marten, house-physician to Addenbrooke's Hospital, Cambridge. Instead of Richardson's bellows, I now employ a Fletcher's foot-bellows. This apparatus enables the operator to use both hands, whilst his foot works the bellows and controls the delivery of air. It thus becomes possible (provided a steam spray-producer is available) to dress an empyema without a trained assistant. The pressure which can be obtained from Fletcher's bellows is in excess of surgical requirements. But it is easy to obviate this difficulty with the help of a screw-clamp placed on the tubing between the chest and the air-filter. By this means the diameter of the tube may be varied so as to regulate to any extent the quantity of air delivered into the chest. I may add that perflation can be applied whether two openings exist into the chest or only one. A free incision (if possible in the posterior or lateral aspect of the chest) will usually suffice. I believe, however, that an

advantage is often gained by making in front a counter-opening, which need not be so large as the posterior wound.

I remain, Sirs, yours truly,

Curzon-street, Mayfair, March 11th, 1887.

WM. EWART.

P.S.—I avail myself of this opportunity of setting at rest the doubts expressed by some Fellows of the Royal Medical and Chirurgical Society as to the nature of the specimen exhibited as gangrenous lung tissue on the 8th inst. Dr. Délepine's microscopical examination confirms the conclusion drawn from naked-eye appearances that this was a portion of the lung, and the surmise that embolism was the cause of the gangrene.

CANCER OF THE BODY OF THE UTERUS.

To the Editors of THE LANCET.

SIRS,—I shall be glad to be allowed to correct an error in my letter on the above subject which appeared on March 5th. In referring to Case 22, described in Dr. John Williams's Harveian lectures, I stated that, according to my recollection, the disease had commenced in one of the fibroids, and had spread thence to the uterine wall. I have since had an opportunity of making a more complete examination of the specimen, and from the appearances presented by deeper sections, it is clear that the uterine wall, and probably the mucous membrane, was the site of the primary disease.

With regard to the question on which our opinions differed—viz., the frequency with which cancer affects the body of the uterus,—I find that at the Middlesex Hospital we have had only two such cases in the post-mortem room during the last eleven years. This may possibly be due, as Dr. Williams suggests to me, partly to the fact that such, not being recognised outside as cases of malignant disease, are not sent into our cancer wards, and that, when diagnosed, they are considered too "good" to be given up to a hospital. On the other hand, Dr. Williams's experience of such cases may be exceptionally large, as they would certainly present difficulties, and on that account would be likely to be submitted for the opinion of a consultant.

I am, Sirs, yours truly,

J. KINGSTON FOWLER, M.D.

Clarges-street, W., March 15th, 1887.

IODOFORM AS AN ANTISEPTIC.

To the Editors of THE LANCET.

SIRS,—The doubts cast upon the antiseptic power of iodoform are surprising, and must surely be contradicted by the experience of every practical surgeon. It really seems absurd to quote instances to prove the superiority of iodoform (through the penetrating character of its volatile particles) to other antiseptics, it is a matter of such daily experience. I will just mention one case, however, if you will permit me. When surgeon on board the Orient steamship *Lusitania* (temporarily an armed cruiser), I had an invalided engineer from the Nile boat corps handed over to my charge at Alexandria, with the worst "poisoned hand" I ever saw. It was riddled with abscesses back and front; it had been deeply incised in several places, the pus burrowing up the tendon sheaths above the wrist; and it was horridly offensive. It had been treated with carbolic baths continuously. I kept it four hours in a carbolic bath, and then syringed it well and dressed it with carbolic solution (1 in 40). At night it was as offensive as ever. I then wrapped the fingers and hand with strips of lint on which was spread a weak iodoform ointment (twenty grains to an ounce of zinc ointment). The hand in the morning was perfectly sweet. The dressing was simply renewed, first twice a day, then daily. The healing process commenced immediately, discharge rapidly diminished, and there was never the slightest odour from the hand after the first dressing. The man's temperature became normal (it had been 101°), appetite good, and he went ashore with his hand healed. I would only say, in conclusion, that, as my supply of this invaluable drug was very small, I learned how exceedingly small an amount was necessary, and that the waste to which I had been accustomed in London hospitals was simply sinful. To powder a wound over or stuff a sinus with pure iodoform is utterly unnecessary. I am convinced that drachms are used when grains would suffice, especially in hospital practice.—I am, Sirs, yours obediently,

March, 1887.

J. P. GLOVER, M.D. Lond.