draught was now repeated, and a common enema was injected, which, however, returned without feculent matter, but there was some blood with it; the man was further directed to take an ounce of castor oil. When Mr. Jeffreys saw the patient, about noon, the spasm of the abdominal muscles had abated, but the tenderness of the abdomen had increased, and the countenance was becoming anxious; he brought up the castor oil soon after taking it. Mr. Jeffreys now directed him to take camphor and calomel, of each five grains, with two grains of opium, in two pills, immediately; and also to take three table spoonful of the following mixture every four hours: one ounce of sulphate of magnesia, one drachm and a half of compound spirit of sulphuric æther, and eight ounces of camphor mixture.—During the afternoon another clyster was administered, which brought away no feces: towards midnight the pulse became quick and fluttering, the countenance sunk and anxious, and the vomiting still continued.

5. Has had a restless night; the pain is still very severe. Twelve leeches were applied to the abdomen last night, and he has been put into a warm-bath this morning, without any relief. Some blood has passed per anum, but nothing like feces have passed. The vomiting still continues, and a brown fluid, of a thick consistence, is ejected. The pulse is quick, 112, weak, and scarcely perceptible; the abdomen is tense, distended, and exquisitely tender on pressure; the patient has not evacuated any urine since yesterday morning; he feels a desire, but has not the power. Mr. Jeffreys attempted to pass the catheter, for a considerable time, but failed.

2 o'clock, P. M. The countenance is sunk, and of a leaden hue; the finger nails are livid, and the feet are cold; the whole body is bedewed with the cold damp sweat, precursory to death. About five o'clock hiccup came on; he then sunk rapidly, and died at seven, P. M.

Post Mortem Examination.

The abdomen having been laid open, the peritoneum, covering the intestine, was found to have been in a high state of inflammation; a large quantity of purulent fluid was discovered, partly in the abdominal and partly in the pelvic cavity. The bladder and rectum, with the sphincter ani, were now removed from the body. On making an incision into the rectum, at its back part, a very extensive ulceration of its mucous membranes was brought into view. The ulceration commencing at the anus, occupied the whole circumference of the gut, for about two inches; above this, was a circle of sound membrane, to about the same extent, the lower edge of which being

thickened and contracted, had formed the stricture. Above the sound part, the ulceration had extended upwards, for at least four inches, and in this portion there was an ulcerated opening in the fore part of the bowel, extending through the peritoneal covering into the cavity of the pelvis. At the sphincter ani there was another ulcerated opening, leading to a cavity, of the size of an egg, between the rectum and bladder. Neither feces nor blood were detected in the abdominal cavity.

OPERATIONS.

On Thursday, the 22d ult. Mr. Keate performed the operation of lithotomy on a boy five years of age. The external incision was made by the scalpel, and the bladder cut into with the *bistouri cachée*. The calculus, which was of a moderate size, was readily extracted, and the boy has done well. On the same day, Mr. Brodie removed some portions of necrosed bone from the tibia of a little girl, and Mr. Rose removed the metatarsal bone of the great toe.

ERYSIPELAS.

To the Editor of THE LANCET.

SIR,—The subject of erysipelas having of late much occupied the pages of your valuable publication, and as I acknowledge myself an advocate for sparing the patient, (when it can be done consistently with safety) the pain necessarily attendant upon incisions made in exquisitely tender and inflamed parts: I have taken the liberty of sending you the following case, to which you will oblige me by giving publicity.

Mary Wensley, a woman of a full healthy habit, was seized, without any assignable cause, with active erysipelatous inflammation, (erysipelas vesiculosum) occupying the whole of the leg, attended with the usual symptoms of irritative fever; in visiting her, the limb was directed to be enveloped in a large linseed meal poultice, powerful aperient medicines were administered, and the antiphlogistic system in every respect rigidly insisted on; notwithstanding this, the erythema continued to extend itself until the entire thigh and leg were involved in one continuous inflammatory action; the same treatment was persisted in, with the occasional use of a spirituous lotion; and although at one period the constitutional symptoms became exceedingly alarming, the diseased action gradually subsided, and the patient is now in a fair way of recovery.

Bristol, Dec. 12th, 1827.

I. EARLE.

Mr. Earle partakes of the common error; this was not a case for incisions.—ED. L.