to entertain the idea that his physical punishment would balance his crimes; these were, without doubt, wholly imaginative. For the first night or two the insomnia was marked, large and repeated doses of chloral giving no sleep; nor were morphia or tincture of opium more successful; but Dr. Clouston's prescription, thirty minims of the tincture of Indian hemp with thirty grains of bromide of potassium, which he has found so valuable in asylum practice, was quite efficacious in a single dose. Nay, more; the tendency to sleeplessness persisting, it was used every night for nearly a month with undiminished effect, by the end of which time the man was at any rate in nearly his proper senses. His diet was, of course, liberal, though it included but little stimulant. The stump was nearly healed in a fortnight, hardly any pus having been produced, and, despite a good deal of bruising during the time of his violence, without any action that could be termed inflammatory.

I append a thermometrical table. It will be seen that the highest temperature was not attained before the fifth day—two days later than is usually the case, but being, however, about the time of his most marked insanity. It will also be noticed that the pulse and respiration reached their maxima before the temperature attained its highest point. All these factors were normal by about the eleventh day, although for some time longer the patient was under the influence of his delusions. On the fifth day, when his general aspect was alarming, and his tongue dry and brownish, the sphygmograph gave a favourable prognosis, the pulse sustaining some pressure without obliteration, and diastole being nearly absent in the tracing. He left at the expiration of about two months, with the stump well consolidated, and apparently sane, though eccentric.

	Temp.	Pulse.	Resp.
March 16th, evening	100.6°	84	18
„ 17th, morning	101.2°	84	18
„ „ evening	100.0°	96	24
„ 18th, morning	100.5°	96	24
„ „ evening	102.4°	108	18
„ 19th, morning	100.0°	96	36
„ „ evening	102.0°	120	48
„ 20th, morning	103.2°	96	24
„ „ evening	103.0°	96	24
„ 21st, morning	102.0°	96	24
„ 22nd, morning	99.7°	84	24
„ 23rd, morning	100.4°	84	18
„ „ evening	102.0°	96	24
„ 24th, morning	99.6°	96	—
„ 25th, morning	99.2°	96	—

PERICARDITIS WITH EFFUSION; ASPIRATION OF PERICARDIUM.

IMMEDIATE RELIEF AND ULTIMATE RECOVERY.

By T. H. BARTLEET, F.R.C.S.,

SURGEON TO THE GENERAL HOSPITAL, BIRMINGHAM.

DR. HARVEY, under whose care the case is, reports that the patient, Henry H—, aged twenty, had been for fourteen days suffering from acute rheumatism, but under his care for four days only. Symptoms of pericarditis were observed on Dr. Harvey's first visit. On Nov. 13th he found the patient suffering from urgent dyspnoea. There was a considerably increased area of cardiac dulness and marked failure of pulse. At noon on Nov. 13th Dr. Russell saw the patient with Dr. Harvey, and reports: Very feeble and rapid pulse; very rapid breathing; much dyspnoea.

Cardiac dulness extended to one inch to the right of the sternum and to the upper border of the first rib above, though here the dulness was not complete. The limit of the dulness on the left side was uncertain, owing to effusion in the chest. On auscultation, the heart-sounds were very feeble and distant; no friction-sound. There was a faint mitral bruit. The character of the heart-sounds was not affected by posture. The impulse of the heart could not be felt, otherwise than as an ill-defined movement of the chest-wall at the region of the apex, spreading over a space the size of a crown-piece. The history of cardiac disorder at a

former period suggested the possibility of pericardial adhesion.

At 8 P.M. on Nov. 13th aspiration was performed. A No. 2 aspirator needle, unguarded, was used. This was passed in the intercostal space between the fourth and fifth ribs, two inches to the left of the central line of the sternum. The needle was used as an exhausted needle; and directly the pericardium was penetrated, as shown by fluid freely passing into the aspirator, the point of the needle was pressed up against the chest-wall as closely as possible. The fluid flowed freely to fourteen ounces, and then stopped flowing. It was deeply tinged with blood, and deposited speedily a scanty coagulum, and subsequently a layer of blood débris. The supernatant fluid, still somewhat tinged, had a specific gravity of 1024.

During the latter part of the operation it was noticed by all present that the needle could be felt when held by the finger to be moved with the contraction of the heart. The patient was breathing deeply from pain or excitement, and thus prevented any cardiographic movements of the needle being seen. During and for half an hour after the operation the patient complained of severe aching pain. This ceased after a dose of fifteen minims of liq. opii. After the operation the line of dulness had receded to close upon the middle line of the sternum. The subclavian dulness had not disappeared, but was much less marked.

Nov. 14th.—Had passed a good night. Dr. Harvey thought the dulness a little increasing.

15th.—Reported considerably better. Lying down with ease; no dyspnoea; marked improvement in his look; pericardial dulness hardly reaches the right edge of the sternum. The first and second intercostal spaces are clear; third rib partly so.

Dec. 3rd.—Dr. Harvey reports that the pericardial dulness has nearly disappeared, and the pleuritic fluid has been mostly absorbed. Now that the heart-sounds can be plainly heard, a loud regurgitant mitral sound is developed.

10th.—Dr. Harvey reports to-day that the patient is able to sit up and walk about in the house.

Remarks.—This case shows the ease and safety with which paracentesis pericardii can be performed, for the relief of urgent symptoms resulting directly or indirectly from pericardial effusion. There was no difficulty in the operation itself, nor was there any subsequent symptom to mar the steady progress of the case to recovery. One point more is of especial interest—namely, that no peculiar or unfavourable importance need be attached to a free admixture of blood with the fluid withdrawn.

Birmingham.

A Mirror

OF

HOSPITAL PRACTICE,

BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum, tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

ST. THOMAS'S HOSPITAL.

CASES OF ANEURISM CURED BY DIGITAL COMPRESSION.

(Under the care of Mr. MAC CORMAC.)

AT page 693 we published the notes of a case of popliteal aneurism cured at the above-named hospital in twenty-three hours by means of digital compression. In the first of the subjoined cases pulsation was completely arrested in seventeen hours; but in the second case, which was traumatic, digital compression, although ultimately successful, was not so satisfactory as in the other cases. The mode of treatment adopted in the latter case may be contrasted with that which was employed in some cases of traumatic aneurism the notes of which we reported last week from Guy's Hospital. Two of these were instances of traumatic aneurism of the radial artery, and both were treated by opening the sac and tying both ends of the vessel. In one case the patient left the hospital cured in fourteen days, and in the

second case the patient was discharged cured on the sixth day after the operation. On the other hand, in the second of the subjoined cases, the patient remained in the hospital one month; and the aneurism was ultimately cured only after ninety-four hours of digital compression. Such compression, it is needless to say, could not be carried on except where there are a large number of assistants. It would appear, therefore, that, under some circumstances, the opening of the sac and ligature of the artery is not only a safe but a speedy mode of treating traumatic aneurism, especially in superficial arteries. Where there are plenty of willing assistants to carry on prolonged digital compression it may be an advantage to the patient to be saved the risks that attend, in a greater or less degree, all active surgical interference; and in this manner this patient may have been a gainer.

For the following notes we are indebted to Mr. Samuel Osborn, surgical registrar.

CASE 1. *Popliteal aneurism; digital compression; cure.*—J. W—, a labourer, aged forty, was admitted into Albert ward on October 10th, suffering from a popliteal aneurism in the left leg. The swelling was first noticed about the beginning of August, and was from the commencement the seat of great pain, for which the patient consulted a medical man, who recommended the application of poultices. As neither the pain nor the swelling subsided under this mode of treatment, the patient applied to another medical practitioner, who, however, advised a continuance of the treatment, but stated that if the swelling did not soon come to a "head," he would open it. The origin of the swelling the patient attributed to pushing heavy weights, sometimes as much as half a ton, with his feet. He had, however, acquired syphilis about fifteen years ago.

When he applied at the hospital for admission, a pulsating tumour, as large as a good-sized orange, and simulating very closely an abscess, was found occupying the entire left ham, and blocking up the whole of the popliteal space overlapping the biceps tendon on the outer side and the muscles forming the inner hamstring on the inner side. The skin appeared red and inflamed, and the covering over the aneurism was very thin, particularly over a small spot on the fibular side of the leg,—so much so that the danger of the tumour bursting seemed imminent. The pulsation was extremely violent, and was doubtless made worse by walking to the hospital.

Mr. MacKellar, the resident assistant-surgeon, who admitted the patient, ordered his immediate removal to the ward on a stretcher, fearing that he should have to ligature the femoral artery immediately. Fortunately at the time numerous students were about the hospital, and volunteers were easily obtained to carry on digital compression, which was commenced at 3 P.M. The following day, at 8 A.M., no pulsation was perceptible. Compression was, however, kept up for seven hours longer, and a watch was continued until 12.15 A.M. on the 12th, after a space of forty-five hours and fifteen minutes.

CASE 2. *Traumatic ulnar aneurism; digital compression; cure.*—J. J—, a sadler, aged thirty, was admitted into Leopold ward on May 1st with an aneurism of the ulnar artery. One month ago he cut his left hand with a knife; the wound was situated on the palmar surface at the junction of the hand and the forearm, and extended transversely across the lower end of the ulnar artery. There was at the time no hæmorrhage to speak of; the wound healed up quickly, but a swelling remained in that situation, which one week before coming to the hospital began to throb and to be very painful.

On admission, a swelling about the size of a walnut existed over the ulnar artery, just where it divides to form the superficial palmar arch, pulsation in it being very distinct, and the skin covering it extremely thin.

After digital compression had been applied to the brachial artery for thirty-two hours, all pulsation was found to have ceased, and the aneurism felt tolerably firm. On May 6th, however, distinct pulsation was again noticeable in the aneurism, and digital compression was again applied to the brachial artery for thirty-eight hours, until all pulsation had ceased. On the 11th pulsation again returned, and was again arrested by compression on the radial artery. Digital compression was applied to this vessel for twenty-four hours, after which time the tumour remained solid, and the patient left the hospital cured on June 1st.

MIDDLESEX HOSPITAL.

SEVERE UTERINE HÆMORRHAGE; SO-CALLED HYDATIDS; RECOVERY.

(Under the care of Dr. EDIS.)

THE following case illustrates a form of menorrhagia which is frequently overlooked at the time. At page 694 we published a case of menorrhagia which occurred in a person the subject of Bright's disease; and at page 761 another case was recorded in which the menorrhagia was due to a granular condition of the os and the cervix of a gravid uterus, which was cured by the application of nitrate of silver and the internal administration of ergot, without any interference with the progress of utero-gestation. Although these cases do not illustrate all the forms of menorrhagia, they serve to teach one fact of great importance—namely, that every case of menorrhagia should be made the subject of special study, and should be treated according to the indications peculiar to it.

C. J—, aged seventeen; single. Catamenia regular until eight months since, when they suddenly ceased. She had taken pills, pennyroyal, hiera picra, and various other remedies, with the view to bring on "the courses," the patient more than suspecting that there was good reason for their non-appearance. On August 4th, whilst out walking, she experienced a sharp attack of hæmorrhage from the vagina, unaccompanied by pain or other inconvenience. She was assisted home, and the hæmorrhage ceased when she lay down.

On August 12th she went for a long drive in a cart. She had just returned, when she had a second and much more severe attack of hæmorrhage. Her friends brought her in a cab at once to the hospital, when Dr. Langmore, the resident obstetric assistant, in the absence of Dr. Hall Davis, sent for Dr. Edis. When first seen, the patient was in a state of extreme syncope. The face was blanched, the expression anxious, and the pulse small, soft, jerking, and over 120. The patient was restless, and the extremities were cold.

On examination the mammae were found to be well developed, the areolæ very dark, and the follicles enlarged. A thin milky fluid could be expressed from the nipple. The abdomen was enlarged, and occupied by a tense ovoid elastic tumour in the mesial line, extending to an inch and a half above the umbilicus. No foetal movements, nor, in fact, any trace of a foetus could be detected, nor were any foetal heart-sounds audible. On examining per vaginam, the os uteri was detected high up, posteriorly, soft and dilatable. There was little or no cervical portion, and on passing the tip of the finger through the os, nothing beyond clots could be felt; no presentation of any foetal structure.

The hæmorrhage having now been checked, and the patient's exhausted condition not warranting any immediate interference, an opiate was administered, brandy, milk, &c., being given to restore her strength, and she was carefully watched.

At midnight, the patient having turned in bed, a gush of bright blood ensued, followed very shortly by another. Very slight uterine contractions were present. The pulse had moderated to 96; the temperature 100° F. A No. 3 Barnes's bag was introduced within the os, and inflated.

At 5 A.M. renewed hæmorrhage took place; uterine action more marked, but pains weak and irregular. A larger bag was then inserted, and ergot was given. After a short time a further examination was made, when two or three small vesicular bodies were extruded with some clots; and the nature of the case, which had been more than suspected, was made evident. The os being now fairly dilated, a single blade of the forceps was passed into the uterus, and nearly a washhand-basinful of so-called hydatids was extracted.

Brandy, milk, and ergot were administered, and with the exception of some slight vomiting, inability to void urine, and a few rigors, the patient made a good recovery.

A small, firm mass of placental structure was all that was found to indicate the fact of utero-gestation having taken place.

The specimen presented the ordinary form of hydatidiform or cystic degeneration of the villi of the chorion—viz., botryoidal or grape-like clusters of small cysts, varying in size from a pea to a large marble.