after the accident there was no change in the boy's condition, total unconsciousness remaining. At the end of twenty-four hours the pulse was just perceptible, and, as it was earnestly requested by the parents, permission was granted for his removal home, fresh dressings in the shape of lint soaked in cold water being previously applied; these were renewed every day for a week; each time there was an escape of brain from the large wound and ear. Altogether more than two ounces of brain substance were lost. The temperature ranged from 99° to 101°, and the pulse from 100 to 120. On the 27th slight consciousness returned, the patient opening his eyes when spoken to. At this time I perceived that there was total paralysis of the right side; the urine passed involuntarily, and the bowels had to be relieved by enemata. On April 4th and on several subsequent occasions Dr. Gisborne, senior consulting surgeon to the Derbyshire General Infirmary, saw the child with me, and he regarded it as the most extraordinary case he had ever known. On May 19th, eight weeks after the accident, the little patient was quickly getting well; the wound on the head was two inches long and one inch wide. There were two fragments of parietal bone projecting; these, though loose, were not removed at the time on account of the close proximity of the middle meningeal artery, which could be seen pulsating, and raising the pieces with each pulsation. There was great irregularity over the surface of the left temporal bone, which somewhat projected. The pupils were alike and contracted to light. The child sat up, and ate and slept well. The right leg had regained some power, and the arm a little; and although he could only repeat one or two words said to him, he looked fairly intelligent. On the 23rd one of the fragments was removed; it was diamond-shaped, measured an inch and a quarter in its greatest diameter, and weighed two scruples. On my visit on the 30th, I found the little fellow out playing in a field, the paralysis having almost entirely disappeared; his speech was better, and in every way he was rapidly improving.

P.S. (July).—Since the above was written I have removed another large piece of the left parietal bone, and there is another portion which I shall take away in a few days. The child has gone on well throughout, has had no fit or any bad symptom. The paralysis of the right upper extremity has entirely disappeared; he drags the right leg slightly in running, but is daily acquiring more power in it. He articulates perfectly, plays in the park with his brothers and sisters; he eats, digests, and sleeps well, and appears quite well. The father of the boy keeps a lodge in the park of T. W. Evans, Esq., M.P. for South Derbyshire; and the kindness of that gentleman in supplying everything necessary, during the entire time, has had much to do with the child's recovery.

Duffield.

DEPRESSED AND COMMUNUTED FRACTURE OF THE SKULL; RECOVERY.

By F. A. SAUNDERS, L.R.C.S., L.R.C.P. Ed., &c.

ANDREW S—, aged ten years, of Kingsbarns, Fifeshire, was, on September 1st, 1882, playing near a large farm cart, when the horse suddenly moved and the wheel crushed his head between it and the wall, which rendered him perfectly unconscious for a few minutes, but after his arrival at home he soon recovered consciousness. I saw the patient two hours after the injury, and found on examination, under chloroform, an incision extending from the left parietal eminence to the right superciliary ridges. The scalp was detached from its connexions, and turned down over the left ear, leaving bare the left half of the cranium, and exposing a very considerable surface of bone. The periosteum was completely stripped off the frontal bone for the distance of three inches. The supra-orbital and frontal arteries were torn across and twisted; the left temporal muscle was torn from its connexions with the bone and hanging loose, with the posterior temporal artery also torn and twisted. Upon reflecting the flap of skin back over the ear I felt and saw that the left parietal bone, one inch above the squamous suture, was fractured, comminuted, and depressed, and a splinter of bone was easily detached. He was then in bed, conscious but restless, moaning and suffering from great pain in the head. The pulse was quick (120); temperature 102.6°. The pupils dilated, but contracted slowly under the stimulus of light. Considering his age, and as there were

no prominent symptoms of compression, having ligatured a branch of the anterior temporal artery that was bleeding, I simply applied cold water cloths to the wound, and determined to wait before I brought the flaps of skin in contact. The next morning, September 2nd, I asked Dr. J. Moir of St. Andrews to see the case with me, which he kindly did. The temperature had fallen to 101.2°, the pulse to 108. He was quite conscious, and did not complain of much pain. We then determined not to trephine, but to follow out the expectant plan of treatment.

Dr. Moir having placed the patient under chloroform, I removed the greater portion of the temporal muscle and another small splinter of bone, and then brought the flaps of skin together as much as possible, taking care that perfect drainage was established. Ice was applied continuously for three weeks. During the first seven days the temperature oscillated between 101° and 102.8°. On the eighth day, when it rose to 103°, twenty grains of quinine were administered. On the ninth day the temperature was 103.4°, and twenty grains of quinine were again administered. On the tenth day the temperature was 104°; twenty grains of quinine given. On the eleventh day the temperature was 104.2°; the same quantity of quinine administered. On the twelfth day the temperature began to fall, going down to 102°, and continued between 102° and 101° until Sept. 28th, when it kept at about 99° until Nov. 2nd. On that date I opened up a sinus by a two-inch incision, having felt dead bone, and removed three pieces of necrosed bone. The temperature then went up to 103° for two days, after which, in the following week, two similar pieces of bone came away. On Nov. 27th the wounds had healed sufficiently for me to leave the dressings to the care of the mother. The boy has done well since. The interesting features in this case are:—First, that although there was such a large surface of bone stripped of its periosteum there was no exfoliation; secondly, it is another example of Sir A. Cooper's, Abernethy's, and Dupuytren's advice, not to interfere unless there are marked symptoms of compression accompanying a depressed fracture.

Craill, Fifeshire.

A Mirror

OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nu. la autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

GUY'S HOSPITAL.

CASES OF DISEASE OF THE UPPER JAW.

(Under the care of Mr. BRYANT.)

CASE 1. *Periosteal (?) Sarcoma of Anterior Surface of Upper Jaw; Removal; Recovery.* (Reported by Mr. Franklyn Brown.)—Edith C—, aged nine years, was admitted into Lydia ward on June 30th, 1881. The patient's father, aged thirty, was healthy, but her mother, aged thirty-two, had suffered from rheumatism. Six months before admission the patient received a blow on the right cheek, and soon afterwards noticed a small swelling, which for a time remained stationary, but for the last three weeks it had increased. About two months before admission it was punctured, but with no result.

On admission, there was a slight swelling on the right cheek over the malar bone, deeply placed and movable over the bone "with an elastic feel." The superficial structures were freely movable over the tumour. The finger could be placed between the tumour and the teeth. Size about .75 in. by .5 in. Nostril free, lacrymal duct free. On the right side she had permanent incisors and the six-year old molar, also the temporary canine and temporary second molar. The first molar had been shed. On July 1st the tumour was removed from inside by scissors. It was easily separated from the bone. Iodine water was applied and a sponge inserted. The tumour was solid and a good deal of muscle was removed. On section it had a uniform translucent