PROCEEDINGS OF THE NINETEENTH ANNUAL MEETING,
HELD AT NEW YORK, N. Y., DECEMBER 27, 28, AND 29,
INCLUDING PROCEEDINGS OF THE EIGHTEENTH AN-
NUAL MEETING OF THE CORDILLERAN SECTION, HELD
AT STANFORD UNIVERSITY, CALIFORNIA, DECEMBER
28 AND 29, 1906

EDMUND OTIS HOVEY, *Secretary*

CONTENTS

	Page.
Session of Thursday, December 27.....	559
Report of the Council.....	559
Secretary's report	559
Treasurer's report	564
Editor's report	568
Librarian's report	568
Election of officers.....	569
Election of Fellows.....	570
Memoir of William Buck Dwight; by F. J. H. Merrill.....	571
Memoir of Samuel Lewis Penfield (with bibliography); by Joseph P. Iddings	572
Memoir of Israel C. Russell (with bibliography); by Bailey Willis...	582
Memoir of Nathaniel Southgate Shaler (with bibliography); by John E. Wolff	592
Graded surfaces [abstract]; by F. P. Gulliver.....	609
Session of Friday, December 28.....	611
Seventeenth annual report of the Committee on Photographs.....	611
Resolutions concerning retiring Secretary and Treasurer.....	611
Resolution concerning investigation of volcanoes.....	613
Memorial of A. R. C. Selwyn [abstract]; by H. M. Ami.....	614
Dome structure in conglomerate [abstract]; by Ralph Arnold.....	615
Relations between climate and terrestrial deposits [abstract]; by Joseph Barrell	616
Personal reminiscences of Sir William E. Logan [abstract]; by Robert Bell	622
Session of Saturday, December 29.....	623

	Page
Report of the Auditing Committee.....	623
Permo-Carboniferous climatic changes in South America [abstract]; by David White	624
Controlling factors of artesian flows [abstract]; by Myron L. Fuller..	626
Volcanoes of Colima, Toluca, and Popocatepetl [abstract]; by E. O. Hovey	635
Probable age of the Meguma (gold-bearing) series of Nova Scotia [abstract]; by J. Edmund Woodman.....	636
Lebanon glacier [abstract]; by G. Frederick Wright.....	637
Ice present during the formation of glacial terraces [abstract]; by F. P. Gulliver.....	640
Glacial lake Memphremagog [abstract]; by C. H. Hitchcock.....	641
Earth-flows at the time of the San Francisco earthquake [abstract]; by Robert Anderson.....	643
Genetic relations of some granitic dikes; by Alfred C. Lane.....	644
Ophitic texture [abstract]; by A. C. Lane.....	648
Charles Willson Peale's painting, "The exhuming of the first American mastodon"; by Arthur Barneveld Bibbins.....	650
Resolution as to division of meetings into sections.....	654
Register of the New York meeting, 1906.....	655
Session of the Cordilleran Section, Friday, December 28, 1906.....	656
Notes on the structure of the Santa Cruz range, California [abstract]; by J. F. Newsom.....	657
Notes on the topography of the Seward peninsula, Alaska [abstract]; by J. F. Newsom.....	657
Transportation of detritus by Yuba river [abstract]; by G. K. Gilbert.	657
Crocidolite-bearing rocks of the California coast ranges [abstract]; by George D. Louderback and Wm. J. Sharwood.....	659
Primitive characters of American Triassic Ichthyosaurs [abstract]; by John C. Merriam.....	659
Origin of South American bears [abstract]; by John C. Merriam...	660
Two mountain ranges of southern California [abstract]; by W. C. Mendenhall	660
Session of the Cordilleran Section, Saturday, December 29, 1906.....	661
Notes on the geology of the Mount Hamilton quadrangle [abstract]; by J. F. Newsom and Roderic Crandall.....	661
Cretaceous stratigraphy of the Santa Clara Valley region [abstract]; by Roderic Crandall.....	661
Physiographic features of south central Oregon [abstract]; by G. A. Waring	662
General geological features of the Truckee region east of the Sierra Nevada [abstract]; by George D. Louderback.....	662
Register of the meeting of the Cordilleran Section.....	670
Accessions to the library from October, 1906, to October, 1907.....	671
List of Officers and Fellows of the Geological Society of America.....	681
Index to volume 18.....	693

SESSION OF THURSDAY, DECEMBER 27

The Society was called to order by Acting President William M. Davis, in the absence of President Israel C. Russell, deceased, at 4.30 o'clock p m, in the lecture hall of Schermerhorn hall, Columbia University. A brief but cordial address of welcome was delivered by President Nicholas Murray Butler, of Columbia University, to which response was made by Acting President Davis.

The report of the Council was called for, and was presented by the Secretary, Herman Leroy Fairchild, in print, as follows:

REPORT OF THE COUNCIL

To the Geological Society of America,

in Nineteenth Annual Meeting assembled:

The stated annual meeting of the Council was held at Ottawa conjointly with the meeting of the Society. It has not been found necessary to call any special meeting, but some business relating to nomination of officers and recommendation for fellowship has been done by correspondence.

The details of administration for the eighteenth year in the history of the Society are given in the following reports of the officers:

SECRETARY'S REPORT

To the Council of the Geological Society of America:

Meetings.—The proceedings of the Ottawa Winter Meeting, 1905, will be recorded in the closing brochure of volume 17 of the Bulletin.

Membership.—Since the last printing of the list of Fellows four Fellows have been removed by death: W. B. Dwight, S. L. Penfield, I. C. Russell, and N. S. Shaler. The names of 16 Fellows, elected at the Ottawa meeting, have been added to the list, making the present enrollment 283, or 12 more than at the last printing. On the ballot now before the Society are the names of 14 candidates, while several nominations are awaiting action by the Council.

During recent years our membership has been increasing. In November, 1901, it was 247; in January, 1904, it was only 253. The present enrollment, with the addition of the 14 candidates on the current ballot, will make a total within three of 300. Another year will carry the list beyond 300. For several years no name has been dropped from the roll for non-payment of dues, and only one resignation has been accepted. The increase in membership has not been due to any lowering of the standard

for admission, but to the reputation of the Society and the attraction of its fellowship for students of earth-science.

Distribution of Bulletin.—At this date 468 pages of volume 17 have been distributed, and the remaining brochures are approaching completion. The irregular distribution of the Bulletin during the past year has been as follows: Complete volumes sold to the public, 58; sold to Fellows, 14; brochures sent to supply deficiencies, 57; sold to the public, 20; sold to Fellows, 44. One copy of volume 16 has been donated and three copies bound for the use of officers and the Library. Three complete sets of the back volumes have been sold to libraries, and a partial set to a Fellow. Not many complete sets will be sold in the future, as the cost is becoming large, the greater libraries are generally supplied, and the sets belonging to deceased Fellows will be thrown on the market.

Bulletin sales.—The receipts from the sale of the Bulletin during the past year appear in the following table:

Receipts from Sale of Bulletin, December 1, 1905, to December 1, 1906

	Complete volumes.			Brochures.			Grand total.
	Public.	Fellows.	Total.	Public.	Fellows.	Total.	
Volume 1..	\$10 00	\$9 00	\$19 00				\$19 00
Volume 2..	10 00	4 50	14 50		\$2 75	\$2 75	17 25
Volume 3..	10 00	4 00	14 00		3 10	3 10	17 10
Volume 4..	10 00	7 00	17 00	\$0 35		35	17 35
Volume 5..	10 00	4 00	14 00	45		45	14 45
Volume 6..	10 00	4 00	14 00				14 00
Volume 7..	10 00	4 00	14 00				14 00
Volume 8..	10 00	4 00	14 00				14 00
Volume 9..	10 00	4 00	14 00				14 00
Volume 10..	15 00	4 00	19 00				19 00
Volume 11..	10 00	4 50	14 50				14 50
Volume 12..	15 00	4 00	19 00		1 20	1 20	20 20
Volume 13..	15 00	4 50	19 50	4 80	4 95	9 75	29 25
Volume 14..	30 00		30 00	80	2 25	3 05	33 05
Volume 15..	25 00	4 50	29 50	1 15	45	1 60	31 10
Volume 16..	215 00		215 00	2 55	1 20	3 75	218 75
Volume 17..	200 00		200 00				200 00
Volume 18..	20 00		20 00				20 00
<hr/>							
				\$10 10	\$15 90	\$26 00	\$727 00
Index	\$635 00	\$66 00	\$701 00				9 25
	7 00	2 25	9 25				
				\$10 10	\$15 90	\$26 00	\$736 25

Receipts for the fiscal year.....	\$736 25
Previous receipts, to December 1, 1905.....	8,866 09

Total receipts to date.....	\$9,602 34
Charged and uncollected.....	100 40

Total Bulletin sales to date.....	\$9,702 74
-----------------------------------	------------

The bills to regular subscribers for volume 17 have not been sent, and the above table includes only advance payments. In another year the total receipts from the sale of the Bulletin will considerably exceed \$10,000.

The publication cost of volumes 1-16 is \$30,629.55, the average cost being \$1,914.35. This does not include the expense of distribution. The receipts to date from the sale of the same volumes are \$9,382.34, or an average of \$586.40 per volume. The receipts from sale of these volumes approach one-third the cost of publication; but for the past six years the annual receipts have averaged \$727.05, which is considerably more than one-third the expense of publication. The present income from sales of the Bulletin and from interest on funds, taken together, amounts to nearly 70 per cent, or more than two-thirds of the cost of the Bulletin. This would seem to be good financing for a society of small membership, depending on moderate dues for its income.

Exchanges.—Three addresses have been added to the list of exchanges—one in Mexico, one in Vienna, and one in Manila. The full list is given with the list of Library accessions, at the close of each volume of the Bulletin.

Expenses.—The following table gives the cost of administration and of Bulletin distribution during the past year:

EXPENDITURE OF SECRETARY'S OFFICE DURING THE FISCAL YEAR ENDING NOVEMBER
30, 1906

Account of Administration

Postage and telegrams.....	\$27 90
Expressage	6 36
Printing (including stationery).....	109 66
Meetings (expenses of Cordilleran Section).....	14 00
Total	\$157 92

Account of Bulletin

Postage	\$81 40
Expressage and freight	109 86
Wrapping material	4 13
Addressograph links	1 91
Binding three copies of volume 15.....	3 00
Printing	3 25
Collection of checks	3 50
Total	207 05
Total expenses for the year.....	\$364 97

As this is my last report, a few words reminiscent and historical may not be amiss.

The character of the Society, the policy, and the formalities of the Secretary's office were established during the first two years, under the wise guidance of Professor Stevenson. As Editor, Doctor McGee gave the Bulletin its exemplary form, only one volume of which had been printed and sent to the Fellows when the writer was called to the Secretary's office, January, 1891. The questions of the distribution and sale of the Bulletin had yet to be considered.

The first circular advertising the Bulletin and soliciting subscriptions from libraries was dated January, 1892. From time to time efforts have been made to enlarge the subscription lists, and some advertising has been done, but with little apparent effect. The character of the publication is such that the demand can not be greatly stimulated. Where desired, the Bulletin sells itself.

Through the Editor and Secretary, the Society has been conducting a publishing business, and the Secretary has been running a sales-room by correspondence and selling the Bulletin in all parts of the world, chiefly, of course, in America.

After deliberation and correspondence, a list of "Exchanges" was made in 1891, with the purpose of placing the Bulletin at focal points of scientific influence over the world. The first list numbered 68 (see Bulletin, vol. 3, page 468). The obtaining of library material in return for the Bulletin was not an object, but was inevitable. The accumulation of exchange material led to the making of a contract in 1894 with the Case Library of Cleveland for the care of the Society's Library (see Bulletin, vol. 6, page 427), and in 1898 Professor Cushing consented to act as Librarian. (His first report is in Bulletin, vol. 10, page 422.)

The Constitution of the Society followed conventional practice in providing for several classes of members: Fellows, Correspondents, and Patrons. At one time the Council made an earnest effort to select a complimentary list of Correspondents, but after compiling and sifting many proposed names, the effort was abandoned and has never been revived. It is quite as well. One strong element of success of the Society has been its democratic spirit, and it is better that all its membership should stand on an equality. If any foreign geologist wishes to place F. G. S. A. after his name, he can do so by becoming a Fellow and paying the regular charges, as the territorial limitations to fellowship were removed in 1894.

Perhaps it is also fortunate that the Society has had no patrons, but has succeeded entirely through the character and the contributions of its fellowship (see the Treasurer's report, following).

The Secretary has preserved in a file of large scrap-books every scrap of printing, outside of the Bulletin, that the Society has ever had, along with all other material that could conveniently be pasted. These books, with the Council minutes and the Bulletin, make a perfect record of the public and the internal work of the Society. All the correspondence (with very few exceptions, of confidential papers) has been preserved. The future historian of the early years of what will some time be the "old" society should find nothing missing.

The writer has never missed a meeting of the Society, nor any meeting of the Council since the winter meeting in 1890. For the past sixteen years he has given a large part of his time and thought to the work of the secretaryship. It has been a labor of love, and the approval of the Fellows has been sufficient reward. To a conscientious executive officer there is a satisfaction in the control of affairs and the direction of policy, as in any work, when these are accepted and commended by those concerned. Unanimous reelection year after year has been interpreted as such approval.*

It is possible that an administration with more initiative and vigor would have produced greater results; but with our scattered membership, of strong individualities and somewhat diverse interests, it seemed safer to pursue a conservative course and let affairs move quietly and smoothly until the Society had acquired strength and prestige. The perfect unity and harmony in the Society and the Council during all the 18 years is a happy fact.

The work of the Secretary is intimately bound with that of the Treasurer and Editor and they are largely to be credited with the success of the Society. For many years of loving and laborious service the Society owes a debt to Stanley-Brown and I. C. White that only appreciation can repay. For his successor the writer bespeaks the continued sympathetic support of the fellowship, which is the best help to his success.

In surrendering office your Secretary sincerely thanks every Fellow for uniform kindness, courtesy, and support. No incident has ever occurred of character so unpleasant as to leave any remembrance. The recollection of the sixteen years of service to the Society will remain the most unalloyed and satisfying memory of a busy life.

Respectfully submitted.

H. L. FAIRCHILD,
Secretary.

ROCHESTER, N. Y., *December 10, 1906.*

* In fifteen elections by secret ballot the total vote has failed of unanimity only by a single vote at one election. Doctor White has always had a unanimous vote.

TREASURER'S REPORT

To the Council of the Geological Society of America:

The Treasurer herewith submits his annual report for the year ending December 1, 1906.

On this latter date two (2) Fellows were liable to be dropped from the roll for non-payment of dues, in accordance with section 3, chapter 1, of the By-laws; six (6) were delinquent for two years, while thirty-one (31) had not yet paid for the present year.

Since December 1 8 of these have paid, leaving only 4 delinquent for 1905 and 1906 and 25 delinquent for 1906.

Five (5) Fellows, A. P. Low, A. P. Brown, R. A. Daly, S. H. Ball, and F. G. Clapp, have commuted for life by the payment of \$100.00 each, thus increasing the total Life Commutations to eighty (80) at the present time.

The Permanent Publication Fund (only interest of which can be used for the current expenses of publication) was both decreased and increased during the year. This Fund amounted to \$9,300.00 at the date of the last statement, of which \$300.00, representing three (3) bonds of the Tunnelton, Kingwood and Fairchance railroad, matured during the year and were redeemed under date of July 28, 1906, at par and accrued interest. To offset this reduction of invested funds, 10 more shares in the Ontario Apartment House Company of Washington, D. C., were purchased by myself and Mr Emmons, the majority of the Committee on Investments, under date of March 24, 1906, thus increasing this fund to exactly \$10,000.00 par value, half of which yields 6 per cent interest and the other half 5 per cent, or a total of \$550.00 annually. The entire cost of the Publication Fund was \$9,342.50, so that the rate of interest on the actual investment is nearly 6 per cent. The total interest received during the year from these permanent investments (\$517.33), increased by that (\$64.38) on monthly balances from the Security Trust Company of Rochester, New York, amounts to \$581.71, while \$120.00 more will accrue (on the 40 shares of the Ontario Apartment House Company stock) January 1, 1907.

A table showing the growth of the Publication Fund and these interest items from the beginning of the Society's financial history will prove both instructive and useful.

The first purchase on account of permanent investments was made during the incumbency of my predecessor in the Treasuryship, Professor H. S. Williams, on April 1, 1891, when a \$1,000.00 seven per cent bond of Tioga township, Kansas, was bought at a cost of \$1,140.25; so that when the present Treasurer assumed the office, January 1, 1892, the in-

vested funds had a par value of \$1,000.00 and an annual interest return of \$70.00.

The following table exhibits the growth of the invested funds and interest items, the latter being increased each year to a small extent by the 4 per cent received on monthly balances from the Security Trust Company of Rochester, New York, where all the cash not needed in paying current accounts has been kept for the last 15 years:

Year.	Cost of investment.	Par value.	Annual interest from investments.	Interest on bank balances.
1889.....				\$16 63
1890.....				51 34
1891.....	\$1,140 25*	\$1,000 00*	\$35 00	65 32
1892.....	900 35†	900 00†	92 50	16 72
1893.....	700 00†	700 00†	135 50	17 65
1894.....			135 50	23 46
1895.....	304 00‡	300 00‡	162 00	8 06
1896.....			168 00	23 00
1897.....	100 00†	100 00†	170 50	49 84
1898.....	1,976 25	2,000 00	275 50	32 98
1899.....			273 00	60 28
1900.....			273 00	97 23
1901.....	1,000 00	1,000 00	282 63	100 61
1902.....			236 11	151 83
1903.....	2,000 00	2,000 00	248 00	148 17
1904.....	2,366 25	3,000 00	483 00	93 98
1905.....	1,000 00	1,000 00	448 00	65 64
1906.....	1,000 00	1,000 00	517 33	64 38
Total	\$12,487 10	\$13,000 00	\$3,935 57	\$1,087 12

Adding the total of interest received on investments, \$3,935.57, to the total received from bank deposits, \$1,087.12, gives a grand total of \$5,022.69 received since the organization of the Society, all of which except \$168.27 has been received during the last 15 years.

At the second annual meeting of the Society, held in New York city, December 26, 1889, the Council, among other recommendations, asked that an effort be made to raise a Publication Fund of \$10,000.00. This demand not having been met, the Council renewed the request in submitting its annual report at the Columbus meeting of December 29, 1891. It was evidently the expectation of the Council that some wealthy friends of the Society would donate the amount requested in a short time, but the hope failed of fruition. The present Treasurer, who was elected at the Columbus meeting (1891) never made any serious effort to raise a Publication Fund in the way suggested, because he preferred to see what

* Kansas bond redeemed at \$1,081.80, April 1, 1904.

† \$1,700 in Cosmos Club bonds redeemed at par May 16 and 25, 1901.

‡ T. K. & F. bonds redeemed July 28, 1906, at par and accrued interest.

the Society could do for itself by careful attention to correct business methods, and hence it is with no small degree of satisfaction that the retiring Treasurer can at this meeting turn over to his successor and to the Society the full amount (\$10,000.00) asked for 15 years ago, invested in first-class interest-bearing securities, with all accrued bills paid to date (except 3 bills for \$572.00 just received from President) and \$2,000.00 in the treasury. The above table illustrates the value of small savings and points the way to a continued increase of the Society's financial resources. The total expenses of the Treasurer's office (printing, stationery, and postage) for the last 15 years amount to \$318.23, or an average of \$21.23 per annum.

The securities now owned by the Society are as follows:

	Par value
March 17 and 25, 1898, two Texas and Pacific Railroad 1st mortgage 5 per cent bonds, cost \$1,976.25.....	\$2,000 00
February 6, 1901, ten shares of the capital stock of the Iowa Apartment House Co., Washington, D. C., cost \$1,000.00.....	1,000 00
April 1, 1903, twenty shares of the capital stock of the Ontario Apartment House Co., Washington, D. C., cost \$2,000.00.....	2,000 00
April 11, 1904, three second mortgage 5 per cent bonds of the U. S. Steel Corporation, cost \$2,366.25.....	3,000 00
May 12, 1905, ten shares of the capital stock of the Ontario Apartment House Co., Washington, D. C., cost \$1,000.00.....	1,000 00
March 24, 1906, ten shares of the capital stock of the Ontario Apartment House Co., Washington, D. C., cost \$1,000.00.....	1,000 00
Total cost, \$9,342.50; total par value.....	\$10,000 00

The general financial condition of the Society is exhibited by the tabular statement (page 567), showing the receipts and disbursements for the year ending December 1, 1906.

This *balance* is distributed as follows:

In the Security Trust Co. of Rochester, N. Y.....	\$1,678 47
In the Bank of Monongahela Valley, Morgantown, West Va.....	318 94
Total	\$1,997 41

Respectfully submitted.

I. C. WHITE,
Treasurer.

MORGANTOWN, WEST VA., *December 20, 1906.*

Statement of Receipts and Expenditures

RECEIPTS		EXPENDITURES	
Balance in the treasury December 1, 1905.....	\$1,609 26	Total receipts brought forward.....	\$6,028 72
Fellowship fees 1903 (4).....	\$40 00	Secretary's office:	
" " 1904 (7).....	70 00	Administration	\$157 92
" " 1905 (31).....	310 00	Bulletin	297 05
" " 1906 (171).....	1,710 00	Allowance	500 00
" " 1907 (1).....	10 00		<u>\$864 97</u>
Initiation fees (16).....	2,140 00	Treasurer's office:	
Life commutations (5).....	160 00	Postage	17 50
Interest on investments:	500 00	Librarian's office	8 71
Tunnelton, Kingwood and Fair-		Publication of Bulletin:	
chance R. R. bonds.....	\$18 00	Printing	\$1,392 13
Iowa Apartment House Co. stock..	60 00	Engraving	494 45
Texas and Pacific R. R. bonds.....	100 00	Editor's allowance (per-	
Ontario Apartment House Co. stock	189 33	sonal and office ex-	
U. S. Steel Corporation bonds.....	150 00	penses)	250 00
Interest on deposits with Security			<u>2,136 58</u>
Trust Company of Rochester, N. Y.	64 38	Investments, ten shares Ontario Apart-	
		ment House Co. stock.....	1,000 00
Investments (Tunnelton, Kingwood and Fair-	581 71	Miscellaneous	3 55
chance bonds) redeemed	301 50		<u>4,031 31</u>
Sales of publications	736 25	Total expenditures	
Total receipts carried forward.....	\$6,028 72	Balance cash on hand December 1, 1906..	\$1,997 41

EDITOR'S REPORT

To the Council of the Geological Society of America:

The Editor was able to make only a preliminary report to the Council at the meeting held in December, 1906, but full statistics were prepared for both volumes 16 and 17 before the latter volume was closed.

J. STANLEY-BROWN,
Editor.

	Average. Vols. 1-12.	Vol. 13.	Vol. 14.	Vol. 15.	Vol. 16.	Vol. 17.	Vol. 18.
	pp. 577. pls. 43.	pp. 583. pls. 58	pp. 609. pls. 65.	pp. 636. pls. 59.	pp. 636. pls. 94.	pp. 785. pls. 84.	pp. 717. pls. 74.
Letter-press.....	\$1,675 14	\$1,647 12	\$1,657 50	\$1,661 21	\$1,817 03	\$2,087 98	\$2,015 68
Illustrations.....	327 62	477 27	431 21	457 76	706 97	608 66	\$486 22
	\$1,902 66	\$2,124 39	\$2,088 71	\$2,118 97	\$2,524 00	\$2,696 66	\$2,501 90
Average per page.....	\$3 30	\$3 64	\$3 43	\$3 33	\$3 96	\$3 37	\$3 42

Classification

Volume.	Areal geology.	Physical geol- ogy.	Glacial geology.	Physiographic geology.	Petrographic geology.	Stratigraphic geology.	Paleontologic geology.	Economic geol- ogy.	Official matter.	Memorials.	Unclassified.	Total.
	Number of pages.											
1.....	116	137	92	18	83	44	47		60	4	4	593 + xii
2.....	56	110	60	111	52	168	47	9	55	1	7	662 + xiv
3.....	56	41	44	41	32	158	104		61	15	1	541 + xii
4.....	25	134	38	74	52	52	14		47	32	2	458 + xii
5.....	138	135	70	54	28	51	107		71	14	9	665 + xii
6.....	50	111	75	39	71	99	1		63	25	4	538 + x
7.....	38	77	105	53	40	21	123	4	66	28	13	558 + x
8.....	34	50	98	5	43	67	58	14	79	8		446 + x
9.....	2	102	138		44	28	64	16	64	12		460 + x
10.....	35	33	96	37	59	62	68	28	84	27	17	534 + xii
11.....	65	110	21	10	54	31	188	7	71	60	46	651 + xii
12.....	199	39	55	53	24	98	5	5	70	2		538 + xii
13.....	125	17	13	24	28	116	42	4	165	32	29	583 + xii
14.....	48	47	48	59	183	118	22	1	80	14	1	609 + xii
15.....	26	124	3	94	36	267			77	17	3	636 + x
16.....	64	111	78	30	102	141	19		67	22	15	636 + xii
17.....	49	161	41	84	47	294	27		71	9	2	785 + xiv
18.....	16	164	141	5	29	246	5		68	40	3	717 + xii

(OLD SPRING HARBOR, N. Y., December 20, 1907.

LIBRARIAN'S REPORT

To the Council of the Geological Society of America:

Accessions to the Library for the past year were duly acknowledged and catalogued, and the list of accessions to September 1 compiled and sent to the Secretary for incorporation in volume 17 of the Bulletin.

There are now 3,000 catalogued numbers in the Library, some 2,000 of which are bound volumes, for the most part sets of the publications of the exchanges of the Society. The binding, which is done by the Case Library, amounts to about 125 volumes a year.

The expense of this office the past year is as follows:

To postage	\$1 46
To express	2 25
To clerk hire	5 00
	<hr/>
	\$8 71

Respectfully submitted.

H. P. CUSHING,
Librarian.

CLEVELAND, OHIO, *December 1, 1906.*

On motion of the Secretary, it was voted to defer the consideration of the Council report to the following day.

As the Auditing Committee to examine and report upon the accounts of the Treasurer, the Society elected A. P. Coleman and N. M. Fenneman.

ELECTION OF OFFICERS

Acting President Davis then announced the result of balloting for officers for 1907, as canvassed by the Council, and declared the following officers elected.

President:

CHARLES R. VAN HISE, Madison, Wis.

First Vice-President:

J. S. DILLER, Washington, D. C.

Second Vice-President:

A. P. COLEMAN, Berkeley, Cal.

Secretary:

EDMUND OTIS HOVEY, New York city.

Treasurer:

WILLIAM BULLOCK CLARK, Baltimore, Md.

Editor:

JOSEPH STANLEY-BROWN, Cold Spring Harbor, N. Y.

Librarian:

H. P. CUSHING, Cleveland, Ohio.

Councilors:

H. E. GREGORY, New Haven, Conn.

H. F. REID, Baltimore, Md.

ELECTION OF FELLOWS

Secretary Fairchild stated that the candidates for fellowship had been elected by the transmitted ballots with but few dissenting votes. The list is as follows:

ALFRED ERNEST BARLOW, B. A., M. A., D. Sc., Ottawa, Canada. Mining Geologist.

RAY SMITH BASSLET, B. A., M. S., Ph. D., Washington, D. C. Assistant Curator Stratigraphic Paleontology, U. S. National Museum.

EUGENE COSTE, B. ès-Sc., E. M., Toronto, Canada. Mining Engineer.

JOHN ALEXANDER DRESSER, B. A., M. A., Montreal, Canada. Principal of Prince Albert School.

RUSSELL D. GEORGE, A. B., A. M., Boulder, Col. Professor of Geology, University of Colorado.

ELLSWORTH HUNTINGTON, A. B., A. M., Milton, Mass. Fellow of Harvard University.

THOMAS AUGUSTUS JAGGAR, JR., A. B., A. M., Ph. D., Cambridge, Mass. Professor of Geology, Massachusetts Institute of Technology.

DOUGLAS WILSON JOHNSON, B. S., Ph. D., Cambridge, Mass. Assistant Professor of Physiography, Harvard University; Assistant Professor of Geology, Massachusetts Institute of Technology.

JOSEPH VOLNEY LEWIS, B. E., S. B., New Brunswick, N. J. Professor of Geology, Rutgers College.

DAVIS HALE NEWLAND, B. A., Albany, N. Y. Assistant State Geologist.

IDA HELEN OGILVIE, A. B., Ph. D., New York city. Tutor in Geology, Barnard College, Columbia University.

WILLIAM A. PARKS, B. A., Ph. D., Toronto, Canada. Associate Professor of Geology, University of Toronto.

WILLIAM JOHN SINCLAIR, B. S., Ph. D., Princeton, N. J. Instructor of Geology, Princeton University.

ARTHUR CLIFFORD VEATCH, Washington, D. C. Geologist, U. S. Geological Survey.

NECROLOGY

On call of the Acting President, memorials of the Fellows who had died since the Ottawa meeting were read as follows:

MEMOIR OF WILLIAM BUCK DWIGHT

BY F. J. H. MERRILL

William Buck Dwight was born in Turkey at Constantinople May 22, 1833. His father was the Reverend Harrison Gray Otis Dwight, who in 1831 settled in the Turkish capital as the first American missionary to the Armenians. His mother's name was Elizabeth Barker.

The subject of this sketch came to America in 1849 and entered Yale College, from which he was graduated with the degree of Bachelor of Arts in 1854. He then entered the Union Theological Seminary in New York and was graduated in 1856. Returning to New Haven and renewing his studies at Yale, he received the degree of Master of Arts in 1857, and two years later was granted the baccalaureate degree in Science.

This broad educational foundation was planned to support him in the undertaking which he and his elder brother had contemplated, of founding at Constantinople an unsectarian Christian institution for higher education.

This idea was, however, soon after, absorbed and developed by others, and William B. Dwight turned his energies to the field of teaching at home.

In 1859 he married Miss Eliza Howe Schneider and in the same year founded and became principal of the Englewood, New Jersey, Female Institute, in which he remained until 1865. From 1865 to 1867 he was occupied with work in mining geology in Virginia and Missouri, in the course of which he examined and reported on the now familiar Mine La Motte, and in the latter year opened a school for the children of the Military Officers at West Point, New York, which he conducted until 1870. He was then appointed Associate Principal and Professor of Natural Sciences in the Connecticut State Normal School, at New Britain. There he became actively engaged in teaching the branches to which he subsequently devoted his life, and in 1878 was called to the chair of Natural History in Vassar College, which he held until his death, in August, 1906. From 1878 to 1890 he was also in charge of the Zoological Department in the Marthas Vineyard Summer Institute, at Cottage City, and not long since was actively engaged, in cooperation with Professor N. S. Shaler, in editing the definitions in geology for the Standard Dictionary.

Shortly after taking up his residence at Poughkeepsie, Professor Dwight became interested in the geology of the surrounding district. With this began a fruitful investigation of the stratigraphy and paleontology of the limestone belts of Dutchess county, of which the age was up to that time undefined by recognized fossils, although it had been rather

generally assumed as Lower Silurian. The results of this work appeared chiefly in the *American Journal of Science* between 1879 and 1890 and established for the author a reputation as a most careful observer of obscure paleontologic detail. He was also the successful designer of mechanical appliances for making thin-sections of the much altered fossils of these semicrystalline rocks from which metamorphism had nearly obliterated the evidences of organic life. It was the privilege of the writer to become acquainted with him in 1885, when in the midst of success in the development of a new field of investigation, and his enthusiasm and self-sacrificing devotion to his work were such as to inspire the highest admiration.

At the time when Professor Dwight took up his work it was commonly considered by trustees of colleges that only chemists and physicists were entitled to laboratories and apparatus; other workers in science were supposed to elaborate their conclusions chiefly from the alcoves and dust of the college library. It was therefore at that time a more strenuous undertaking to accomplish the result in question than to do similar work in these enlightened days of more liberal allowances for college work in science. As a teacher, Professor Dwight filled a wide field and will long be remembered as a strong and beneficent influence in the progress of natural science. He was content to give much of his time to elementary instruction, but will prove to have been of broader helpfulness than some who have specialized in higher branches of geology.

His end, which came at Cottage City, Massachusetts, on August 29, 1906, was sudden, peaceful, and probably painless, perhaps the best that one can desire. Three children survive him with the comfort of an honored father's name.

MEMOIR OF SAMUEL LEWIS PENFIELD

BY JOSEPH P. IDDINGS

Samuel Lewis Penfield, known to the world of science as mineralogist and investigator, remembered by hundreds of graduates of the Sheffield Scientific School as the Professor of Mineralogy and Instructor in Blow-pipe Analysis, was to his colleagues and friends a genial and lovable companion, whose cheerfulness, generosity, steadfastness, and absolute honesty in thought and action form his most memorable characteristics.

Born in the village of Catskill-on-the-Hudson on January 16, 1856, his boyhood was spent at home, in attending school and in the enjoyment of such sports as a boy may find on the river or the neighboring mountains. His father, George H. Penfield, was a shipping merchant, prominent in business and in religious affairs, a man noted for his generosity

Samuel L. Penfield

and given to hospitality. His mother was Ann A. Cheesman, of Connecticut ancestry. By inheritance and early training he acquired that cheerful and amiable spirit which made him so companionable, and those generous and sympathetic impulses that bound him so closely to his friends. From them also sprang that native honesty which was the foundation of his character, underlying his judgment, occasioning his frankness and sincerity of speech, and forming the controlling element of his scientific work. To them must be ascribed that simple and unfaltering confidence in a beneficent Deity which accompanied him through life and permitted him to endure the strain of his prolonged illness with cheerful patience and Christian courage.

While yet a boy his fondness for study led him to hope for a college education and caused him to go to the academy at Willbraham, Massachusetts, to prepare himself for the Sheffield Scientific School, which he entered in the autumn of 1874. His tastes and talents led him to success in mathematics and the natural sciences, and at the end of his freshman year he selected the course in chemistry. In this he distinguished himself as an analyst, graduating with honors in 1877. To his classmates he was a modest, manly comrade, devoted to his studies and enthusiastic in his laboratory work.

Upon graduating he became an assistant in the chemical laboratory of the Sheffield Scientific School, and in addition to his duties as an instructor occupied himself with analytical work, especially on mineral compounds. His first publication is dated the year of his graduation and is on "The chemical composition of triphylite from Grafton, New Hampshire." His career as a mineralogist must be considered the natural result of his environment, for there was an invigorating atmosphere about the place like the clear, brilliant air that surrounds the summits of mountains. The young, enthusiastic investigator felt the inspiration that came from his association with such men as George J. Brush, James D. Dana, Edward S. Dana, and George W. Hawes, men of clear cut, vigorous ideas and brilliant attainments. It acquired specific direction through the opportunity offered him for cooperation with Professors Brush and E. S. Dana in their study of the remarkable group of minerals found in the pegmatite at Branchville, Connecticut. The analytical chemical work was intrusted to him in part, and he determined the composition of five new species of magnesian phosphates, eosphorite, triploidite, dickinsonite, fairfieldite, and fillowite, that of two other new species being determined by his classmate and associate in the chemical laboratory, Horace L. Wells. In addition to these he analyzed a number of other minerals from this locality and from elsewhere. About this time

also he contributed to the improvement of analytical processes, publishing in 1879 a paper "On a new volumetric method of determining fluorine."

It appears that up to this time his interests were centered in the purely chemical phase of his work, and it is doubtful whether he looked upon the study of mineralogy as the probable occupation of his life, for in 1880 he went to Strasburg to attend the lectures and laboratory of Professor Rudolph Fittig for the purpose of acquiring advanced methods of work in organic chemistry. He also attended some of the lectures in mineralogy given by Professor P. Groth. While in Strasburg he published, with Professor Fittig, a paper on some of the hydrocarbon compounds investigated in his laboratory.

Whatever may have been his intention at this time, his course as a mineralogist was determined upon his return to New Haven, in the autumn of 1881, by the opportunity offered him to become instructor in mineralogy in the Sheffield Scientific School in place of Doctor George W. Hawes, who had been called to Washington, D. C., to take charge of the geological department of the National Museum.

Assuming the responsibilities of this position four years after his graduation, he threw into the work of the course in determinative mineralogy and blowpipe analysis all the energy and enthusiasm of his nature, injecting into the laboratory methods the skill and precision that characterize his own analytical operations. At first his interest in mineralogy was chiefly in its chemical side, since the work with the students largely involved the chemically determinative phase of the subject, and his publications continued to treat of the chemical composition of various minerals; but an increasing interest in the physical properties of minerals and in their crystallography led him in 1884 to go to Germany for the second time, and to study with Professor Rosenbusch at Heidelberg the methods of investigation so successfully employed by him in the petrographical laboratory. This experience was followed by a broadening of his methods of work and of the range of his investigations, so that the physical characters of minerals, especially the optical properties and the crystal form, were studied, in addition to their chemical composition. In these operations he developed the same skillfulness of manipulation and thoroughness which characterize his chemical work.

In 1886 he became responsible for all of the mineralogical instruction given in the Sheffield Scientific School, owing to the fact that the school had increased to such a size as to require in its management all of the time and energy of its director, Professor Brush. In 1888 Penfield was appointed assistant professor, and in 1893 he was created a professor, and in consequence became a member of the governing board of the Sheffield Scientific School.

His success as a teacher followed from his devotion to the work and his personal interest in the individual student, and particularly from his painstaking efforts to aid the student in every way by presenting the subject in the simplest manner and by assisting the imagination by ingenious models and devices for illustrating crystal forms and structures. His success was also due to his systematic and thorough methods of work, and to a considerable extent undoubtedly to his great patience, persistency, and amiability.

That he succeeded in imparting his zeal and enthusiasm as well as his skill to those who worked in his laboratory as advanced students is shown by the successful mineralogists who have been trained by him. Evidence of his thought and ingenuity in the arrangement of his collections and apparatus is furnished by the appointments of the mineralogical laboratories in Kirtland Hall, which was built three years before his death; and it is a source of great satisfaction to his friends, as it was of happiness to himself, that the last two years of his life were spent in the enjoyment of his ideal of a mineralogical laboratory.

That he fulfilled successfully his obligations as a teacher and also achieved yet greater fame as an investigator was due to his extreme diligence as a worker, devoting to his studies all of his reserve of time and energy; and it was also due to his clear comprehension of mineralogical problems and to his ability to execute with deftness and ease the mechanical operations required in their solution. The latter showed itself in the manipulation of apparatus, whether in chemical analysis or in optical or crystallographic determinations, in which he became especially interested. He was for these reasons not only able to accomplish much in comparatively short periods of time, and thereby traverse a wide field of operation, but he obtained astonishingly good results from what might have proved inadequate material in less skillful hands. Some of his chemical analyses were carried out successfully on very small amounts of mineral, in one case upon one-tenth of a gram, and some of his crystallographic measurements were made on surprisingly small crystals. The uniformly high quality of his work, as already said, is due to his absolute honesty and conscientiousness. The exact facts and all of them that could be obtained by any means at his command were the least that would satisfy the demands of his nature, though of this exaction he was himself probably unconscious. It was an exaction applied to his work and did not extend to his general intercourse with others, toward whom he was always lenient, even indulgent.

Much of his work was done in cooperation with others, chiefly his laboratory assistants or graduate students, and he was generous in his

treatment of them and in the recognition of their work. He was also a liberal assistant of other investigators, whom he helped by advice or by contributions of analytical or of other determinative work.

In 1898 he published a revised and almost completely re-written edition of the *Manual of Determinative Mineralogy and Blowpipe Analysis*, originally the work of Professor George J. Brush. The list of publications include over eighty titles, all of which are distinct contributions to the science of mineralogy or chemistry. Their value consists in part in being determinations of new species; in part in being corrections based on more exact analyses of numerous species whose composition had been previously imperfectly and unsatisfactorily determined. Of the entirely new mineral species which were described by him, there are fourteen, namely, bixbyite, canfieldite, clinohedrite, gerhardtite, glaucocroite, graftonite, hamlinite, hancockite, leucophœnicite, nasonite, nesquehonite, pearceite, roebblingite, and spangolite. There are twenty-one species whose correct composition he was able to establish: alurgite, amblygonite, argyrodite, aurichalcite, childrenite, chondrodite, clinohumite, connellite, cookeite, ganomalite, hanksite, herderite, howlite, humite, monazite, ralsstonite, staurolite, sulphohalite, topaz, tourmaline, and turquoise.

His last investigative work, which was being carried on in association with Mr F. S. Stanley, was aimed at the elucidation of the problem of the composition of the amphiboles.

Among his specific achievements may be mentioned the recognition of the isomorphic rôle played by fluorine and hydroxyl in the structure of numerous minerals and the consequent necessity for their recognition in the composition and formulæ of such minerals; also the determination of the relations between the proportions of these constituents in a mineral series and the physical properties of the minerals, as in topaz and in the members of the humite group—humite, chondrodite, and clinohumite.

Another important achievement was the recognition of germanium in a silver ore from Bolivia and the correct crystallographic determination of the mineral argyrodite, in which the element was first discovered; also the demonstration of the control exerted upon the character of the crystal symmetry in such a chemically complex mineral as tourmaline by the acid portion of the molecule, which appears to be largely independent of the nature of the bases combined with the acid.

His contributions to crystallography consist not only of numerous accurate determinations and descriptions of crystal forms, but in the elaboration of graphical methods of representing crystals and of calculating angular relationships and also in devising simple drawing instruments and maps for use in spherical projection. His studies in these methods

led him to advocate the application of stereographic projection to geographical map making and to demonstrate the advantages to be derived from its use.

He did not confine his investigations and experiments to his laboratory, but broadened his horizon by working in the field, spending two summers in the Yellowstone National Park as an assistant on the United States Geological Survey, and other summers in New York state, North Carolina, and Colorado, as well as in various parts of Europe, visiting at the same time the most celebrated mineral collections and becoming acquainted with foreign mineralogists.

The high quality of his work and the value of his contributions to the science of mineralogy are attested by the esteem in which he was held both abroad and in his own country, as well as by the honors bestowed upon him. Beginning with the year 1893, these rapidly succeeded one another in the following order: Associate Fellow of the American Academy of Arts and Sciences in Boston; in 1896, Foreign Correspondent of the Geological Society of London, Master of Arts in Yale University; in 1900, Member of the National Academy of Sciences; in 1902, Fellow of the American Association for the Advancement of Science, Corresponding Member of the Royal Society of Sciences at Göttingen, Member of the Scientific Society of Christiania; in 1903, Corresponding Member of the Geological Society of Stockholm and Foreign Member of the Mineralogical Society of Great Britain. In 1904 he received the honorary degree of Doctor of Laws from the University of Wisconsin. He was a member of the Connecticut Academy of Sciences and a fellow of the Geological Society of America.

These honors he accepted with such charming modesty that many of them remained hidden and unsuspected by even those most intimate with him. His achievements followed as the natural results of his ceaseless activities, which sprang from a love of research for its own sake.

For twenty years after his graduation he remained unmarried, devoting his time, thought, and energy to his scientific work. During the greater part of these years he lived with a small company of bachelor colleagues in the attic of Sheffield Hall, familiarly known as "Old Sheff"; here his genial and generous nature won him the admiration and affection of his associates and friends. In these homely halls, low-roofed and bare, there grew up a group of men who seem to have acquired from their surroundings certain sturdy qualities which are to be reckoned among their fortunate characteristics.

In 1897 Professor Penfield married Miss Grace Chapman, of Albany, who brought to him great happiness, and whose devotion to him through

the years of his illness strengthened his courage and undoubtedly prolonged his life.

As the man is more than his achievements, so the character of Samuel Lewis Penfield must be accorded higher praise than his labors. Honest, and modest, and true; sympathetic, and generous, and loving; steadfast and simple in his affections and in his beliefs, his beneficent nature has affected the lives of many who have come within its influence, and, being transmitted beyond the range of his own experience, will contribute its share to the sum of human goodness. And, as his works and writings will be found in print long after his personality shall have faded from the minds of men, may this suggestion of his noble character be found in place beside the record of his accomplishments.

BIBLIOGRAPHY *

1877. On the chemical composition of triphylite from Grafton, New Hampshire. *American Journal of Science* (3), xlii, 425-427.
1878. Analyses of eosphorite, triploidite, and dickinsonite (by George J. Brush and Edward S. Dana). *Ibid.*, xvi, 40, 45, 117.
1879. On the chemical composition of triphylite. *Ibid.*, xvii, 226-229.
Analyses of fairfieldite and fillowite (by George J. Brush and Edward S. Dana). *Ibid.*, xvii, 362, 365.
Analyses of chabazite and rhodochrosite (by George J. Brush and Edward S. Dana). *Ibid.*, xviii, 50.
On a new volumetric method of determining fluorine. *Amer. Chem. Jour.*, i, 27-29.
On the chemical composition of amblygonite. *American Journal of Science* (3), xviii, 295-301.
1880. On the chemical composition of childrenite. *Ibid.*, xix, 315-316.
Analyses of some apatites containing manganese. *Ibid.*, 367-369.
Analyses of spodumene, β -spodumene, eucryptite, cymatalite, muscovite, microcline, and killinite (by George J. Brush and Edward S. Dana). *Ibid.*, xx, 259-263, 268, 271-274.
1881. Analysis of jarosite from the Vulture mine, Arizona. *Ibid.*, xxi, 160.
1882. Occurrence and composition of some American varieties of monazite. *Ibid.*, xxiv, 250-254.
Ueber die Phenylhomoparaconsäure (with R. Fittig). *Ann. der Chem.*, ccxvi, 119-127.
1883. Scovillite: a new phosphate of didymium, yttrium, and other rare earths from Salisbury, Connecticut (with G. J. Brush). *American Journal of Science* (3), xxv, 459-463.
Analyses of the two varieties of lithiophilite. *Ibid.*, xxvi, 176.
On a variety of descloizite from Mexico. *Ibid.*, 361-365.
1884. Identity of scovillite with rhabdophane (with G. J. Brush). *Ibid.*, xxvii, 200-201.

* This list has been taken from the memoir published by Professor L. V. Pirsson in the *American Journal of Science*, fourth series, vol. xxii, 1906. It has been revised by Professor H. L. Wells, and is believed to be practically complete.

- On the occurrence of alkalies in beryl. *Ibid.*, xxviii, 25-32.
- Ueber Erwärmungsversuche an Leucit und anderen Mineralien. *Neues Jahrb. für Min.*, ii, 224.
1885. Crystallized tiemannite and metacinnabarite. *American Journal of Science* (3), xxix, 449-454.
- Gerhardtite and artificial basic cupric nitrates (with H. L. Wells). *Ibid.*, xxx, 50-57.
- Crystals of analcite from the Phoenix mine, Lake Superior copper region. *Ibid.*, 112-113.
- Mineralogical notes (with E. S. Dana). *Ibid.*, 136-139.
1886. Brookite from Magnet Cove, Arkansas. *Ibid.*, xxxi, 387-389.
- Chemical composition of herderite and beryl (with D. N. Harper). *Ibid.*, xxxii, 107-117.
- On hitherto undescribed meteoric stones (with E. S. Dana). *Ibid.*, 226-231.
- On pseudomorphs of garnet from Lake Superior and Salida, Colorado (with F. L. Sperry). *Ibid.*, 307-311.
- On the chemical composition of ralstonite (with D. N. Harper). *Ibid.*, 380-385.
- Crystallized vanadinite from Arizona and New Mexico. *Ibid.*, 441-443.
1887. Phenacite from Colorado. *Ibid.*, xxxiii, 130-134.
- On the chemical composition of howlite (with E. S. Sperry). *Ibid.*, xxxiv, 220-223.
- Triclinic feldspars with twinning striations on the brachypinacoid (with F. L. Sperry). *Ibid.*, 390-393.
1888. On the crystalline form of polianite (with E. S. Dana). *Ibid.*, xxxv, 243-247.
- Bertrandite from mount Antero, Colorado. *Ibid.*, xxxvi, 52-55.
- Mineralogical notes (with E. S. Sperry). *Ibid.*, 317-331.
1889. On the crystalline form of sperrylite. *Ibid.*, xxxvii, 71-73.
- On some curiously developed pyrite crystals from French creek, Chester county, Pennsylvania. *Ibid.*, 209-212.
- Crystallized bertrandite from Stoneham, Maine, and mount Antero, Colorado. *Ibid.*, 213-216.
- Results obtained by etching a sphere and crystals of quartz with hydro-fluoric acid (with Otto Meyer). *Trans. Conn. Acad.*, viii, 158-165.
1890. On lansfordite, nesquehonite, a new mineral, and pseudomorphs of nesquehonite after lansfordite (with F. A. Genth). *American Journal of Science* (3), xxxix, 121-137.
- On spangolite, a new copper mineral. *Ibid.*, 370-378.
- On hamlinite, a new rhombohedral mineral from the herderite locality at Stoneham, Maine (with W. E. Hidden). *Ibid.*, 511-513.
- Fayalite in the obsidian of Lipari (with J. P. Iddings). *Ibid.*, xl, 75-78.
- On connellite from Cornwall, England. *Ibid.*, 82-86.
- Crystallographic notes (with F. A. Genth). *Ibid.*, 199-207.
- Chalcopyrite crystals from French creek, Pennsylvania. *Ibid.*, 207-211.
- Anthophyllite from Franklin, Macon county, North Carolina. *Ibid.*, 394-397.
- On the beryllium minerals from mount Antero, Colorado. *Ibid.*, 488-491.

1891. Chemical composition of aurichalcite. *Ibid.*, xli, 106-109.
 Crystallographic notes (with F. A. Genth). *Ibid.*, 394-400.
 The minerals in hollow sperulites of rhyolite from Glade creek, Wyoming (with J. P. Iddings). *Ibid.*, xlii, 39-46.
1892. On caesium trihalides (by H. L. Wells) and their crystallography. *Ibid.*, xliii, 17-32.
 Crystallographic notes (with F. A. Genth). *Ibid.*, 184-189.
 Crystallography of the rubidium and potassium trihalides. *Ibid.*, 475-487.
 On polybasite and tennantite from Mollie Gibson mine, Aspen, Colorado (with S. H. Pearce). *Ibid.*, xliv, 15-18.
 Crystallography of the alkali-metal pentahalides. *Ibid.*, 42-49.
 On herderite from Hebron, Maine (with H. L. Wells). *Ibid.*, 114-116.
 Crystalline form of RbCl.HIO_3 and CsCl.HIO_3 . *Ibid.*, 132-133.
 Crystallography of double halides of silver and alkali-metals. *Ibid.*, 155-157.
 Crystallography of caesium and rubidium chloraurates and bromaurates. *Ibid.*, 157-162.
 Crystallography of the caesium-mercuric halides. *Ibid.*, 311-321.
 Crystallographic notes (with F. A. Genth). *Ibid.*, 381-389.
1893. On cookeite from Paris and Hebron, Maine. *Ibid.*, xlv, 393-396.
 Mineralogical notes (zunyite, xenotime). *Ibid.*, 396-399.
 On pentlandite from Sudbury, Ontario. *Ibid.*, 493-497.
 On canfieldite, a new germanium mineral, and on the chemical composition of argyrodite. *Ibid.*, xlv, 107-113.
 Minerals from the Manganese mines of Saint Marcel, Piedmont. *Ibid.*, 288-295.
1894. Chemical composition of staurolite and on its inclusions (with J. H. Pratt). *Ibid.*, xlvii, 81-89.
 Chemical composition of chondrodite, humite, and clinohumite (with W. T. H. Howe). *Ibid.*, 188-206.
 Crystallization of willemite. *Ibid.*, 305-309.
 Crystallization of herderite. *Ibid.*, 329-339.
 Chemical composition and related physical properties of topaz (with J. C. Minor). *Ibid.*, 387-396.
 On argyrodite and a new sulphostannate of silver (canfieldite) from Bolivia. *Ibid.*, 451-454.
 On thallium triiodide and its relation to the alkali-metal triiodides (with H. L. Wells). *Ibid.*, 463-466.
 On some methods for the determination of water. *Ibid.*, xlviii, 30-37.
 Mineralogical notes. *Ibid.*, 114-118.
 Mineralogical notes and separation of minerals of high specific gravity (with D. A. Kreider). *Ibid.*, 141-144.
1895. Note on the crystallization of calaverite. *Ibid.*, i, 128-131.
 Effect of the mutual replacement of manganese and iron on the optical properties of lithiophilite and triphylite (with J. H. Pratt). *Ibid.*, 387-390.
 Devices for the separation of minerals of high specific gravity. *Ibid.*, 446-448.

1896. Fayalite from Rockport, Massachusetts, and the optical properties of the chrysolite group (with E. H. Forbes). *Ibid.* (4), i, 129-135.
Occurrence of thaumasite at West Paterson, New Jersey (with J. H. Pratt). *Ibid.*, 229-233.
On pearceite, a sulpharsenite of silver, and on the crystallization of polybasite. *Ibid.*, ii, 17-29.
1897. On roeblingite, a new silicate from Franklin Furnace, New Jersey, containing SO₂ and lead (with H. W. Foote). *Ibid.*, iii, 413-415.
Identity of chalcostibite (wolfsbergite) and guejarite and on chalcostibite from Huanchaca, Bolivia (with A. Frenzel). *Ibid.*, iv, 27-35.
On bixbyite, a new mineral, and on the associated topaz (with H. W. Foote). *Ibid.*, 105-108.
Note on the composition of ilmenite (with H. W. Foote). *Ibid.*, 108-110.
Chemical composition of hamlinite and its occurrence with bertrandite at Oxford county, Maine. *Ibid.*, 313-316.
1898. On clinohedrite, a new mineral from Franklin, New Jersey (with H. W. Foote). *Ibid.*, v, 289-293.
Crystallographic note on kreunerite from Cripple Creek, Colorado. *Ibid.*, 375-377.
Note on sperrylite from North Carolina (with W. E. Hidden). *Ibid.*, vi, 381-383.
Manual of determinative mineralogy and blowpipe analysis by G. J. Brush. Revised and enlarged, with new tables. 312 pp. New York: John Wiley & Sons.
1899. On the chemical composition of tourmaline (with H. W. Foote). *American Journal of Science* (4), vii, 97-125.
On the chemical composition of parisite and a new occurrence at Ravalli county, Montana (with C. H. Warren). *Ibid.*, viii, 21-24.
On some new minerals from the zinc mines at Franklin, New Jersey (hancockite, glaucocroite, nasonite, leucophœnicite, and note on chemical composition of ganomalite (with C. H. Warren). *Ibid.*, 339-353.
1900. On graftonite, a new mineral from Grafton, New Hampshire, and its intergrowth with triphylite. *Ibid.*, ix, 20-32.
Siliceous calcites from the Bad Lands, Washington county, South Dakota (with W. E. Ford). *Ibid.*, 352-354.
Chemical composition of sulphohalite. *Ibid.*, 425-428.
The interpretation of mineral analyses, a criticism of recent articles on the constitution of tourmaline. *Ibid.*, x, 19-32.
On some interesting developments of calcite crystals (with W. E. Ford). *Ibid.*, 237-244.
1900. Contactgoniometer und transporteur von Einfacher Construction. *Zeitschr. für Kryst.*, xxxiii, 548-554.
On the chemical composition of turquois. *American Journal of Science* (4), x, 346-350.
1901. The stereographic projection and its possibilities from a graphical standpoint. *Ibid.*, i, 1-24.
Contributions to Mineralogy and Petrography. Yale Bicentennial Pub-

- lication (with L. V. Pirsson). 482 pp. Charles Scribner's Sons. New York.
- On calaverite (with W. E. Ford). *Ibid.*, xii, 225-246.
1902. New occurrence of sperrylite (with H. L. Wells). *Ibid.*, xiii, 95-96.
- Use of the stereographic projection for geographical maps and sailing charts. *Ibid.*, xiii, 245-275, 347-376.
- On the solution of problems in crystallography by means of graphical methods based on spherical and plane trigonometry. *Ibid.*, xiii, 249-284.
- Some additions to the alunite-jarosite group of minerals (with W. F. Hillebrand). *Ibid.*, xiv, 211-220.
1903. Tables of minerals, including the uses of minerals and statistics of the domestic production. 8°, 77 pp. (New Haven, Connecticut).
1905. On crystal drawing. *American Journal of Science*, xix, 39-75.
- On tychite, a new mineral from Borax lake, California, and on its artificial production and its relations to northupite (with G. S. Jamieson). *Ibid.*, xx, 217-224.
1906. On the drawing of crystals from stereographic and gnomonic projections. *Ibid.*, xxi, 206-215.
- Filter tubes for collection of precipitates on asbestos (with W. M. Bradley). *Ibid.*, 453-466.
- On stibiotantalite (with W. E. Ford). *Ibid.*, xxii, 61-77.
1907. On the chemical composition of amphibole (with F. C. Stanley). *Ibid.*, xxiii, 23-51.

MEMOIR OF ISRAEL C. RUSSELL

BY BAILEY WILLIS

A native of New York by birth, Israel C. Russell was at home everywhere, through the intrepidity of his spirit and the breadth of his interests. The arid plains of our West, or the snowy scarps of Saint Elias, or the ash-strewn slopes of Pelée were to him equally alluring. The unexplored challenged the inborn instinct of the explorer, which dominated the aims of his life; he delighted to attempt the unknown; he was independent and at ease in the wildest environment; and where in the interest of science difficulty or danger were to be met, there he was most resourceful, cool, and happy. On the other hand, he was restless under routine; he chafed in harness; he rather endured than enjoyed the comforts of conventional life, with the obligations they imposed.

It is told of a colonial Russell ancestor that when calling Sunday evenings on the lady whom he afterward married he would confine his remarks to a greeting and a phrase of thanks on leaving for the pleasant visit he had enjoyed. Silence ran in the family, but so also did a genial sense of the pleasure of conversation. Outwardly the Russell we knew was a silent, self-contained man, from whom flashed occasionally a sally of keen wit; but the external reserve covered a fertile mind and a warm kindly

Isaac C. Russell

nature. In the exchange of scientific thought he gave generously; in the field he was a considerate and loyal comrade, as men near and true to Nature are; and in his home, in those intimate relations that are the highest test of character, he has left a memory which sustains his family and is a heritage of love and inspiration for his children.

Russell's career falls into four epochs, through all of which runs the chain of exploration. After graduating from the University of New York in 1872, he sought an opportunity to accompany the U. S. Transit of Venus expedition to New Zealand in 1874 and 1875, and finding the opportunity limited to the position of photographer, he learned the then difficult art and received the appointment. This journey around the world opened to him the great field of physical geography in which he afterward labored continuously. He drifted for five years after his return. Two years as assistant professor of geology at Columbia, a season in New Mexico, a journey to Europe—these completed the first epoch of restless preparation. When I was a student in Columbia he was constantly at work among the collections—a slender, silent figure, a typical student, in whom one would not suspect the power of initiative, the capacity for daring and doing, which characterized the man.

The second epoch covered the years from 1880 to 1885, during which, as geologist of the U. S. Geological Survey, in association with Gilbert, he studied the Pleistocene history of the Great basin. Here he was in his element—free, face to face with Nature, and in frequent intercourse with one who may be characterized as one of the wisest organizers of American geologic science of any period. While Gilbert interpreted the history of lake Bonneville, Russell developed that of lake Lahontan; the two studies ran parallel and reacted each upon the other. The senior's influence is seen in the character of the Lahontan monograph, which is a systematic, complete, and essentially final work, peculiar among Russell's writings in being thus highly organized and finished; but the acute observations regarding the chemical history of the waters and its bearing on variations of the lake are Russell's independent contribution.

When in 1885 Gilbert surrendered his promising investigations of the Great basin to undertake the study of the Appalachian province, Russell was also transferred and their association continued. Cogent as the administrative reasons for the transfer were, the change of tasks was unfortunate for the scientists and for science. Gilbert was burdened with executive affairs, and Russell found in the southern Appalachians an uncongenial field. Nevertheless he worked with characteristic activity and completed the investigation of the section assigned him in Gilbert's systematic plan for studying the Appalachians; but he did not bring the

work to a condition for publication and it was inherited by Hayes. Several of the folios are still unpublished on account of the difficulty of working out the metamorphic rocks.

Russell's earliest geologic studies had been on the Triassic rocks of New Jersey and Connecticut, and they were continued on his return to the East in North Carolina and Virginia. When Powell planned the correlation papers he was the best equipped man available for the Triassic. He extended the fieldwork in 1888-1889, visiting the sections in the far West, but was interrupted by an assignment to exploration in Alaska. His observations beyond the eastern districts being thus left incomplete, the correlation essay was confined to the Newark formation. It is a carefully prepared abstract of the literature, accompanied by original chapters on the stratigraphy, paleontology, diastrophism, and degradation of the formation. Later studies have contributed additional details of stratigraphy and structure of the Newark, but they have not modified Russell's conclusions in regard to its former distribution and the geographic conditions of the epoch.

A logical development of his studies of western Pleistocene was the investigation of subaerial decay of rocks, and the Triassic invited his attention directly to the cause of the red color in certain deposits.

Thus closed the third epoch of his career, during which he worked against his inclination and reaped but little result from nearly four years of industrious application to the tasks assigned him.

Alaska was a field which was the antithesis of the Appalachian for a man of Russell's abilities. He first visited it in 1889, as the representative of the U. S. Geological Survey on the expedition of the U. S. Coast and Geodetic Survey to determine the boundary with British Columbia. After many weeks of steamer travel through the Yukon flats, with little chance for geology, he cut loose from the expedition on the upper Yukon and with three miners came out across the ranges to the coast. Alert to observe the features of stratigraphic and surface geology, he made the most of the somewhat limited opportunity and contributed valuable observations on the general structure of the Yukon basin and the limitations of Pleistocene glaciation. His article on the surface geology of Alaska lays down for that subject the principal lines of investigation which have since been elaborated by others.

Mount Saint Elias, then supposed to be the highest peak in United States territory, had for several years been the pole of Russell's purpose. His imagination was fascinated by the snowy heights that rise in alpine grandeur near the sea, yet remained untrodden. In contrast with this inaccessible pinnacle of unknown geologic interest, the rich fields of

geology within easy reach had no attraction for him. Better an hour on that frost-fettered virgin height than a year in the haunts of men. Through the cooperation of the National Geographic Society and the U. S. Geological Survey, his purpose of exploration was realized and he twice visited the region. He did not attain the highest summit, being driven back repeatedly by storms, and the last time when all the difficulties of the ascent had been overcome; but he yielded only to irresistible conditions and made a record of indomitable pluck and perseverance. The experience which he gathered at cost of extreme effort and risk he placed freely at the service of others, and a share in the honor of the successful ascent made by the Duke of the Abruzzi is justly granted by the latter to Russell.

In the course of his second expedition Russell was separated from his companions on the slopes of mount Saint Elias and proceeded alone to a camp they had established above the ice cascades of the glaciers beneath the cliffs that define the peak. The others were to join him there, but a heavy storm interfered and Russell was isolated among the wildest scenes of alpine crags and ice. His tent was crushed and he burrowed in the snow, where he measured out his scant supplies into eight days' rations and prepared himself to await the issue. Afterward, in discussing Chamberlin's suggestion that heat waves might be transmitted through glacial ice, he once remarked that none penetrated the snowbank in which he passed three days. Storm winds whirled through the amphitheater; avalanches shot from the cliffs; the isolation was utter and to many men would have been terrible; but Russell's was a spirit that rose with the battling elements and exulted in challenging Nature's most savage mood.

The expeditions to Saint Elias were made across the Malaspina, the great confluent glacier which receives the alpine ice-streams of the range and extends to the coast, where it is in part covered by primeval forest. The opportunity to study glacial phenomena was unrivaled and was fully exploited by Russell, who thus by direct observation made us acquainted with the piedmont type of ice-sheet, which was formerly of common occurrence throughout the northern hemisphere.

The exploration of mount Saint Elias was the most striking, but not the most fruitful of Russell's campaigns against the unknown. He continued his sallies into unexplored regions during the years following 1901, going out each summer, as the vacation season permitted. He repeatedly made reconnaissances in Washington, Oregon, and Idaho for the U. S. Geological Survey, and there are extensive areas which we know only through his publications.

In 1892 Russell succeeded Alexander Winchell in the chair of Geology

at the University of Michigan and established his home at Ann Arbor. Five years previously (1887) he had married Julia Augusta Olmstead, who, with their three daughters and one son, survives him. The home circle was one in which it was a happy privilege to be welcomed, and in that atmosphere Russell, talking of glowing scenes of the deserts and snow peaks he knew so intimately, revealed beneath the quiet exterior the enthusiasm of an artist.

Russell's place in science was determined by the epoch in which he lived and by the strong impulses of his own nature. The conditions of the one and the tendencies of the other were in harmony when he began his career; later the conditions changed, but Russell did not, except in outward compliance, and he remained ever a frontiersman of science.

In the eighteen seventies American geography and geology were still sciences of exploration; the refinements of surveying were undeveloped; the specialization of the general subject had scarcely begun. The Fortieth Parallel Survey, which is now regarded as high grade reconnaissance work, was the standard piece of geologic surveying. Geology was a relatively simple study, which offered vast opportunities to keen observers and explorers, but made relatively slight demands upon the systematist and the specialist. The opportunities were those which Russell was most able to exploit brilliantly; the demands he could afford to overlook; and throughout his career he always sought the fresh opportunity, he did his best work in virgin fields. He takes his place in the group of geographers and geologists, of whom it suffices to name Newberry, Hayden, Powell, and Richthofen—men who did pioneer work, which is the essential foundation of later research; men who have blazed out the way, both physically and intellectually, for the road-builders who follow; men to whom all honor is due for the courage and high purpose they had in the face of difficulties, and to whom we owe a debt of gratitude for the greater opportunities they have opened to us.

BIBLIOGRAPHY

- New Zealand flax. *American Naturalist*, vol. 10, 1876, pp. 18-21.
Lake Wakatipu. New Zealand. *American Naturalist*, vol. 10, 1876, pp. 385-392.
Notes on the ancient glaciers of New Zealand. New York Lyceum Natural History, Annals, vol. 11, 1876, pp. 251-265, map 19.
On the formation of lakes. *Popular Science Monthly*, vol. 9, 1876, pp. 539-546.
The giant birds of New Zealand. *American Naturalist*, vol. 11, no. 1, 1877, pp. 11-21.
Concerning footprints. *American Naturalist*, vol. 11, 1877, pp. 406-417.

- On the intrusive nature of the Triassic trap sheets of New Jersey. *American Journal of Science*, 3d series, vol. 15, 1878, pp. 277-280.
- On the occurrence of a solid hydrocarbon in the eruptive rocks of New Jersey. *American Journal of Science*, 3d series, vol. 16, 1878, pp. 112-114. Abstract, *American Naturalist*, vol. 13, 1879, pp. 198-199 (2/3 p.).
- On the physical history of the Triassic formation in New Jersey and the Connecticut valley. New York Academy of Science, Annals, vol. 1, 1879, pp. 220-254. Reviewed by J. D. Dana, *American Journal of Science*, 3d series, vol. 17, 1879, pp. 328-330, and P. Frazer, *American Naturalist*, vol. 13, 1879, pp. 284-292.
- A new form of compass-clinometer. New York Academy of Sciences, Annals, vol. 1, 1879, pp. 263-264.
- A sketch of New Zealand, with pen and pencil. *American Naturalist*, vol. 13, 1879, pp. 65-77.
- The geological museum of the School of Mines, Columbia College. *American Naturalist*, vol. 13, 1879, pp. 502-513.
- The fertilization of the Wistaria. *American Naturalist*, vol. 13, 1879, pp. 648-649.
- On the former extent of the Triassic formation of the Atlantic states. *American Naturalist*, vol. 14, 1880, pp. 703-712.
- Sulphur deposits in Utah and Nevada. New York Academy of Science, Trans., vol. 1, 1882, pp. 168-175. *Engineering and Mining Journal*, vol. 35, 1883, pp. 31-32. Abstract, *American Journal of Science*, 3d series, 1883, vol. 25, p. 158 (1/3 p.).
- Sketch of the geological history of lake Lahontan, a Quaternary lake of north-western Nevada. U. S. Geological Survey, J. W. Powell, Director; 3d Annual Report, for 1881-1882, pp. 189-235. Washington, 1883. Abstracts, *American Journal of Science*, 3d series, 1884, vol. 27, pp. 67-68; *American Naturalist*, vol. 19, 1885, pp. 152-153, plates; *Scientific American Supplement*, vol. 18 (no. 450), 1884, p. 7187; *Science*, vol. 4, 1884, pp. 64-66.
- The geology of Hudson county, New Jersey. New York Academy of Science, Annals, vol. 2, 1883, pp. 27-80, pl. 2. In part *Science* (edited by J. Michels), vol. 2, 1881, pp. 63-65.
- Playas and playa lakes. *Popular Science Monthly*, vol. 22, 1883, pp. 380-385. Abstract, *Science*, vol. 1, 1883, pp. 77-78 (1/4 col.).
- A geological reconnaissance in southern Oregon. U. S. Geological Survey, J. W. Powell, Director; 4th Annual Report, 1882-1883, pp. 431-464, plates. Washington, 1884. Abstract, *Science*, vol. 6, 1885, pp. 58-59 (1/2 p.).
- Lakes of the Great Basin. *Science*, vol. 3, 1884, pp. 322-323.
- Deposits of volcanic dust in the Great Basin. [Abstract.] Bulletin Washington Philosophical Society, vol. 7, 1885, pp. 18-20. Abstract, *Science*, vol. 3, 1884, p. 555 (1/4 p.).
- North Carolina coal fields. [Review of report of H. M. Chance.] *Science*, vol. 6, 1885, pp. 548-549.
- The existing glaciers on the high sierra of California. [Abstract.] Bulletin Washington Philosophical Society, vol. 7, 1885, pp. 5-8.
- Existing glaciers of the United States. U. S. Geological Survey, J. W. Powell, Director; 5th Annual Report, 1883-1884, pp. 303-355, 1885.

- What is a glacier? Abstract, Philosophical Society of Washington, Bulletin, vol. 7, 1885, p. 37.
- Geological history of lake Lahontan, a Quaternary lake of northwestern Nevada. U. S. Geological Survey, J. W. Powell, Director; monograph XI, 288 pages, 44 plates, 4°. Washington, 1885 [1886]. Abstract by R. McLintoch. Liverpool Literary and Philosophical Society, Proceedings, vol. 41, 1890, pp. 339-342. Abstracts, *Science*, vol. 10, 1887, pp. 79-80. *Scottish Geographical Magazine*, vol. 3, 1887, pp. 466-472.
- Natural gas and coal in Chesterfield county, Virginia. *The Richmond Dispatch*, February 20, 1887.
- Notes on the faults of the Great basin and of the eastern base of the Sierra Nevada. Bulletin, Washington Philosophical Society, vol. 9, 1887, pp. 5-7. Abstract, *Neues Jahrbuch*, Bd. 2, 1887, pp. 317-318.
- The Jordan-Arabah depression and the Dead sea. *Geological Magazine*, dec. iii, vol. 5, 1888, pp. 337-344, 387-395.
- The Great Basin. *Overland Monthly*, second series, vol. 11, April, 1888, pp. 420-426.
- Quaternary history of Mono Valley, California. U. S. Geological Survey. Stth Report, J. W. Powell, 1889, pp. 261-394, plates 16-44. Abstracts, *American Geologist*, vol. 6, pp. 54-56; *American Journal of Science*, 3d series, vol. 39, 1889, p. 402 ($\frac{1}{2}$ p.).
- Subaerial deposits of the arid regions of North America. *Geological Magazine*, Decade III, vol. 6, 1889, pp. 289-295, 342-350.
- The Newark system. *American Geologist*, vol. 3, 1889, pp. 178-182. Reviewed by C. H. Hitchcock, *American Geologist*, vol. 5, 1890, pp. 200-202.
- Subaerial decay of rocks and origin of the red color of certain formations. Bulletin no. 52, U. S. Geological Survey, 65 pp., plates 1-5. Abstracts, *American Geologist*, vol. 5, 1890, pp. 110-111; *Canadian Record of Science*, vol. 4, 1889, pp. 74-75; *Engineering and Mining Journal*, vol. 49, 1889, pp. 307-308 ($\frac{2}{3}$ col.); *Popular Science Monthly*, vol. 36, 1889, p. 567 (1/5 col.). Reviewed by J. D. Dana, *American Journal of Science*, 3d series, vol. 39, 1889, pp. 317-319.
- Notes on the surface geology of Alaska. Bulletin, Geological Society of America, vol. 1, 1890, pp. 99-162, pl. 2. Discussed by N. S. Shaler and T. C. Chamberlin, pp. 155-156. Abstracts, *American Geologist*, vol. 5, pp. 118-119 (1/5 p.); *American Naturalist*, vol. 24, 1890, p. 208 (4 lines).
- Ice cliffs on Kowah river, Alaska, observed by Lieutenant Cantwell. *American Geologist*, vol. 6, 1890, pp. 49-50. Presenting a letter from J. C. Cantwell.
- An expedition to mount Saint Elias, Alaska. *National Geographic Magazine*, vol. 3, 1891, pp. 53-194, pls. 2-20. Abstracts, *American Journal of Science*, 3d series, vol. 42, 1891, pp. 171-172 ($\frac{2}{3}$ p.); *American Geologist*, vol. 8, 1891, p. 120.
- Two expeditions to mount Saint Elias. Part 2. *The Century Magazine*, vol. 41, April, 1891, pp. 872-884.
- Explorations in Alaska. *American Geologist*, vol. 7, 1891, pp. 33-38.
- Are there glacial records in the Newark system? *American Journal of Science*, 3d series, vol. 41, 1891, pp. 499-505. Abstract, *American Naturalist*, vol. 25, 1891, p. 739 (5 lines).
- Has "Newark" priority as a group name? *American Geologist*, vol. 7, 1891, pp. 238-241.

- Origin of the gravel deposits beneath Muir glacier, Alaska. *American Geologist*, vol. 9, 1892, pp. 190-197.
- Climatic changes indicated by the glaciers of North America. *American Geologist*, vol. 9, 1892, pp. 322-336.
- Mount Saint Elias and its glaciers. *American Journal of Science*, 3d series, vol. 43, 1892, pp. 169-182. Abstract, *American Geologist*, vol. 9, 1892, pp. 340-341.
- Height and position of mount Saint Elias. *National Geographic Magazine*, vol. 3, 1892, pp. 231-237.
- Mount Saint Elias revisited. *The Century Magazine*, vol. 44, June, 1892, pp. 190-203.
- Correlation papers. The Newark system. Bulletin no. 85, U. S. Geological Survey, 1892. 344 pp., pls. i-xiii, figs. 1-4. Abstracts, *Journal of Geology*, vol. i, pp. 740-744; *American Geologist*, vol. 12, pp. 402 ($\frac{1}{3}$ p.); *American Naturalist*, vol. 27, 1893, pp. 987-988.
- Malaspina glacier. *Journal of Geology*, vol. 1, 1893, pp. 219-245. Abstract, *American Geologist*, vol. 12, pp. 121-122 ($\frac{2}{3}$ p.).
- Geological history of the Laurentian basin. *Journal of Geology*, vol. 1, 1893, pp. 394-408.
- Second expedition to mount Saint Elias in 1891. U. S. Geological Survey, 13th Annual Report, 1893, pt. 2, pp. 7-91, pls. iii-xxi, figs. 1-6.
- A geological reconnaissance in central Washington. Bulletin U. S. Geological Survey, no. 108, 1893, pp. 1-108, pls. i-xii, figs. 1-8.
- Alaska: its physical geography. *The Scottish Geographical Magazine*, vol. 10, no. 8, August, 1894, pp. 393-413.
- Present and extinct lakes of Nevada. National Geographic Monographs, vol. 1, no. 4, 1895, pp. 101-132, 4 figs., 3 maps.
- Review of "Report on surface geology," by R. D. Salisbury. *Journal of Geology*, vol. 3, 1895, pp. 358-364.
- Lakes of North America, a reading lesson for students of geography and geology. Boston, 1895, 125 pp., 23 pls.
- The influence of debris on the flow of glaciers. *Journal of Geology*, vol. 3, 1895, pp. 823-832.
- Reports of a conference on geography. American Geographical Society, Bulletin, vol. 27, no. 1, 1895, pp. 30-41.
- A journey up the Yukon River. American Geographical Society, Bulletin, vol. 27, no. 2, 1895, pp. 143-160.
- The Newark system. *Science*, new series, vol. 1, 1895, pp. 266-268.
- Igneous intrusions in the neighborhood of the Black hills of Dakota. *Journal of Geology*, vol. 4, 1896, pp. 23-43, pls. i-iii.
- On the nature of igneous intrusions. *Journal of Geology*, vol. 4, 1896, pp. 177-194.
- Mountaineering in Alaska. American Geographical Society, Bulletin, vol. 28, no. 3, 1896, pp. 217-228.
- Igneous intrusions and volcanoes. *Popular Science Monthly*, vol. 50, 1896, pp. 240-250.
- A reconnaissance in southeastern Washington. U. S. Geological Survey, Water Supply and Irrigation Papers no. 4, 1897, 96 pp. pls. i-vii, figs. 1-3.

- Principal features of the geology of southeastern Washington. *Mining*, vol. 3, 1897, pp. 163-165.
- "Plasticity" of glacial ice. *American Journal of Science*, 4th series, vol. 3, 1897, pp. 344-346. Abstract, *Journal of Geology*, vol. 5, 1897, pp. 104-105.
- Review of "Glacier bay and its glaciers," by H. F. Reid. *Journal of Geology*, vol. 5, 1897, pp. 203-206.
- Glaciers of North America, a reading lesson for students of geography and geology. Boston and London, 1897, 210 pp., 22 pls. *Journal of Geology*, vol. 5, 1897, pp. 302-303, review by T. C. Chamberlin; *National Geographic Magazine*, vol. 8, 1897, pp. 124-125, review by W. J. McGee; *Science*, new series, vol. 5, 1897, pp. 660-661, review by H. F. Reid; *American Geologist*, vol. 19, 1897, p. 278 ($\frac{1}{4}$ p.), review by N. H. Winchell.
- Volcanoes of North America, a reading lesson for students of geography and geology. New York, 1897, xiv, 346 pp., 16 pls. Review, *Nature*, vol. 57, 1897, pp. 70-71.
- A new geographical magazine. *Science*, new series, vol. 5, 1897, pp. 477-478.
- Impressions of mount Rainier. *Scribner's Magazine*, vol. 22, no. 2, August, 1897, pp. 169-176.
- Principal features of the geology of southeastern Washington. Abstract, *Journal of Geology*, vol. 5, 1897, pp. 107-109.
- The wheat-lands of Washington. *Harper's Weekly*, vol. 41, 1897, pp. 735-740.
- The future meetings of the Geological Society of America. (An editorial.) *Journal of Geology*, vol. 5, 1897, pp. 194-195.
- Glaciers of mount Rainier, with a paper on the rocks of mount Rainier, by George Otis Smith. U. S. Geological Survey, 18th Annual Report, pt. 2, 1898, pp. 355-423, pls. lxxv-lxxxii.
- The great terrace of the Columbia and other topographic features in the neighborhood of lake Chelan, Washington. *American Geologist*, vol. 22, 1898, pp. 362-369.
- Topographic features due to landslides. *Popular Science Monthly*, vol. 53, 1898, pp. 480-489, 3 figs.
- Geography of the Laurentian basin. Bulletin American Geographic Society, vol. 30, 1898, pp. 226-254, 6 figs.
- Rivers of North America, a reading lesson for students of geography and geology. New York, 1898. xix, 327 pp., 17 pls.
- Review of Outlines of the Earth's History, by N. S. Shaler. *Science*, new series, vol. 8, 1898, pp. 712-715.
- Review of Lavas and Soils of the Hawaiian Islands, by Walter Maxwell. *American Naturalist*, vol. 32, 1898, pp. 537-539.
- The glaciers of North America. *Geographical Journal*, vol. 12, no. 6, December, 1898, pp. 553-564.
- Remarks on the use of the term plutonic plugs. *Journal of Geology*, vol. 7, 1899, pp. 96-97.
- Review of "The physiography and geology of the Nicaraguan Canal route." by C. Willard Hayes. *American Naturalist*, vol. 33, 1899, pp. 679-688.
- Geology of Cascade mountains of Washington. Abstract, *American Geologist*, vol. 23, p. 96 ($\frac{1}{2}$ p.); *Science*, new series, vol. 9, 1899, pp. 103-104.
- A preliminary paper on the geology of the Cascade mountains in northern Washington. U. S. Geological Survey, 20th Annual Report, pt. 2, 1900, pp. 89-210, pls. viii-xx, figs. 3-4.

- Deposits of calcareous marl in Michigan. Abstract, *Science*, new series, vol. 11, 1900, p. 402 ($\frac{1}{2}$ p.).
- Topographic atlas of the United States. *Science*, new series, vol. 12, 1900, pp. 1003-1004.
- Geology and water resources of Nez Perces county, Idaho, part i. U. S. Geological Survey, Water Supply and Irrigation Papers no. 53, 1901, pp. 1-85, 10 pls., 4 figs. Abstract, *American Geologist*, vol. 28, 1901, pp. 319-321.
- Geology and water resources of Nez Perces county, Idaho, part ii. U. S. Geological Survey, Water Supply and Irrigation Papers no. 54, 1901, pp. 95-141, 10 figs.
- Report to the National Geographic Society on the recent volcanic eruptions in the West Indies. *National Geographic Magazine*, vol. 13, 1902, pp. 267-285, 8 figs.
- Volcanic eruptions on Martinique and Saint Vincent. *National Geographic Magazine*, vol. 13, 1902, pp. 415-436, 10 figs.
- Geology and water sources of the Snake River plains of Idaho. Bulletin no. 190. U. S. Geological Survey, 1902, 192 pp., 25 pls., 6 figs.
- The Portland cement industry in Michigan. U. S. Geological Survey, 22d Annual Report, pt. 3, 1902, pp. 629-685, 3 pls.
- Geology of the Snake River plains, Idaho. Abstract, *Science*, new series, vol. 15, 1902, pp. 85-86.
- Notes on the geology of southwestern Idaho and southeastern Oregon. Bulletin no. 217, U. S. Geological Survey, 1903, 83 pp., 18 pls., 2 figs.
- Preliminary report on artesian basins in southwestern Idaho and southeastern Oregon. U. S. Geological Survey, Water Supply and Irrigation Paper no. 78, 1903, 51 pp., 2 pls., 3 figs.
- Volcanic eruptions on Martinique and Saint Vincent. Smithsonian Institution, Annual Report for 1902, pp. 331-349, 11 pls., 1903. Reprinted by permission, after revision by the author, from the *National Geographic Magazine*, vol. 13, no. 12, December, 1902. See above.
- Glacier cornices. *Journal of Geology*, vol. 11, 1903, pp. 783-785, 1 fig.
- The Pelé obelisk. *Science*, new series, vol. 18, 1903, pp. 792-795.
- Geography and international boundaries. American Geographical Society, Bulletin, vol. 35, 1903, pp. 147-159.
- Criteria relating to massive, solid volcanic eruptions. *American Journal of Science*, 4th series, vol. 17, 1904, pp. 253-268, 3 figs.
- Physiographic problems of today. *Journal of Geology*, vol. 12, 1904, pp. 524-550. Congress of Arts and Science, Universal Exposition, St. Louis, 1904.
4. 1906, pp. 627-649.
- North America (Appleton's World Series: The regions of the world). New York, 1904. 435 pp., 8 pls., 39 figs.
- The topographic survey of Michigan. Fifth Annual Report of the Michigan Academy of Science, 1904, pp. 149-165.
- Douglass Houghton. Michigan Academy of Science, 4th Report, 1904, pp. 160-162 (por.).
- Bela Hubbard. Michigan Academy of Science, 4th Report, 1904, pp. 163-165 (por.).

- Research in state universities. A paper read before the Research Club of the University of Michigan, January 20, 1904. *University Bulletin*, Ann Arbor, Michigan, new series, vol. 5, no. 5, 1904, 24 pp.
- Review of The Harriman Alaska Expedition. Vol. iii, Glaciers and Glaciation. *Science*, new series, vol. 19, 1904, pp. 783-787.
- Timberlines. *National Geographic Magazine*, vol. 15, no. 1, 1904, pp. 47-49.
- Abstract, Bulletin Geological Society of America, vol. 14, 1904, pp. 556-557.
- Biographical notice of William Henry Pettee. *American Geologist*, vol. 35, 1905, pp. 1-4, 1 pl. (por.).
- Preliminary report on the water supply of the Ann Arbor Water Company. Ann Arbor, Michigan, Council Proceedings, November 13, 1905, pp. 153-174.
- The influence of caverns on topography. *Science*, new series, vol. 21, 1905, pp. 30-32.
- Hanging valleys. Bulletin Geological Society of America, vol. 16, 1905, pp. 75-90.
- Preliminary report on the geology and water resources of central Oregon. Bulletin no. 252, U. S. Geological Survey, 1905, 138 pp., 24 pls.
- Drumlin areas in northern Michigan. Abstract, *American Geologist*, vol. 35, 1905, pp. 177-179; *Science*, new series, vol. 21, 1905, pp. 220-221.
- The Pelé obelisk once more. *Science*, new series, vol. 21, 1905, pp. 924-931, 1 fig.
- A geological reconnaissance along the north shore of lakes Huron and Michigan. Michigan Geological Survey, Report for 1904, pp. 33-112, 3 pls., 1 map, 1905.
- Memoir of William Henry Pettee. Bulletin Geological Society of America, vol. 16, 1906, pp. 558-560.
- Drumlin areas in northern Michigan. Seventh Annual Report of the Michigan Academy of Sciences, 1905, pp. 36-37. Abstract, Bulletin Geological Society of America, vol. 16, 1906, pp. 577-578.
- Concentration as a geological principle. Presidential address. Bulletin Geological Society of America, vol. 18, 1907, pp. 1-28.

MEMOIR OF NATHANIEL SOUTHGATE SHALER

BY JOHN E. WOLFF

Nathaniel Southgate Shaler was born at Newport, Campbell county, Kentucky, February 20, 1841, and died in Cambridge, Massachusetts, April 10, 1906. His father's people migrated from England to Jamaica, and from there to New York state and Connecticut. His father, Doctor Nathaniel Burger Shaler, graduated from Harvard College in 1827 and from the Medical School in 1829; he then went to Cuba, and from there to Newport, Kentucky, opposite Cincinnati, where he settled and practiced his profession. He married Ann Hind Southgate, of a prominent family of that name, of Virginia origin. Doctor Shaler was not only a highly educated man in the usual sense, but he had a taste for natural history

There is a world of work on hand.

Faithfully yours,

N S Palmer

and knew something of geology. He owned what we might call an experimental farm a few miles from Newport, where in his spare time he cultivated new varieties of seed, tobacco, and grapes, and also experimented with breeds of sheep. The son was much about with his father in the open, often galloping out to the farm from town on a pet mare, and was encouraged by his father in making experiments of his own with plants and in collecting specimens interesting to a boy, such as minerals, arrow-heads, and so on. The boy's health was delicate, so his school attendance was irregular and his formal training was acquired from a learned German named Escher, with whom his studies were of the philosophic and literary kind. From his mother he inherited his intense energy and quickness of repartee, and from her father he absorbed much of general culture, for his grandfather was a good Greek and Latin scholar, of a wide acquaintance with things intellectual, to whose discerning talk the boy listened eagerly. Thus he grew up constantly in the open and trained in out-of-door accomplishments, familiar with plants and animals and loving Nature, while at home his appreciation for the things in books was growing under this personal environment.

Young Shaler entered the Lawrence Scientific School in 1859, when 18 years old, registering in the course in zoology and geology, under Louis Agassiz, whose fame attracted students from the entire country. Among his classmates are found the names of Alexander Agassiz, Daniel C. Eaton, Alpheus Hyatt, F. W. Putnam, A. E. Verrill, Burt G. Wilder, A. S. Bickmore, A. S. Packard, and others, men who have spread the study of natural history in this country. While a student, he with Verrill and Hyatt made an excursion in 1861 to the island of Anticosti, studying the geology and collecting fossils, the first of his many wanderings over the surface of North America. He received his degree of B. S. in 1862, when 21 years old; but, without waiting for commencement, enlisted in the Federal army in Kentucky and was commissioned as captain of the Fifth Kentucky battery, called "Shaler's," and served two years, mainly in Kentucky. A volume of poetry published since his death deals with this phase of his life. Returning in 1864 to the Museum of Comparative Zoology, he was appointed "Assistant in Paleontology," with general charge, under Louis Agassiz, of that department. In this year, when 23 years old, he began his first public course of lectures at the Museum, on the "Geological succession of the brachiopoda" and on the "Elevation of continents." In 1866 he was obliged to suspend his work, owing to ill health, and spent over a year traveling in Europe, visiting many museums, seeing classical geological districts, and meeting such men as Darwin, Lyell, and Elie de Beaumont.

An incident illustrating Mr. Shaler's love of fun is that when calling on Darwin he brought with him a spirited American cartoon representing the old ape instructing the youthful Darwin, which delighted the great naturalist.

Soon after resuming his work at Cambridge he received his first formal appointment from the corporation of Harvard College as "Lecturer on paleontology and animal life considered in its geological relations," and a year later, in 1869, when 28 years old, was appointed "Professor of Paleontology" (changed in 1888 to "Professor of Geology"). His second exploration, namely, an excavation for fossil bones at Big Bone lick, Kentucky, was made in 1868; and now, besides giving lectures to undergraduates in zoology and paleontology, he began to supplement Louis Agassiz's lectures in geology by field excursions. Gradually the latter turned over to him much of the instruction in these subjects as ill health or his many undertakings necessitated.

A third excursion was made with students in 1870 to the James river, Virginia, and to other states to collect fossils. Resigning to others instruction in zoology, Professor Shaler now concentrated his teaching on paleontology and geology, especially the latter, and, having seen the establishment of the Summer School of Zoology at Penikese in 1873, he inaugurated the first summer school of geology himself, with the famous session of 1875, at Cumberland Gap, Virginia.

In 1873 he was appointed State Geologist of Kentucky, retaining this office seven years. The original survey, begun in 1854 by David Dale Owen, had lapsed, most of the records had been destroyed by fire, and nearly all of the assistants had passed away, and yet it was specified in the law establishing the new survey that it should begin where Owen's left off. This appealed to Mr Shaler's sense of humor, but he proceeded to carry out the plan as far as he could. The results are shown in five volumes, published under his direction, dealing with the resources, geology, and natural history of the state. During the years of this directorship he spent his summers largely in Kentucky and adjacent parts of Virginia, permitting a number of advanced students of geology to accompany the survey parties, who not only paid their own expenses, but often assisted in the work in any humble way permitted and were proud of the privilege. At the same time the geological department which he had founded at Harvard was rapidly growing and William M. Davis began to take a share of the instruction.

In 1881-1882 Shaler was in Europe, spending part of the time in Italy. Soon after his return he became one of the commissioners for the topographical survey of Massachusetts, which, in cooperation with the U. S.

Geological Survey under Major Powell, soon completed the topographical map of the state.

In 1884 Professor Shaler was appointed in charge of the Atlantic Coast Division of the National Survey and continued this connection until 1900, in the last seven years as a separate party. The original field of this division included the investigation of coastal lands, changes of level, phosphate beds, bog ores, etc., near the coast, but it soon broadened to cover many areas of rock geology, especially the Mount Desert island, the Narragansett coal-field and adjacent territory, and the Richmond, Virginia, coal-field. The surface geology of nearly the whole of New England was ultimately included; so that in the sixteen years of his connection with the U. S. Geological Survey he and his associates contributed many large and small papers to the annual reports and bulletins of the Survey and to outside publications.

In connection with this and earlier work done for the Coast Survey, Mr Shaler used to say he had traveled on foot nearly the whole coastline from Maine to Florida, which a hearer who was familiar with his long strides had no difficulty in believing. He made two explorations of the Dismal swamp, a trip through the marsh district about the mouth of the Mississippi, and a visit to the swamps of Michigan, Wisconsin, Minnesota, and Dakota.

In the winter of 1887-1888 he went to Florida, examining the phosphate beds southerly from Charleston, and then the Everglades and 360 miles of the east coast of Florida. On this trip the boat in which the party were sailing capsized during the night near an uncharted reef. The shore was reached with the greatest difficulty, and then followed a long walk to the nearest habitation. Mr Shaler published an article on this unexpectedly discovered reef.

Among his more important Survey publications resulting from this period may be mentioned:

In 1885: "Preliminary report on seacoast swamps of the eastern United States."

1888: "Report on the geology of Marthas Vineyard."

1889: "The geology of Cape Ann, Massachusetts;" "The geology of the island of Mount Desert," and "The geology of Nantucket."

1890: "General account of the fresh-water morasses of the United States, with a description of the Dismal Swamp district of Virginia and North Carolina."

1892: "The nature and origin of soils."

1893: "The geological history of harbors."

1895: "Peat deposits."

1898: "Geology of the Cape Cod district."

And in joint authorship with others:

In 1888: "On the geology of the Cambrian district of Bristol county, Massachusetts."

1896: "The glacial brick clays of Rhode Island and southeastern Massachusetts."

1899: "Geology of the Narragansett basin" and "Geology of the Richmond basin, Virginia." In the course of this work Mr Shaler discovered the Cambrian fossils of Attleboro, Massachusetts, and the Cretaceous of Marthas Vineyard.

In this same period, in addition to his service to the state of Massachusetts in connection with the topographic map, Mr Shaler served on two other commissions, the national importance of which is for different reasons now apparent. The first was as a member of a committee of the State Board of Agriculture for the suppression of the gypsy moth. He entered with vigor into the fight against this pest, taking a prominent part in the organization of the work and in attempting to impress on the public the need of doing it in time. The second commission had a more grateful task, namely, that of improving the highways of Massachusetts, and Mr Shaler was one of the original three members of the Massachusetts Highway Commission, which in its ideals and methods of work has served as a model for other states.

I will quote from a letter of the present chairman, Mr W. E. McClintock, dealing with Mr Shaler's part in the beginning of this great movement. He says:

"Professor Shaler was an enthusiastic advocate of good roads, and his numerous writings and talks brought a very large number of educated thinking men to the support of the movement, and to him more than to any one man is due the strong support of the commission by this class of men. His facile pen made the annual report of the commission held in such high repute that the usual editions were exhausted before the demand ceased."

The Geological Survey published a paper of his in 1895 on the geology of the road-building stones of Massachusetts and elsewhere. In this field he had organized a laboratory for physical tests of road materials, a work which was subsequently taken up by the U. S. Department of Agriculture.

In addition to such work, allied to economic geology, he at various times reported and advised on mines and quarries, and in his long list of publications are found several dealing with building stones. In the latter years he became interested in the question of working low-grade placer deposits, and finally organized and directed a large and successful undertaking of this kind in Montana. These various interests, theoretical and

practical, took him to many parts of this continent, including Cuba and Alaska, so that it would perhaps have been difficult to find any one who had seen understandingly more of the country and its people.

These travels and investigations, covering thirty-five crowded years, had been carried on simultaneously with his university duties, which were constantly growing in extent and importance.

No instructor at Harvard has held the position which Shaler did in the eyes and in the hearts of the students. His elementary course of lectures on the principles of geology ("Natural History," afterwards "Geology 4") reached the high-water mark of attendance, with between five and six hundred hearers, and many working geologists of the country can say with the writer that their first interest in natural history, especially in geology, was inspired by this elementary course. He also taught elementary paleontology and advanced geology, always aiming to present underlying principles, but with less attention to detail. It is estimated that about seven thousand students came under his direct instruction in this long term of years, probably a greater number than had before been taught this subject by any one man. Until his administrative duties prevented, he was much in the field, both summer and winter, with advanced students.

His teaching, however, was only a part of his activity in the university; for, besides the summer schools, he originated and directed many other ways of assisting the students, especially those who were strangers or were poor. He was ready with advice and sympathy, often of a material kind, for any who came to him in trouble.

The great university task and achievement of the last fifteen years of his life was the successful reorganization and development of the Lawrence Scientific School, which had been overshadowed by the development of Harvard College. Mr Shaler became dean in 1891, and vigorously took up the work of enlarging the instruction and resources and of attracting students; not only were the existing departments remodeled, but new ones were added from time to time, such as, for instance, that of mining and metallurgy, which at the time of his death had become a flourishing and well equipped department. The result was seen in an increase in attendance from 118 in 1891 to 504 in 1906. His friend, Gordon Mackaye, left his great fortune for the endowment of applied science in the university, and at the time of his death Mr Shaler was actively planning the development to follow from the application of this fund.

Mr Shaler's personality was striking and original. He was tall and well proportioned, neither slender nor stout, with erect, active carriage; he had prominent features, with large blue-gray eyes and thick hair, brushed back from his high forehead. In lecturing or speaking he rarely

used gestures, his favorite pose being to rest both hands on a book. And what *flashing* speaking that was, whether formal or informal; never hurried or undignified, but in turn impressive or humorous or picturesque and always interesting. The charm extended to all kinds of people. I have seen the rough mountaineers of the south listening fascinated to his words of explanation to the students with him. Perhaps some intuition of the kindness of heart which underlay his words and actions had to do with this, for Mr Shaler was above all a lover of his fellow-men and what concerned them; this was well expressed when in 1903 the degree of LL.D. was conferred on him by Harvard as "naturalist and humanist."

With all this activity in so many different lines, he yet found time to practice a warm hospitality at his home in Cambridge. He was married in 1862 to Miss Sophia Page, a resident of Kentucky, who with two daughters, Mrs Willoughby L. Webb and Mrs Walter L. Page, survives him. The host and hostess welcomed to their home not only those who might fairly be expected to offer some mutual entertainment, but also with true kindness the young and often bashful newcomers in the college.

Professor Shaler's publications dealing with geology and kindred topics number nearly two hundred titles besides many popular articles in the magazines. They deal with paleontology (especially in the earlier years), glaciation, mountain-building, shorelines, earthquakes, the elevation of continents, swamps and peat-deposits, soils, phosphate beds, petroleum, coal, building stones, changes of climate, caverns, and many others. He published various separate works, such as text-books and popular works dealing with various features of geology, always emphasizing the relation to man. His "First Book of Geology" was translated into Polish. His last important scientific publication, "A comparison of the features of the earth and the moon," published in 1903 by the Smithsonian Institution, with superb plates, is an interesting attempt to explain lunar features by the experience of geology. It illustrates an unexpected persistency in following certain lines of research, for his work with the telescope had been done thirty years earlier. I am told that for several years before his death he had been studying the coast charts of the world in continuation of his earlier work on shorelines and had even visited Egypt with this in mind.

Among his miscellaneous writings mention should be made of a series of philosophical essays entitled "The citizen, a study of the individual and the government;" "The individual, a study of life and death;" "The neighbor, the natural history of human contacts," and others, which deal with these subjects in an original way from the point of view of the naturalist.

Lastly, within a few years Mr Shaler ventured into the field of general literature and poetry. His drama, "Elizabeth of England," has been highly praised as a noble work of imagination and diction. A posthumous volume of poetry, "From old fields," has been mentioned.

It is too early to estimate the value of his formal contributions to American geology, of which but an imperfect account has been given, but there can be no doubt as to the immense influence he exerted as a teacher and expositor of the subject. He recognized the preeminent place accorded to him in this respect by choosing as the subject of his presidential address to this Society in 1895 "The relations of geologic science to education." It is perhaps as one of the great teachers of geology that he will be best remembered by us, and may the inspiration he gave long continue to bear fruit.

BIBLIOGRAPHY*

1861. Lateral symmetry in brachiopoda. *Proc. Bost. Soc. Nat. Hist.*, vol. 8, 1861-1862, pp. 274-279.
1862. On the Geology of Anticosti island, gulf of Saint Lawrence. *Proc. Bost. Soc. Nat. Hist.*, vol. 8, 1862, pp. 285-287.
1865. List of the brachiopoda from the island of Anticosti, sent by the Museum of Comparative Zoology to different institutions in exchange for other specimens, with annotations. *Bull. Mus. Comp. Zool.*, vol. 1, 1865, pp. 61-70.
1866. Preliminary notice of some opinions concerning the mode of elevation of continental masses. *Proc. Bost. Soc. Nat. Hist.*, vol. 10, 1866, pp. 237-239.
 On the modifications of ocean currents in successive geological periods. *Proc. Bost. Soc. Nat. Hist.*, vol. 10, 1866, pp. 296-302.
 On the formation of the excavated lake basins of New England. *Proc. Bost. Soc. Nat. Hist.*, vol. 10, 1866, pp. 358-366.
1868. Notes on the position and character of some glacial beds containing fossils at Gloucester, Massachusetts. *Proc. Bost. Soc. Nat. Hist.*, vol. 11, 1868, pp. 27-30.
 On the formation of mountain chains. *Proc. Bost. Soc. Nat. Hist.*, vol. 11, 1868, pp. 8-15.
1869. On the nature of the movements involved in the changes of level of shorelines. *Proc. Bost. Soc., Nat. Hist.*, vol. 12, 1869, pp. 128-136.
 On the disappearance of the cane from the central part of the Ohio valley. *Proc. Bost. Soc. Nat. Hist.*, vol. 12, 1869, pp. 136-137.
 Considerations concerning the absence of distinct evidences of glacial action in the valley of the Yukon river, Alaska. *Proc. Bost. Soc. Nat. Hist.*, vol. 12, 1869, pp. 145-149.
 Notes on the concentric structure of granitic rocks. *Proc. Bost. Soc. Nat. Hist.*, vol. 12, 1869, pp. 289-293.

* Owing to the unusual range of subjects covered by Professor Shaler, it has been thought worth while to place on record as complete a bibliography as possible of his writings.

- On changes in geographical distribution of the American buffalo. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, p. 136.
- Note on the occurrence of the remains of *Tarandus rangifer* Gray at Big Bone Lick, in Kentucky. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, p. 167.
1870. On the phosphate beds of South Carolina. U. S. Coast Survey Report for 1870, pp. 182-189. Washington, 1870.
1871. Note on the glacial moraines of the Charles River valley, near Watertown. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, pp. 277-279.
- On the phosphate beds of South Carolina. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, pp. 222-236.
- On the parallel ridges of Glacial drift in eastern Massachusetts, with some remarks on the Glacial period. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, pp. 196-204.
- On the relations of the rocks in the vicinity of Boston. *Proc. Bost. Soc. Nat. Hist.*, vol. 13, 1871, pp. 172-177. Abstract, *American Naturalist*, vol. 4, 1871, p. 238.
1872. On the causes which have led to the production of cape Hatteras. *Proc. Bost. Soc. Nat. Hist.*, vol. 14, 1872, pp. 110-121. Abstract, *American Naturalist*, vol. 5, 1871, pp. 178-181.
- On the geology of the island of Aquidneck and the neighboring parts of the shores of Narragansett bay. *American Naturalist*, vol. 6, 1872, pp. 518-528, 611-621, 751-760, map. From Report of U. S. Coast Survey.
- Source of boulders in Ohio, Kentucky, etc. *Proc. Bost. Soc. Nat. Hist.*, vol. 14, 1872, p. 386.
1873. Note on the origin of our domestic cat. *Proc. Bost. Soc. Nat. Hist.*, vol. 15, 1873, pp. 159-162.
- On the effects of the vertical position in man. *Proc. Bost. Soc. Nat. Hist.*, vol. 15, 1873, pp. 188-191.
- On elongation of pebbles. *Proc. Bost. Soc. Nat. Hist.*, vol. 15, 1873, p. 2.
- Geology of Marthas Vineyard and Nantucket. *Proc. Bost. Soc. Nat. Hist.*, vol. 15, 1873, p. 219.
- Mixed populations of North Carolina. *North American Review*, vol. 116, 1873, pp. 150-166.
1874. Recent changes of level on the coast of Maine, with reference to their origin and relation to other similar changes. *Mem. Bost. Soc. Nat. Hist.*, vol. 2, 1874, pp. 321-340.
- On the geology of the region about Richmond, Virginia, etc. *Proc. Am. Acad. of Arts and Sciences*, vol. 8, 1874, pp. 307-308.
- Report of special committee [of experts, to a committee of the state board of education, on a proposed scientific survey of Massachusetts]. [Benjamin Peirce: Report of special committee, etc., 1874.]
1875. Notes on some of the phenomena of elevation and subsidence of the continents. *Proc. Bost. Soc. Nat. Hist.*, vol. 17, 1875, pp. 288-292.
- Some considerations on the possible means whereby a warm climate may be produced within the Arctic circle. *Proc. Bost. Soc. Nat. Hist.*, vol. 17, 1875, pp. 332-337.
- Note on some points connected with tidal erosion. *Proc. Bost. Soc. Nat. Hist.*, vol. 17, 1875, pp. 465-466.

- Note on the geological relations of Boston and Narragansett bays. *Proc. Bost. Soc. Nat. Hist.*, vol. 17, 1875, pp. 488-490.
- Remarks on the geology of the coast of Maine, New Hampshire, and that part of Massachusetts north of Boston. U. S. Coast Survey, Coast Pilot for the Atlantic seaboard, Gulf of Maine, and its coast from Eastport to Boston, pp. 883-888, Washington, 1875.
- Question guide to the environs of Boston. Designed for the use of beginners in geology in the classes of Harvard University. Pt. 1. Somerville and Cambridge, 1875.
- Frequent reports on paleontology and geology in the annual reports of the trustees of the Museum of Comparative Zoology at Harvard College, in Cambridge, for the years 1864-1900.
1876. On the prehistoric remains of Kentucky. (Lucien Carr and N. S. Shaler.) Kentucky Geol. Survey Mem., vols. 1, 4. Cambridge, 1876.
- Brief statement of the objects, methods, and results of the survey, 1873-1876. Kentucky Geol. Survey.
- On the fossil brachiopods of the Ohio valley. Kentucky Geol. Survey Mem., etc., vol. 1. Pt. 1. Cambridge.
- On the antiquity of the caverns and cavern life of the Ohio valley. Kentucky Geol. Survey Mem., vol. 1, pt. 1, 13 pp. Cambridge, 1876.
- Abstract, *American Journal of Science*, 3d series, vol. 13, 1877, pp. 226-227.
- Report on the forest timber of Greenup, Carter, Boyd, and Lawrence counties. (N. S. Shaler and A. R. Crandall.) Kentucky Geol. Survey, N. S., vol. 1, pt. 1, 1876, pp. 1-26.
- Introduction to report on the botany of Barren and Edmonson counties by John Hussey. Kentucky Geol. Survey, N. S., vol. 1, pt. 2, 1876, pp. 27-58.
1877. On the origin of the galena deposits of the Upper Cambrian rocks of Kentucky. Kentucky Geol. Survey, reports of progress, N. S., vol. 2, 1877, pp. 277-292.
- A general account of the Commonwealth of Kentucky, prepared by the Geological Survey of the Commonwealth. Kentucky Geol. Survey, reports of progress, N. S., vol. 2, 1877, pp. 361-468, map. Published separately, Frankfort, 1876. Map republished in black in Report of Kentucky Commission of Agriculture, 1877.
- On the geographical and geological distribution of the genus *Beatricea*, and of certain other fossil corals in the rocks of the Cincinnati group. Title in *Proc. A. A. A. S.*, vol. 26, 1877.
- On the recent formation of a small anticlinal axis in Lincoln county, Kentucky. Title in *Proc. A. A. A. S.*, vol. 26, 1877.
- On the continuation of the fold of the Alleghany chain to the north of the Delaware river. Title in *Proc. A. A. A. S.*, vol. 26, 1877.
- Account of some lunar phenomena. Title in *Proc. Am. Acad. Arts and Sciences*, vol. 38, 1903.
- Geological Survey of Kentucky, reports of progress, N. S., vol. 3, 1877, 451 pp., 5 maps.
- General report of the Geological Survey of Kentucky. Kentucky Geol. Survey, N. S., vol. 3, pt. 1, 1877, pp. 1-30.

- History of the operations of the survey in 1874 and 1875. Kentucky Geol. Survey, N. S., vol. 3, pt. 2, pp. 31-127.
- Notes on the investigations of the Kentucky Geological Survey during the years 1873, 1874, and 1875. Kentucky Geol. Survey, N. S., vol. 3, pt. 3, pp. 129-240.
- Annual report of N. S. Shaler for the year 1876. Kentucky Geol. Survey, N. S., vol. 3, pt. 4, 1877, pp. 283-315.
- The transportation routes of Kentucky and their relation to the economic resources of the commonwealth. Kentucky Geol. Survey, N. S., vol. 3, pt. 5, pp. 316-346.
- Description of the preliminary topographical and geological maps of Kentucky, edition of 1877. Kentucky Geol. Survey, N. S., vol. 3, pt. 6, pp. 347-364.
- Annual report of N. S. Shaler, state geologist, for 1877. Kentucky Geol. Survey, N. S., vol. 3, pt. 7, pp. 365-414.
- Report on the unfinished work of the survey of the commonwealth under the direction of Dr David Dale Owen. Kentucky Geol. Survey, N. S., vol. 3, pt. 8, pp. 415-420.
- On the cause and geological value of variation in rainfall. *Proc. Bost. Soc. Nat. Hist.*, vol. 18, 1877, pp. 176-182.
- Propositions concerning the motion of continental glaciers. *Proc. Bost. Soc. Nat. Hist.*, vol. 18, 1877, pp. 126-133.
- Notes on the age and the structure of the several mountain axes in the neighborhood of the Cumberland gap. *American Naturalist*, vol. 11, 1877, pp. 385-392.
- On the existence of the Alleghany division of the Appalachian range within the Hudson valley. *American Naturalist*, vol. 11, 1877, pp. 627-628.
1878. Thoughts on the nature of intellectual property and its importance to the state. Boston, 1878.
- Mammoth Cave. Johnson's New Universal Encyclopedia.
- Recent changes of level of the coast of Maine. *Mem. Bost. Soc. Nat. Hist.*, vol. 2, 1878, pp. 320-341.
1879. Petroleum. Kentucky Geol. Survey. Bull. no. 1, pp. 5-20.
- On the Improvement of the rivers of Kentucky. Kentucky Geol. Survey, Bull. no. 2, pp. 13-21.
- On the Importance of Improvement in the Navigation of the Kentucky River to the mining and manufacturing interests of Kentucky. N. S. Shaler and J. R. Procter. Kentucky Geol. Survey, Bull. no. 3, pp. 22-45.
1880. Summary of the work of the Geological Survey for the years 1878-1879. Kentucky Geol. Survey.
- Outline of the geology of Boston and its environs. (Justin Winsor, editor of The Memorial History of Boston, etc., vol. 1, pp. 1-8.)
- Proposition concerning the classification of lavas, considered with reference to the circumstances of their extrusion. *Bost. Soc. Nat. Hist.*, Anniversary Memoirs, 1880, 15 pp.
- Preliminary report on the natural resources of the country. Cambridge, Massachusetts. 4 pp.

- Preliminary report concerning the resources of the country adjacent to the line of the proposed Richmond and Southwestern railway. Map. Cambridge.
1881. The island of Campobello. Preliminary report. Cambridge, 1881, vol. 1, pp. 11.
- Notes on the submarine coast-shelf, or hundred-fathom detrital fringe. *Proc. Bost. Soc. Nat. Hist.*, vol. 20, pp. 278-282.
- Illustrations of the earth's surface. I. Glaciers. (N. S. Shaler and W. M. Davis.) Boston, 1881.
- Notes on certain evidences of a gradual passage from sedimentary to volcanic rocks shown in Brighton district of Boston. *Proc. Bost. Soc. Nat. Hist.*, vol. 20, 1881, pp. 129-133.
- Great Kanawha, West Virginia, iron ores and coals. The Black Band Iron and Coal Company. The Virginias, vol. 2, 1881, pp. 154-155.
1883. Report on the Croton magnetic iron mines. Cambridge, Massachusetts. 25 pp.
- On the recent advances and retrocessions of glaciers. *Proc. Bost. Soc. Nat. Hist.*, vol. 21, 1883, pp. 162-167.
1884. General report on the building stones of Rhode Island, Massachusetts, and Maine. 10th Census U. S., report on the building stones of the U. S. and statistics of the quarry industries for 1880, vol. 10, 1884, pp. 107-116.
- Report on the resources of the region adjacent to the proposed Cincinnati and Southwestern railway. Cambridge, Massachusetts. 28 pp. Map.
1885. Preliminary report on seacoast swamps of the eastern United States. U. S. Geol. Survey, 6th Annual Report, 1884-1885, pp. 353-398. Washington, 1885. Abstract, *American Geologist*, vol. 1, 1888, pp. 258-259.
- Kentucky: A pioneer commonwealth. Boston, Massachusetts, 1885. 16°.
- Report of the Atlantic coast division. U. S. Geol. Survey, 6th Annual Report, 1884-1885, pp. 18-22. Washington, 1885.
1886. A series of twenty-five colored geological models and twenty-five photographs of important geological objects, each accompanied by letterpress description. (N. S. Shaler, W. M. Davis, and T. W. Harris.) Boston, Massachusetts, D. C. Heath & Co.
- Preliminary report on the geology of the Cobscook Bay district, Maine. *American Journal of Science*, 3d series, vol. 32, 1886, pp. 35-60. Abstract, *American Naturalist*, vol. 20, 1886, p. 969.
1887. A first book in geology, designed for the use of beginners. Boston, 1887. The same. Boston, 1897.
- On the original connection of the eastern and western coal fields of the Ohio valley. *Mem. Mus. Comp. Zool.*, vol. 16, no. 2, 1887, pp. 1-11.
- Fluviatile swamps of New England. *American Journal of Science*, 3d series, vol. 33, 1887, pp. 210-221. Abstract, *Popular Science Monthly*, vol. 33, 1887, pp. 142-143.
- Field geology. *Popular Science Monthly*, vol. 21, June, 1887, pp. 80-82; July, 1887, pp. 94-96.
- The stability of the earth. *Scribner's Magazine*, 1887, pp. 259-279.

- Notes on the *Taxodium distichum*, or bald cypress. On the original connection of the eastern and western coal-fields of the Ohio valley. *Mem. Mus. Comp. Zool.*, 1887, vol. 16, pp. 1, 2.
- On the forests of North America. *Scribner's Magazine*, vol. 1, no. 5.
- On the origin and nature of rock gas. *The Forum*, April, 1887.
- On the instability of the atmosphere. *Scribner's Magazine*, vol. 2, no. 2.
- Caverns and cavern life. *Scribner's Magazine*, vol. 2, no. 5, pp. 449-472.
1888. Report on the geology of Marthas Vineyard. U. S. Geol. Survey. 7th Report. 1885-1886, pp. 297-363, pls. 19-29. Abstract, *Science*, vol. 13, p. 343, *American Geologist*, vol. 4, 1889, pp. 104-106.
- The crenetic hypothesis and mountain building. *Science*, vol. 11, 1888, pp. 280-281.
- Origin of the divisions between the layers of stratified rocks. *Proc. Bost. Soc. Nat. Hist.*, vol. 23, 1888, pp. 408-419.
- Report Atlantic Coast Division of Geology. U. S. Geol. Survey. 7th Report, 1885-1886, pp. 61-65. Washington, 1888.
- Volcanoes. *Scribner's Magazine*, 1888, pp. 201-226.
- Animal agency in soil working. *Popular Science Monthly*, vol. 32, no. 4, pp. 484-487.
- On the study of nature. *Popular Science Monthly* and *Boston Journal of Chemistry*, vol. 22, no. 3, p. 34, and no. 5, pp. 49-50.
- Introduction. Nature and origin of deposits of phosphates of lime. by R. A. F. Penrose, Jr. Bulletin no. 46, U. S. Geol. Survey, vol. 7, pp. 483-494. Washington, 1888. Abstract, *Science*, vol. 13, 1888, pp. 144-146.
- On the geology of the Cambrian district of Bristol County, Massachusetts. *Bull. Mus. Comp. Zool.*, vol. 16, no. 2, 1888, pp. 13-26. Map.
- Dzieje ziemi, czyli Początki geologii. Przełoży z angielskiego z upoważnienia autora Heuryk Wernic. Przerzrzedli i dopenili dopiskami Antoni Słóarski i Józef Siemeradzki. Warszawa. (A translation of his First Book in Geology.)
- Rivers and valleys. *Scribner's Magazine*, vol. 4, no. 2, 1888, pp. 131-155.
- On the origin of kames. *Proc. Bost. Soc. Nat. Hist.*, vol. 23, 1888, pp. 36-44.
- The Law of Fashion. *The Atlantic Monthly*, vol. 61, no. 365, pp. 386-398.
1889. The geology of Nantucket. Bulletin no. 53, U. S. Geol. Survey. vol. 8, pp. 601-653, 10 plates. Washington, 1889. Abstracts, *American Geologist*, vol. 5, 1890, pp. 111-114. *Popular Science Monthly*, vol. 36, 1890, p. 567.
- Physiography of North America. Narrative and critical history of America. Boston [1884], vol. 4, pp. i-xxx.
- The Geology of the island of Mount Desert, Maine. U. S. Geol. Survey. 8th Report, pp. 987-1061, plates 64-76. Washington, 1889. Abstract, *American Geologist*, vol. 6, 1890, pp. 197-198.
- Report, Atlantic Coast Division. U. S. Geol. Survey. 9th Report, pp. 71-74. Washington, 1889.
- Aspects of the earth: A popular account of some familiar geological phenomena. xix, 344 pages, 15 plates. New York, 1889. The same. Illustrated. New York, 1900.

- Report, Division of Coast Line Geology. *U. S. Geol. Survey, 8th Report*, pp. 125-128. Washington, 1889.
- On the occurrence of fossils of the Cretaceous age on the island of Marthas Vineyard, Massachusetts. *Bull. Mus. Comp. Zool.*, vol. 16, 1889, pp. 89-97, plates 1, 2.
- The Geology of Cape Ann, Massachusetts. *U. S. Geol. Survey, 9th Report*, pp. 529-611, plates 32-37. Washington, 1889. Abstract, *American Geologist*, vol. 7, 1891, p. 201.
- The athletic problem in education. [Boston, etc., 1889], 8°, pp. 10. *Atlantic*, 1889, 63, 79-88.
- Chance or design? *The Andover Review*, August, 1889, pp. 1-17.
- Effects of permanent moisture on certain forest trees. *Science*, March, 1889, pp. 176-177.
1890. General account of the fresh-water morasses of the United States, with a description of the Dismal Swamp district of Virginia and North Carolina. *U. S. Geol. Survey, 10th Report*, pp. 255-339, plates 6-19. Washington, 1890. Abstract, pp. 15-16.
- Remarks on conditions attending a Pleistocene submergence on the Atlantic coast. *Bull. Geol. Soc. Am.*, vol. 1, 1890, p. 409.
- The topography of Florida. *Bull. Mus. Comp. Zool.*, vol. 16, pp. 139-156, plate. Abstract, *American Naturalist*, vol. 24, 1890, p. 768.
- Note on glacial climate. *Proc. Bost. Soc. Nat. Hist.*, vol. 24, 1890, pp. 460-465. Abstract, *American Geologist*, vol. 5, 1890, p. 124.
- Rock gases. *The Arena*, 1890, vol. 1, pp. 631-642.
- Report. Atlantic Coast Division. *U. S. Geol. Survey, 10th Report*, pp. 117-119. Washington, 1890.
- Soils of Massachusetts. (Lecture delivered at the annual meeting of the Massachusetts State Board of Agriculture, February 6, 1890.) Boston, 1890.
- Note on the value of saliferous deposits as evidence of former climatal conditions. *Proc. Bost. Soc. Nat. Hist.*, vol. 24, 1890, pp. 580-585.
- Tertiary and Cretaceous deposits of eastern Massachusetts. *Bull. Geol. Soc. Am.*, vol. 1, 1890, pp. 443-452. Abstract, *American Geologist*, vol. 5, p. 118; *American Naturalist*, vol. 24, p. 210; *Science*, vol. 15, p. 16. All 1890.
- The nature of the negro. *The Arena*, 1890, pp. 23-35.
- The economic future of the new south. *The Arena*, August, 1890, pp. 257-268.
1891. Report of the Massachusetts Topographical Survey Commission. January, 1891.
- Atlas of Massachusetts. (N. S. Shaler, F. A. Walker, and H. L. Whiting.) Fifty-five sheets. January, 1891.
- The antiquity of the last Glacial period. *Proc. Bost. Soc. Nat. Hist.*, vol. 25, 1891, pp. 258-267.
- The history of our continent. 290 pages. Boston, 1891.
- Nature and man in America. New York, 1891. The same. New York, 1899.
- Individualism in education. *Atlantic Monthly*, vol. 67.

- College examinations. *Atlantic Monthly*, July, 1891.
1892. The origin and nature of soils. U. S. Geol. Survey, 12th Report, pt. 1. 1892, pp. 219-345. Abstract, *American Journal of Science*, 3d series, vol. 45, 1893, pp. 163-164; *American Geologist*, vol. 14, 1894, pp. 114-115.
- The depths of the sea. *Scribner's Magazine*, vol. 12, 1892, pp. 77-95.
- The story of our continent: A reader in the geography and geology of North America. Boston, 1892. The same. Boston, 1897.
- The border state men of the civil war. *Atlantic Monthly*, 1892, 69 pp. 245-257.
- [Remarks on the life of Samuel Dexter.] *Proc. Bost. Soc. Nat. Hist.*, vol. 25, 1892, pp. 365-366.
- The betterment of our highways. [Boston, 1892], 8°, pp. (12). *Atlantic Monthly*, October, 1892.
1893. The interpretation of nature. (Contains, with slight modification, the course of lectures on the Winkley foundation, delivered before the students of Andover Theological Seminary in 1891. Preface.) Boston, Massachusetts, 1893.
- The geological history of harbors. U. S. Geol. Survey, 13th Report, pt. 2. 1893, pp. 99-209, pls. xxii-xlv, figs. 7-15.
- High buildings and earthquakes. *North American Review*, vol. 156, 1893, pp. 338-345.
- Man and the Glacial period. *American Geologist*, vol. 11, 1893, pp. 180-184.
- The conditions of erosion beneath deep glaciers, based upon a study of the bowlder train from Iron hill, Cumberland, Rhode Island. *Bull. Mus. Comp. Zool.*, vol. 16, no. 11, pp. 185-225, plates i-iv, and map. Abstracts, *American Geologist*, vol. 12, 1893, pp. 191-192; *American Naturalist*, vol. 27, 1893, p. 662.
- With G. S. Perkins and W. E. McClintock. Report of the Commission to Improve the Highways of the Commonwealth of Massachusetts. Boston, 1893, pp. 238.
1894. Pleistocene distortions of the Atlantic seacoast. *Bull. Geol. Soc. Am.*, vol. 5, 1894, pp. 199-202.
- Relation of mountain growth to formation of continents. *Bull. Geol. Soc. Am.*, vol. 5, 1894, pp. 203-206.
- Phenomena of beach and dune sands. *Bull. Geol. Soc. Am.*, vol. 5, 1894, pp. 207-212.
- Discussion on faceted pebbles. *Bull. Geol. Soc. Am.*, vol. 5, p. 608.
- Report of the Commission of the Topographical Survey of the Commonwealth of Massachusetts for the year 1893 to his Excellency Frederic T. Greenhalge. House, no. 74, pp. 3-30. (N. S. Shaler, H. L. Whiting, and D. Fitzgerald.)
- Sea and land: Features of coast and oceans with special reference to the life of man. New York, 1894.
- Editor geological terms in Standard dictionary of the English language. (N. S. Shaler and W. B. Dwight.) Vol. 1, New York, 1894.
- The United States of America: A study of the American commonwealth, its natural resources, people, industries, manufactures, commerce

- and its work in literature, science, education and self-government. New York, 1894. 2 volumes. (Editor.)
- On the distribution of earthquakes in the United States since the close of the Glacial period. *Proc. Bost. Soc. Nat. Hist.*, vol. 26, 1894, pp. 246-256. Abstract. *American Geologist*, vol. 14, 1894, pp. 396-397.
1895. Beaches and tidal marshes of the Atlantic coast. *Nat. Geol. Soc. Mon.*, vol. 1, no. 5, 1895, pp. 137-168.
- Dislocations of the Cretaceous and Tertiary rocks of Marthas Vineyard, Massachusetts. *Bull. Geol. Soc. Am.*, vol. 6, 1895, p. 7.
- Origin, distribution, and commercial values of peat deposits. U. S. Geol. Survey, 16th Annual Report, pt. 4, 1895, pp. 305-314.
- Editor geological terms in the Standard dictionary of the English language. New York, vol. 2, 1895. (N. S. Shaler, W. B. Dwight, and J. B. Woodworth.)
- Sea and land; Features of coasts and oceans, with special reference to the life of man. New York, Charles Scribner's Sons, 1895.
- Evidences as to change of sealevel. *Bull. Geol. Soc. Am.*, vol. 6, 1895, pp. 141-166.
- Certain features in the jointing and veining of the Lower Silurian limestones near Cumberland gap, Tennessee. Title in *Bull. Geol. Soc. Am.*, vol. 6, 1895, p. 443.
- Conditions and effects of the expulsion of gases from the earth. Title *Bull. Geol. Soc. Am.*, vol. 7, p. 11. *Proc. Bost. Soc. Nat. Hist.*, vol. 27, 1896, pp. 89-106.
- The share of volcanic dust and pumice in marine deposits. *Bull. Geol. Soc. Am.*, vol. 7 (abstract), 1895-1896, pp. 490-492.
- Second annual report of the Massachusetts Highway Commission. (N. S. Shaler, G. H. Perkins, and W. C. McClintock.) Public document no. 54, January, 1895, pp. 68.
- Some causes of the imperfection of the geologic record. *Science*, vol. 2, 1895, pp. 858-859.
- The geology of the road-building stones of Massachusetts, with some consideration of similar materials from other parts of the United States. U. S. Geol. Survey, 16th Annual Report, part 2, 1895, pp. 277-341, pls. xviii-xxiv.
- Preliminary report on the geology of the common roads of the United States. U. S. Geol. Survey, 15th Annual Report, 1895, pp. 259-306.
- Domesticated animals; their relation to man and to his advancement in civilization. New York, 1895, 8°.
1896. Third annual report of the Massachusetts Highway Commission (N. S. Shaler, G. A. Perkins, and W. E. McClintock). Public document no. 54, Boston, January, 1896, pp. 126.
- The economic aspects of soil erosion. *National Geographic Magazine*, vol. 7, 1896, pp. 328-338, 368-377.
- The glacial brick clays of Rhode Island and southeastern Massachusetts. U. S. Geol. Survey, 17th Report, 1896, pp. 957-1004, plates lxi-lxii, figs. 34-43. (With J. B. Woodworth and C. F. Marbut.)
- Discussion regarding low temperature Gradients in Mines. Abstract, *American Geologist*, vol. 17, 1896, p. 100.

- Discussion on the carriage of bowlders by the Indians. Abstract, *American Geologist*, 1896, p. 104.
- American highways; a popular account of their conditions and of the means by which they may be bettered. New York.
1897. Water supply of eastern Massachusetts. Abstract, *Science*, N. S., vol. 5, 1897, p. 703.
- Fourth annual report Massachusetts Highway Commission (N. S. Shaler, T. C. Mendenhall, and W. E. McClintock).
- Nansen's heroic journey. *Atlantic Monthly*, May, 1897, vol. 79, pp. 610-617.
1898. Geology of the Cape Cod district (Massachusetts). U. S. Geol. Survey, 18th Annual Report, pt. 2, 1898, pp. 503-593, pls. xcvi-civ, figs. 86-92.
- Outlines of the earth's history: A popular study in physiography. D. Appleton & Co., New York, 1898. (8° and 12 mo.)
1899. Loess deposits of Montana. *Bull. Geol. Soc. Am.*, vol. 10, 1899, pp. 245-252.
- Formation of dikes and veins. *Bull. Geol. Soc. Am.*, vol. 10, 1899, pp. 253-262.
- Spacing of rivers with reference to hypothesis of base-leveling. *Bull. Geol. Soc. Am.*, vol. 10, 1899, pp. 262-276.
- Geology of Narragansett basin. (N. S. Shaler, J. B. Woodworth, and A. F. Foerste.) U. S. Geol. Surv. Mon., vol. 33, 1899, 402 pp., 31 pls., 30 figs.
- Geology of the Richmond basin, Virginia. (N. S. Shaler and J. B. Woodworth.) Nineteenth Annual Report U. S. Geol. Survey, pt. 2, 1899, pp. 385-520, pls. xviii-liv, figs. 90-116.
1900. Influence of the sun upon the formation of the earth's surface. *Int. Mon.*, vol. 1, 1900, pp. 41-82.
- The individual; a study of life and death. New York, 1900, 8°.
- The negro since the civil war. *Popular Science Monthly*, 57, 1900, pp. 29-39.
- The future of the negro in the southern states. *Popular Science Monthly*, 57, 1900, pp. 147-156.
1901. Broad valleys of the Cordilleras. *Bull. Geol. Soc. Am.*, vol. 12, 1901, pp. 271-300.
- The Niagara Book. (W. D. Howells, Mark Twain, and N. S. Shaler.) New and revised edition, New York, 1901.
1902. Reports of Professor N. S. Shaler on the marshes and swamps of northern Long Island, between Port Washington and Cold Spring Harbor. North Shore Imp. Association reports on plans for the extermination of mosquitoes on the north shore of Long Island, etc. New York, 1902.
1903. A comparison of the features of the earth and the moon. Smithsonian Contributions, no. 1438, 1903, vol. 34, 79 pp., 25 pl.
- General description of the moon. Smithsonian Institution Annual Report, 1903, pp. 103-113. 1904.
- Elizabeth of England: a dramatic romance. Boston and New York.
1904. The citizen: A study of the individual and the government. New York, 1904.

The neighbor; the natural history of human contacts. Boston and New York.

1905. Man and the earth. New York, 1905.

After the memorials had been read the report of the Photograph Committee was deferred and the Secretary made sundry announcements regarding registration, luncheon, and division into sections for the reading of papers, on account of the long program. He stated that when the Society adjourned, adjournment would be to 9.30 o'clock Friday morning, at the American Museum of Natural History.

F. J. H. Merrill announced that the annual dinner of the Society would be held Friday, December 28, at 7.00 o'clock p m, at the "Arena" restaurant, 29 and 31 West Thirty-first street.

The Auditing Committee having raised the question as to the custody of the bonds belonging to the Society, the Acting President ruled, in the absence of Treasurer-elect Clark, that the securities were in the care of the Treasurer.

The Acting President then called for the reading of scientific papers. The first two papers on the printed program were read by title. They were

CUTTING OF THE MISSISSIPPI AND MISSOURI RIVER GORGES

BY N. M. FENNEMAN

LATERAL EROSION ON SOME MICHIGAN RIVERS

BY MARK S. W. JEFFERSON

This paper is printed as pages 333-350 of this volume.

GRADED SURFACES

BY F. P. GULLIVER

[Abstract]

This paper called attention to the need felt by the writer and others for a general term to use in physiographic descriptions for every "nearly level surface produced in the past or being now formed by some process of aggradation or degradation."

The writer showed that in a recent study of the complicated land forms of central Pennsylvania, as seen on a horseback trip with a pupil, no general term could be found to use for any given observed form without implying some hypothesis in regard to the origin of said land form. For purposes of reconnaissance study, there is a great need of some such general form term that will imply no genetic hypothesis in regard to its

formation. Then later, when the origin of the graded surface is determined, the appropriate genetic term may be applied.

The following terms were discussed, with references showing their uses by various writers: Terrace, grade, slope, bench, grade-level, grade-plain, peneplain, base-level, and bevel.

The conclusion was reached that some compromise must be made by physiographic writers in order to secure uniformity of usage. The choice would seem to lie between terrace, grade-level, grade-plain, and bevel. If one of these terms could be agreed upon by physiographic writers to use as a general form term, to imply no genetic hypothesis, greater clearness would result.

The next paper was read by title:

THE NEW MADRID EARTHQUAKE

BY M. L. FULLER AND E. M. SHEPARD

The following paper was then read:

PHYSIOGRAPHY OF THE LOWER HUDSON VALLEY

BY J. F. KEMP

The paper was discussed by Bailey Willis, W. M. Davis, and the author.

The next paper was

RELATIONS OF PHYSIOGRAPHY TO STRUCTURE AT MANHATTAN ISLAND AND VICINITY

BY ALEXIS A. JULIEN

The discussion was participated in by W. H. Hobbs, W. M. Davis, and the author.

The Society adjourned to Friday, as announced.

No session of the Society was held Thursday evening, out of deference to the American Association for the Advancement of Science, the retiring President of which, Professor C. M. Woodward, delivered the annual address at eight o'clock. This address was followed by a reception tendered by Columbia University to all the visiting scientists. The reception was followed by an informal smoker at the Faculty Club on the Columbia campus.

SESSION OF FRIDAY, DECEMBER 28

The Society met at 9.45 o'clock a m, in the west assembly hall of the American Museum of Natural History, Acting President Davis in the chair.

Announcements were made by the chair, by E. O. Hovey for the local committee, and by F. J. H. Merrill for the dinner committee.

The report of the Council was taken up and adopted without alteration.

The report of the Photograph Committee was read, as follows:

SEVENTEENTH ANNUAL REPORT OF THE COMMITTEE ON PHOTOGRAPHS

During the year 1906 there has been but little change in the collection of photographs belonging to the Society. No new views have been obtained, but, through the kindness of the Director of the Geological Survey, about 60 old prints have been replaced by new ones, which are printed in a superior manner and mounted on muslin. By this means the bulk of the collection has been considerably diminished.

The photographs are now stored in glass cases in my office, in the building of the Geological Survey, at Washington convenient for reference. Several members of the Society have obtained prints for use in reports and text-books and it is believed that there ought to be a very much wider use of the photographs for this purpose. It is expected that during the coming year a large number of new photographs will be added to the collection, selected from the numerous views which have been taken by members of the Geological Survey during the past few years. Contributions for the collection are desired, but care should be taken that they are views of general interest and illustrate geologic phenomena rather than scenery. A high technical standard is also requested.

Respectfully submitted.

N. H. DARTON,
Committee.

On motion, the report was adopted and the usual appropriation of \$15 for the use of the committee was voted.

RESOLUTIONS CONCERNING RETIRING SECRETARY AND TREASURER

John M. Clarke then presented resolutions regarding the Secretary and the Treasurer; which, on motion, were unanimously and most heartily adopted. The resolutions were as follows:

After sixteen years of service as Secretary of the Geological Society of

America, Professor H. L. Fairchild meets with us at this session for the last time in his official capacity.

These years of his service have witnessed the robust and vigorous growth of this Society. To his fidelity, enthusiasm, patience, conservatism, and lofty ideals we must ascribe in very large part the virile and promising condition of this Society today.

The Fellows of the Geological Society of America desire to enter on its permanent records this expression of appreciation of such devoted service and this acknowledgment of and gratitude for so willing a sacrifice on behalf of the progress of geological science in America.

Since 1891 Doctor I. C. White, now retiring from office, has served as Treasurer of this Society. For these sixteen years he has given to the management of its finances the benefit of his clear, practical judgment and his experience, and he has so carefully nursed its slender income that today the treasury of the Society presents a most substantial and very gratifying showing.

The Fellows of the Society desire to record an expression of their appreciation of this watchful, faithful, and profitable service.

On motion, the Society voted that after the reading of the next two papers there should be a division into two sections, in one of which should be read the papers on physical and structural topics and in the other the papers on economic, biographic, and glacial subjects.

The first paper of the day was

GEOLOGIC MAP OF NORTH AMERICA

BY BAILEY WILLIS

Remarks were made to the paper by W. M. Davis and W. H. Hobbs.

The next paper was

THE GEOLOGIC FOLIO

BY WILLIAM HERBERT HOBBS

The paper was discussed by C. D. Walcott.

The following paper was then read by title:

GEOLOGICAL MAP OF MASSACHUSETTS AND RHODE ISLAND

BY B. K. EMERSON

Although the paper was read by title only, the map was displayed at the meeting.

At this point in the program the Society separated into two sections. The first section, that including papers on physical and structural geology, remained in the west assembly hall and listened first to the paper,

CAVE SANDSTONE DEPOSITS OF THE SOUTHERN OZARKS

BY A. H. PURDUE

This paper has been published as pages 251-256 of this volume. The paper was discussed by A. W. Grabau, Bailey Willis, H. C. Hovey, W. M. Davis, M. L. Fuller, T. C. Hopkins, and the author.

Former President H. S. Williams was called to the chair.

The next paper was

CURRENT METHODS OF OBSERVING VOLCANIC ERUPTIONS

BY T. A. JAGGAR, JUNIOR

Remarks were made by H. F. Reid and Bailey Willis.

RESOLUTION CONCERNING INVESTIGATION OF VOLCANOES

The following resolution was then presented by T. A. Jaggar, Jr.:

WHEREAS Canada, the United States, and Mexico possess, in the Cordilleran belt, Alaska, the Philippines, the Hawaiian islands, Porto Rico, the Windward islands, and the Canal zone, a wide field for the investigation of earth movement and active volcanoes; and

WHEREAS geologic science needs permanent records, made in the field, of physical phenomena accompanying earthquakes and eruptions, both before and after the event; and

WHEREAS such records have direct bearing on prediction and on protection of life and property; therefore be it

Resolved, That the Geological Society of America strongly recommends to the several governments and to private enterprise the establishment of permanent volcano and earthquake observatories.

The resolution was referred to the Council for report the next day.

Acting President Davis resumed the chair.

The following paper was then presented:

EXPERIMENTS ILLUSTRATING EROSION AND SEDIMENTATION

BY T. A. JAGGAR, JR.

The last paper of the morning session of this section was

CHARACTERISTICS OF VARIOUS TYPES OF CONGLOMERATES

BY G. R. MANSFIELD*

* Introduced by W. M. Davis.

This paper was discussed by A. W. Grabau, J. Barrell, W. G. Miller, C. W. Brown, and W. M. Davis. The paper has been published in *The Journal of Geology*, volume xv, pages 550-555, September-October, 1907.

The second section met in the small assembly hall, on the third floor of the Museum building, and organized by electing A. P. Coleman chairman and H. M. Ami secretary.

The first paper read was

THE COBALT, ONTARIO, SILVER AREA

BY WILLET G. MILLER

The paper was discussed by S. F. Emmons, R. Bell, E. R. Buckley, and the author.

The next paper was read by title:

VIRGINIA BARITE DEPOSITS

BY THOMAS LEONARD WATSON

The following paper was read:

GEOLOGY OF SANTA BARBARA AND SUMMERLAND OIL FIELDS, CALIFORNIA

BY RALPH ARNOLD

Remarks were made by R. Bell, G. C. Martin, and the author.

The next paper read was

MEMORIAL OF A. R. C. SELWYN

BY H. M. AMI

[Abstract]

Selwyn was a geologist who attained distinction for his work in Great Britain, Australia, and British America. In 1845 he began his researches in England and Wales, in 1853 was appointed director of the Geological Survey of Victoria, Australia, and in 1869 succeeded Sir William E. Logan as director of the Geological Survey of Canada. Volcanic rocks and their relations to the earlier sedimentary formations formed the principal object of his energies in these three portions of the Empire, while he did much to emphasize the economic relations of these rocks to pure geological investigations. In Canada Selwyn wrought from December, 1869, to January, 1895, a period of twenty-five years. He traversed the continent before the transcontinental lines of railways were built, and directed the efforts of his staff to many of the portions of Canada whose resources to-day prove of such remarkable value. He received many distinctions at home and abroad.

Both sections adjourned for luncheon together, at the Manhattan Square hotel, and reconvened as a body at 2.30 o'clock p m.

Acting President Davis occupied the chair and gave the floor to W. G. Tight, who presented to the Society a most cordial invitation to hold its next meeting at Albuquerque, New Mexico. Two other equally cordial invitations were before the Council, but the Society by a large majority expressed its approval of holding such a western meeting and recommended the invitation to the Council for favorable consideration.

The Acting President then read the posthumous presidential address of Professor Israel C. Russell. This address is entitled

CONCENTRATION AS A GEOLOGICAL PRINCIPLE

and has been printed as pages 1-28 of this volume.

The Society then divided into two sections, as before, and proceeded with the reading of papers.

In the Physical and Structural section, under the chairmanship of the Acting President, the first paper read was

DOMe STRUCTURE IN CONGLOMERATE

BY RALPH ARNOLD

[Abstract]

In Eagle Rock valley, near Pasadena, California, are several dome-shaped structures, developed in lower Miocene conglomerate, of water-worn plutonic rocks. The largest dome is Eagle rock, about 80 feet high. This, in a general way, resembles the granite domes of the Sierras, but is much smaller. Scales, or plates, of the rock from a fraction of an inch to as much as 8 feet in thickness cover much of the sloping dome surface, while large quantities of broken scales from detrital heaps at the base of the rock. Not far distant from Eagle rock are other examples of the dome surface. In all cases the scales are not found on slopes of less than 30 or 40 degrees nor on the exposed southern faces or other portions of the rocks which are kept practically dry. The cracks under the scales always penetrate upward approximately parallel to the outer surface. The incipient cracks are closely followed by a zone of weathering which shows lamination parallel to the crack. Iron and other stains emanate from the cracks.

Two general theories have been advanced in explanation of such peculiar structure, all, heretofore, however, based upon observations of the development of domes in granite. According to one theory, the separation of the granite into curved plates is an original structure, antedating the sculpture of the country and determining the peculiarities of form. According to the other theory, the structure originated subsequently to the form, and was caused by some reaction from the surface. The conclusions reached by the writer regarding the origin of the conglomerate domes are as follows:

In the first place, they are locally hardened portions (gigantic concretions,

if you please) of a practically homogeneous conglomerate. After the ordinary process of erosion uncovers a sufficient surface of the indurated rock, the dome structures are formed by a successive scaling off of blocks, through the development of cracks approximately parallel to the steeply sloping surfaces. These cracks are probably due to expansive force developed by chemical reactions (weathering) produced largely by moisture, the moisture passing upward by capillarity through the incipient cracks caused by expansion, and thus advancing the process. The cracks originate in positions advantageous to the accumulation or retention of the moisture producing the weathering, such, for instance, as that occupied by the detrital material at the base of the slope or in the angle between overhanging blocks and the new dome surface. The direction of the cracks is determined by the configuration of the rock surface, being approximately parallel to it, the departures from strict parallelism being of such a nature as to omit angles and other features of irregularity. This parallelism to the surface is due to the expansive force acting along lines of least resistance, which in this case are practically normal to the outer rock surface. The slope of the surface is the governing function in the removal of the scales because the component of gravity tending to dislodge the separated scales is greater on steep slopes than on low, while the component of the same force tending to counteract the expansion due to weathering is correspondingly less on steep slopes, becoming greater as the angle of declivity lessens.

The paper was published in full in *The Journal of Geology*, volume xv, pages 560-570, September-October, 1907.

The next two papers were read without intermission and were discussed together. They were

RELATIONS BETWEEN CLIMATE AND TERRESTRIAL DEPOSITS

BY JOSEPH BARRELL

[Abstract]*

Contents

	Page
General introduction	616
Part I. Relation of sediments to regions of erosion.....	618
Part II. Relation of sediments to regions of deposition.....	619
Part III. Relations of climate to fluvial transport.....	620
Summary and conclusion.....	620

GENERAL INTRODUCTION

The environment of the lands may be classified into three fundamental and independent factors—the relations to the surrounding seas, the topography which forms their surfaces, and the climates which envelop them—each of major importance in controlling the character of the lands. . . . The third great problem of terrestrial environment, the succession of ancient climates, lags behind the other two in development, but is no less important in a com-

* To be published in full in *Journal of Geology*, vol. xvi, 1908, beginning with no. 2.

plete understanding of the history of the earth and its inhabitants. This backwardness is doubtless due to the intangible nature of climate and the lack of direct record of its geologic changes. When it is considered, however, how fundamental are the relations of continental deposits to the climates in which they are formed, it is seen that the record of geologic climates, while indirect and to a considerable extent awaiting interpretation, is nevertheless in existence. This is exclusive of the significance of salt and gypsum deposits on the one hand or of glacial deposits on the other, which are of course universally recognized, but these are the marks of climatic extremes.

The causes of climatic variations as distinct from the stratigraphic record of climates have in recent years received a large amount of attention, with the result that a considerable body of knowledge has taken the place of previous speculation. But what are the geologic records of that great variety of climates which, excluding the desert belts of the world, reach at present from the equator to the frigid zones? They may be most favorably studied in ancient continental deposits, since these are free from the contributory record of the sea; and, as it is the average climates which it is sought to investigate, it is especially in ancient fluviatile deposits of continental nature rather than in deposits of desert or glacial origin that the record is to be found. In such river deposits each stratum represents an old land surface, the seat of abundant animal and vegetable life, sealed and protected by the laying down of the succeeding strata instead of destroyed by erosion, the dominant process acting on the land surface. In such deposits the evidence most usually studied is that from the teeth and feet of animal fossils, or the nature of vegetable remains. But many continental deposits are without fossils and many groups of organisms show a wide climatic range; so that it is very desirable that other features constantly present, such as the chemical, textural, and structural characteristics of the strata, should be available for the climatic interpretation. The significance of such features has of course not entirely escaped attention; the presence of red in shales or sandstones is sometimes cited as evidence of derivation from a deeply decayed and highly oxidized regolith, or the existence of a conglomerate whose pebbles are of vein quartz as evidence of the thorough decomposition of an ancient soil; but such expressions have been made without a preliminary investigation into all the possible modes of origin, and it will be found that other interpretations are also possible.

The problem, then, in the present paper is to separate the influence of the topographic from the climatic factors in the making of terrestrial sediments. In seeking for data to draw the lines more closely, it is found that the relations of climate to the nature of the land waste and river sediments has attracted the attention of relatively few explorers and scientists. Such exceptions must, however, be noted as Blanford and Oldham, Walther, Hilgard, Merrill, Russell, Davis, and Huntington. These men, with a few others, have largely supplied the data which make the discussion possible.

The first part, dealing with the relations of climate to erosion, is necessarily largely physiographic, and, in the case of material which is carried long distances by rivers, these relations are of less final influence than the conditions under which the sediments are deposited and those of the preceding transportation. The purpose of the whole paper is, however, not physiographic, but

stratigraphic. For this reason it is the relation of physiography to erosion and consequent supply of sediments which is dwelt upon rather than the discussion of the land forms as an end of investigation. Owing to this somewhat unusual use of physiography as well as the desire to make the discussion more complete for students in other branches of geology, it is necessary to go over some ground which is familiar to physiographers. Even from a physiological standpoint, however, it is thought that a brief general discussion of all the climatic factors influencing erosion is not without its value, since it is found to suggest some new points of view upon several old problems. For the elaboration of the second and third parts, those upon the relations of climate to fluvial deposition and the preceding transportation, sufficient reason is found in the existence of the diverse views held by many working geologists upon the significance of various stratigraphic characters, and, furthermore, in the general conclusion of the present paper, that climate is a factor comparable to disturbances of the crust or movements of the shoreline in determining the nature and the variations in the stratified rocks of continental or offshore origin, thus playing a part of large, though but little appreciated, importance in the making of the stratigraphic record.

The investigation was instituted to see to what extent profound climatic variations, complicated doubtless with some tectonic movements, could account for the great contrasts in the Lower and Upper Carboniferous formations of eastern Pennsylvania—formations which had been found by the writer from observations in the field to be continental in nature, and therefore their contrasted features not to be attributed to changes in the relations of land and sea.

An abstract of the body of the paper is as follows:

PART I. RELATION OF SEDIMENTS TO REGIONS OF EROSION

Observations of the nature of river sediment show that it is only in local deposits that the character is markedly due to the kind of rocks undergoing erosion, so that in regions where the sediment has no relation to the adjacent formations but little difficulty will be found in allowing for the more or less unknown lithological nature of the source.

The interrelations of topography and climate to erosion are more complicated, but it is shown from present instances that even where land relief leads to vigorous erosion the influence of arid or humid climates is able to make itself felt in the ratio of coarse to fine material and the freshness or degree of decomposition of the fine. In the mature stage of the topographic cycle these differences become accentuated. The influence of old topography is of less importance, since under such conditions fluvial continental deposits are reduced to a minimum. Between the climatic extremes, however, the results, as are well known, are still conspicuous, leached clays being the mark of rainy and loess and dune sand of arid climates.

Under the relations of temperature to erosion, the influences of vegetation, of frost action, of snow action, and of insolation are discussed. It is noted, as Chamberlin and Salisbury have pointed out, that *increased cold* tends to weaken the power which vegetation possesses in producing decay but preventing erosion. *Increased frost action* is found to greatly increase the rate of

disintegration in young or mature topography without correspondingly increasing the powers of waste-removal. The Gila conglomerate of Arizona is discussed as an example of a great conglomerate formation resulting primarily from a climatic change in which the ratio of erosion to transportation had been increased through increased frost action at times of glaciation in other regions. *Increased snowfall* works in an opposite direction, by protecting the surface from disintegration and promoting transportation.

Under the effects of climatic changes resulting in an *increase of temperature*, it is pointed out that in regions of topographic relief insolation is a weaker force than frost action. In all geologic times, irrespective of climatic variations, it has worked most strongly in the torrid and has been practically absent from the polar zones. In intensity it must have varied considerably, owing largely to geographic as well as possibly to solar conditions. In regions of humid climate rock decay takes the place of insolation and also operates most strongly in the torrid zone; but on account of the mat of vegetation it is not known if erosion is weaker or more rapid than in a temperate climate.

In conclusion, it is pointed out that an examination of the *matrix* of fluvialite conglomerates is of importance for separating the climatic from the topographic conditions of origin, while the *simultaneous development* of conglomerates within many hydrographic basins, as Davis has shown, may point to a climatic as against a tectonic origin. Finally the relative rates of erosion under various climates are discussed and it is concluded that the development of plains of marine denudation depends largely on the climatic factor, taking place most readily in an arid climate. As a result of the discussion of the first part, it is concluded that climatic variation is a major factor, comparing with subordinate tectonic movements, in governing both the kind and quantity of the material supplied to the rivers.

PART II. RELATION OF SEDIMENTS TO REGIONS OF DEPOSITION

In the second part, the influence of topography is first considered, the piedmont slopes being notably better drained than deltas and only covered with carbonaceous deposits in a cool and rainy climate. Over the delta there is a progressive flattening of the grade with the production of swamp areas, to some extent even in desert climates. A number of factors independent of the climate, such as the building of two confluent deltas or a slowly rising water level, tend to increase the area of swamp; but in desert climates there is a close association of the swamp deposits with sharply contrasted evaporation and æolian products. Under all climates the amount of swamp area reaches a maximum on the seaward margin of the delta. Following this preliminary discussion, the effects of five kinds of climates are considered: constantly rainy, arctic, intermittently rainy, semiarid, and arid. Each impresses well marked distinctions upon the flood-plain deposits. The constantly rainy gives rise to carbonaceous and well leached clays and sands, besides beds of peat; the arctic climate differs in the less leached clays, containing especially more potash; in intermittently rainy climates the carbonaceous deposits of the flood-plains are local, the materials are oxidized during seasons of dryness, the clays are slightly calcareous, and fossils are apt to be scanty. It is concluded that such deposits occur to an unappreciated extent in

the geologic record and are apt to be classed with marginal marine formations on account of their lack of definite characteristics. In semiarid climates the deposits show not only a higher oxidation, but the clays contain more potash and are markedly calcareous. As swamp deposits diminish, mud-cracked strata increase and may become noteworthy. Under arid climates potash-bearing and calcareous clays and mud-cracked strata become more prominent; aeolian and especially evaporation deposits of gypsum and salt also occur; oxidation of the delta materials is almost universal, with the result that dark or variegated shales are nearly absent. A discussion of the climatic significance of color follows, and it is concluded that red is the common color of consolidated fluvial deposits which were subjected at the time of their accumulation to seasons of subaerial oxidation, and that red shales and sandstones may therefore arise from either the yellow or red alluvium of intermittently rainy, semiarid, or arid climates. In conclusion it is pointed out that while the varying powers of erosion and transportation are delicate stratigraphic indicators of *climatic fluctuations*, the chemical and organic control accompanying the deposition are the more secure indicators of the *average climatic conditions*.

PART III. RELATIONS OF CLIMATE TO FLUVIAL TRANSPORTATION

This subject is considered last, since except in connection with the conditions of erosion and deposition the climatic relations of transportation are not conspicuous. Laboratory experiments and observations of river action lead to the induction that one of the reasons why rivers carry coarse material such short distances is because of the mechanical weakness of all but quartz and quartzite pebbles and the consequent loading up of the current with finer material. The slope of a graded river is in delicate adjustment between the amount of water and the amount of the load, and consequently every change in these factors throws it out of equilibrium. As a result, sometimes cutting and sometimes depositing is in excess throughout the middle portion of its course. In the case of long rivers flowing from regions in topographic youth or maturity, as those flowing east from the Andes and Rockies, a very great difference in grade exists, according as the river course lies in a semiarid or rainy climate. Under semiarid climates piedmont slopes may be built up to a thickness of a thousand feet or more, and in a climatic variation to one of pluvial nature this may be rapidly carried away, and, since it consists largely of sand and gravel, a great sand or conglomerate formation will be spread over the delta plain. The dissection in the early Pleistocene or latest Tertiary of the high plains fronting the Rocky mountains is cited as an instance. On the other hand, upon a change of climate from humid to semiarid, another sand and conglomerate formation will be laid down close to the mountains as a piedmont slope, building up the river channel to a steeper grade. From this it is concluded that the region of deposit of river gravels and its shiftings is of great climatic significance.

SUMMARY AND CONCLUSION

Summing up the preceding discussion, it is to be concluded that conglomerate and sandstone formations intercalated between others of different nature may be due to three distinct causes:

First. Marine conglomerates and sandstones, due to marine planation and transportation, enabled to reach wide horizontal extent over shallow seas through crustal movements shifting the zone of wave and current action.

Second. Tectonic, conglomerates and sandstones, due to subaerial erosion, owing to a steepening of the river slopes, either from mountain making, crustal warping, or subsidence of the ocean level.

Third. Climatic conglomerates and sandstones, due to climatic change, without necessarily any new and accompanying crustal movement, producing a shifting of the region of accumulation of gravel or sand and a resulting contrast in coarseness, in color, and in chemical composition with the underlying and overlying formations. It is by means of the associated features that climatic conglomerates are to be distinguished from those of tectonic origin. For these two to be sharply separable in nature, the climate must remain stable during an epoch of crustal movement and, *vice versa*, earth movements must be supposed quiescent during a time of climatic change. As tectonic and climatic movements may be intermixed, there may often be a dual cause of conglomeratic formations.

The establishing of a class of climatic conglomerates involves the conception of the shifting location of deposits (*in space*) during an epoch of crustal quiet as being of the same geologic importance as the varying pulses of erosion (*through time*) due to tectonic movements.

In conclusion, it is seen that by means of conglomerate or sandstone formations the important changes are recorded in each of the three fundamental environments of the lands, namely, the relations to the surrounding seas, the topography which forms their surfaces, and the climates which envelop them.

ORIGIN AND SIGNIFICANCE OF THE MAUCH CHUNK SHALE

BY JOSEPH BARRELL

This paper is printed as pages 449-476 of this volume.

The two foregoing papers were discussed by D. White, B. Willis, E. Huntington, A. W. Grabau, W. M. Davis, and the author.

The next paper was

RIVER SEDIMENT AS A FACTOR IN APPLIED GEOLOGY

BY W J MC GEE

The last paper of the afternoon in this section was

ORIGIN OF OCEAN BASINS IN THE LIGHT OF THE NEW SEISMOLOGY

BY WILLIAM HERBERT HOBBS

This paper forms pages 233-250 of this volume.

The second section met with George P. Merrill in the chair and H. M. Ami acting as secretary.

The first paper on the program was read by title:

PERSONAL REMINISCENCES OF SIR WILLIAM E. LOGAN

BY ROBERT BELL

[Abstract]

In his day Logan was one of the most prominent geologists, not only of America, but of the world, being well known in Europe. In the present strenuous times, new men come rapidly upon the stage, and although this great geologist has been dead only about thirty years, his name is not familiar to the younger generation. While the above mentioned paper deals largely with instructive personal incidents and amusing anecdotes, it serves to illustrate the evolution of the geology of eastern North America and the growth of the Geological Survey of Canada.

Logan having pursued scrupulously honorable methods in scientific matters as in everything else, his example was an inspiration to all who were associated with him. His desire to give credit to his assistants for their share in the work secured for him enthusiastic followers, among whom were Murray, Hunt, Billings, and others. His painstaking methods and wonderful powers of observation and reasoning show, in the light of our present knowledge, his marvelous insight as to the relations of the various series of rocks to one another, and his classification and nomenclature of sixty-odd years ago, when geology itself was in its infancy, were adopted two years ago as the best possible, by the International Committee on the Lake Superior rocks.

Logan's life and times are so full of interest to all geologists that it is believed this short sketch at first hand by one who was so intimately associated with him is worth preserving.

This was followed by

A LOWER HURONIAN ICE AGE

BY A. P. COLEMAN

The discussion which followed was participated in by W. G. Miller, R. D. Salisbury, A. C. Lane, R. Bell, F. G. Clapp, and the author. The paper has been published in full in the *American Journal of Science*, volume xxiii, 1907, pages 187-192.

The next two papers were read by title:

GLACIATION OF MANHATTAN ISLAND, NEW YORK

BY ALEXIS A. JULIEN

GLACIAL EROSION IN THE NORTHFIORD

BY MARK S. W. JEFFERSON

This paper has been printed as pages 413-426 of this volume.

The next paper read was

RECENT CHANGES IN THE GLACIERS OF GLACIER BAY, ALASKA

BY F. E. AND C. W. WRIGHT

The paper was discussed by W. Blake (a visitor), H. F. Reid, F. P. Gulliver, G. F. Wright, and F. E. Wright.

The last paper of the session was

RECENT CHANGES IN THE MALASPINA AND OTHER GLACIERS OF THE YAKUTAT BAY REGION, ALASKA

BY RALPH S. TARR

Remarks were made by T. A. Jaggar, Jr., H. F. Reid, A. H. Brooks, and the author. The paper forms pages 257-286 of this volume.

The evening was devoted to the annual dinner of the Society, which was the largest that has been held, 138 Fellows and guests participating. W. M. Davis presided.

SESSION OF SATURDAY, DECEMBER 29

The Society met in general session at 9.45 o'clock a m, with Acting President Davis in the chair.

REPORT OF THE AUDITING COMMITTEE

The Auditing Committee reported that the accounts of the Treasurer had been found correctly cast and properly vouched.

On motion, the report was accepted and ordered placed on file.

The Secretary reported that the Council had voted that the next meeting of the Society be held at Albuquerque, New Mexico, December 30, 1907, to January 1, 1908, and that the resolutions offered by T. A. Jaggar, Jr., had been referred to a committee, consisting of T. A. Jaggar, Jr., J. F. Kemp, and E. O. Hovey, for consideration and report at the next annual meeting of the Society.

After some other announcements the Society divided into two sections, as before, and the scientific program was taken up. The Physical and Structural section remained in the west assembly hall, with Acting President Davis in the chair.

The first paper read was

HYPOTHESIS OF CONTINENTAL STRUCTURE

BY BAILEY WILLIS

Discussion was offered by C. Schuchert, A. Heilprin, B. K. Emerson, F. E. Wright, and the author. The paper appears as pages 389–412 of this volume.

The next paper was read by title:

THE LIMELESS OCEAN OF PRE-CAMBRIAN TIME

BY REGINALD A. DALY

This paper has been published in full in the American Journal of Science, volume xxiii, 1907, pages 93–115.

The following paper was then read:

PERMO-CARBONIFEROUS CLIMATIC CHANGES IN SOUTH AMERICA

BY DAVID WHITE

[Abstract]

The fossil plants collected by the Coal Commission of Brazil not only confirm the existence of the Lower Gondwana flora in Brazil, but they offer striking evidence of climatic changes. At the same time they afford new data relating to Permo-Carboniferous southern land connections and the distribution of Permian glaciation.

The stratigraphy of the Brazilian coal measures is admirably worked out by Dr I. C. White, late chief of the commission, in his paper "Coal Measures and higher beds of South Brazil," which he summarized at this meeting of the Society. For the descriptive and systematic discussions of the fossil plants, embracing 140 species, the reader is referred to the official Brazilian report.*

The fossil plants in the series immediately above the basal conglomerates, which repose unconformably on the ancient eruptives, comprise a typical Lower Gondwana flora, including its principal and characteristic species as well as genera. This flora is so far identical with that associated with the glacial boulder-bearing sediments in Australia and overlying the glaciated conglomerates in India and South Africa as to leave no room for doubt as to the contemporaneous existence of a similar cold climate in South America. The consequent inference as to glaciation in that continent is lithologically proven according to Doctor White, who confirms the observations and tentative views offered in 1889 by Derby.

The beds 130 meters above the base of the Tubarão series show numerous megaspores apparently of Sigillarian origin. The evidence of the return of representatives of the northern or cosmopolitan flora is more abundant in the region of 175 meters above the base, where we find *Sigillaria brardii*,* *S.*

* Commissao dos Estudos das Minas de Carvao de Pedro do Brasil. Rio de Janeiro, 1907.

australis. *Lepidodendron pedroanum*,* and *Lepidophloios laricinus* again assuming the rôle of coal-makers. Mingled with them are species of Gangamopteris, Glossopteris, and Phyllothea, of the Gangamopteris flora. It is plain, therefore, that an amelioration of climate sufficient to permit the return of some of the northern types had already taken place; and it is equally evident that the absence of these types during the deposition of the lower beds was due to climatic inhospitality rather than to a marine or other barrier.

The paleobotanical material from the region of the Iraty black shale and the succeeding beds of the Passa Dois series is confined to fossil stems. This is the zone of the *Psaronius brasiliensis* described by Brongniart, and of the *Lycopodiopsis derbyi* and the *Dadoxylon pedroi* respectively described by Renault and Zeiller, as well as of the *Stereosternum tumidum* Cope and *Mesosaurus brasiliensis* McGregor. The Passa Dois series has also furnished another large Lycopodialean tree, probably belonging to Sigillaria, and two new gymnosperms referred to Dadoxylon. The presence of Psaronius would seem to indicate the return of Pecopterids, presumably of the Cladophleboïd group, to the southern Brazilian basins. This indication of further moderation of the climate is corroborated by the woody structure of the fossil trees, in which there are extremely obscure, if any, traces of annual rings, the growth of the trees being apparently uninterrupted by seasonal changes. It is therefore permissible to conclude that at this time a mild and equable climate once more prevailed, so that the surviving elements of the Permian cosmopolitan flora were able freely to migrate and mingle with the survivors of the Gangamopteris flora.

The date of South American glaciation, which on the paleobotanical evidence appears to have been contemporaneous with the Dwyka in South Africa, the Talchir of India, and possibly the later Newcastle ice in Australia, would seem to be at or extremely near the close of the Carboniferous. That the severe cold was of short duration is shown by the early return of some of the exiled northern Permo-Carboniferous plants. The fact that the post-glacial moderation of the climate occurred some time before the extinction of Lepidophloios and Lepidodendron conclusively shows that the climatic refrigeration could not have occurred later than the very early Permian. It is interesting to note that the types first to return, and therefore presumably the hardiest, belong to the Lycopods.

The distribution in India, Australia, South Africa, and South America of a terrestrial flora of varied elements like the older Gondwana flora necessitates a distribution of the land masses in such a way as to permit land plant migration with absolute freedom between these regions. The testimony of vertebrates and plants is mutually corroborative on this point. The author inclines to recognize the important agency of an Antarctic land mass or continent, of which southern South America or South Africa may have been mere lobes, in facilitating this free intermigration.

The abstraction of carbon dioxide from air and sea and the storage of the same in the coals, carbonaceous shales, and limestones of the Upper Carboniferous has never been equaled in any other period of geological history.

* The employment of capital initials for names of species directly derived from names of persons is not permitted by the rules for printing early adopted by this Society.

The hypothesis of Arrhenius and Chamberlin would therefore appear to find circumstantial support in the occurrence of glaciation at the time of the post-Pennsylvanian uplift. If the efficiency of CO₂ as a climatic factor be admitted, not only may the occurrence of refrigeration itself be explained, but perhaps the singular geographical distribution of the ice action as well. The latter would appear to have been the result of complicated influences, very important among which would be the great exaggeration of temperature differences attending differences in altitude in the atmospheric column were there a depletion of the CO₂ of that column. The temperature gradient would be very much steepened and the frost line brought down nearer to sealevel. To this fact, which does not appear to have received due consideration at the hands of geologists, in connection with the probable great size and considerable height of the older Gondwana land masses, as shown by the great thickness of coarse sediments and the great size of the continental basins, is, in the writer's opinion, chiefly due the extension of the Permian glaciation within the tropics.

The next paper was

COAL MEASURES AND HIGHER BEDS OF SOUTH BRAZIL

BY I. C. WHITE

The next paper was read by title:

NORMAL PRESSURE FAULTING IN THE ALLEGHENY PLATEAUS

BY GEO. H. ASHLEY

The following two papers were read without intermission:

CONDITIONS OF CIRCULATION AT THE SEA MILLS OF CEPHALONIA

BY M. L. FULLER

This paper has been printed as pages 221-232 of this volume.

Mr Fuller's paper was discussed by J. F. Kemp and W. M. Davis.

CONTROLLING FACTORS OF ARTESIAN FLOWS

BY MYRON L. FULLER

[Abstract]*

Contents

	Page
Introduction.....	627
Underground water reservoirs.....	627
Types of water reservoirs.....	627
Sources of underground water.....	629
Confining agents	629
Nature of artesian circulation.....	630
Factors in artesian circulation.....	630

* The full paper is in press as a bulletin of the United States Geological Survey.

	Page
Requisites of artesian flows.....	631
Objections to common requisites.....	631
Chamberlin's requisite conditions of artesian flows.....	631
Pervious stratum	632
Impervious bed below.....	632
Impervious bed above.....	632
Inclination of beds.....	632
Outcrop of porous stratum.....	633
Adequate rainfall	633
Points of escape.....	633
Essentials of artesian flows.....	633
Modifying factors	634
Secondary factors of artesian flows.....	634

INTRODUCTION

It is now more than twenty years since the publication of the admirable paper by Professor T. C. Chamberlin on the "Requisite and qualifying conditions of artesian wells,"† during which period it has remained the standard in this country. It deals, however, almost entirely with the porous bedded rocks, such as sandstones, taking little account of the drift or limestones, and practically none at all of the crystalline rocks. Late investigations, especially in the East, and the growing demand for ground water for industrial and public purposes or for summer resort supplies have led to a very extensive development of wells in each of the classes of material mentioned, and have shown their great economic as well as scientific importance, emphasizing the need of a reexamination into the requisites of artesian flow and a discussion of their controlling and modifying factors.

In view of the number of different ways in which the term artesian has been used in the past, it will not be out of place to state that in the present paper the term is applied in the sense adopted by the United States Geological Survey after consultation with leading geologists of the country, namely, to designate the hydrostatic principle by which confined waters tend to rise in virtue of the pressure of the overlying water column, irrespective of whether or not this pressure is sufficient to lift the water to the surface and produce a flow.*

Artesian flows may be said to depend upon the nature of the reservoirs, the conditions of confinement, and the sources of the water. An enumeration of the factors are given below.

UNDERGROUND WATER RESERVOIRS

By an underground water reservoir is meant the opening or system of openings within the rocks in which the water is contained. The more important types may be summarized as follows:

TYPES OF WATER RESERVOIRS

I. Original forms.

1. Original pores.

† Fifth Annual Report U. S. Geological Survey, 1885, pp. 125-173.

* M. L. Fuller: Significance of the term "Artesian." Water supply and irrigation paper no. 160, 1906, pp. 9-15.

2. Lamination planes.
 3. Bedding planes.
 4. Vesicles (in igneous rocks only).
- II. Secondary forms.
1. Secondary pores.
 - A. Pores resulting from leaching and solution.
 - B. Pores resulting from recrystallization.
 2. Solution openings.
 - A. Isolated cavities.
 - B. Tubular channels.
 - C. Sheet openings.
 3. Mechanical eroded reservoirs.
 - A. Tubular channels.
 - B. Pocket openings.
 - C. Sheet openings.
 4. Fracture openings.
 - A. Irregular openings.
 - a. Desiccation cracks.
 - b. Contraction fissures.
 - c. Tortion fractures.
 - d. Shearing breaks.
 - e. Vibration fractures.
 - f. Explosion ruptures.
 - B. Joints.
 - a. Vertical joints.
 - b. Horizontal joints.
 - c. Parallel joints.
 - d. Intersecting joints.
 - e. Joint breccias.
 - C. Faults.
 - a. Single fault planes.
 - b. Parallel fault planes.
 - c. Irregular faults.
 - d. Intersecting faults.
 - e. Fault breccias.
 5. Vein contacts.
 6. Igneous contacts.
 7. Shearing planes.
 8. Cleavage planes.
 9. Foliation and schistosity planes.

Most of the forms enumerated are not limited to any particular class of rock, but may be found in the stratified, metamorphic, and igneous types. The vesicles of igneous rocks, however, have no exact counterpart in stratified rocks, although the pores containing included water of sedimentation approach them in nature. Solution channels are likewise usually, although not necessarily, found only in sedimentary rocks, foliation and schistosity are mainly features of igneous or metamorphic rocks, while lamination, as the term is here used, is a feature of stratified rocks.

SOURCES OF UNDERGROUND WATER

The probable source of underground waters has been widely discussed, and while every one would doubtless agree that by far the greater part is derived from rainfall, there is a considerable variation of opinion as to the relative importance of the other sources, and especially as to the part played by the sea and aqueous magmatic emanations. The common sources of underground waters are as follows:

List of Sources of Underground Water

- I. Atmosphere.
 - A. Direct.
 - a. Precipitation.
 - b. Condensation.
 - B. Indirect.
 - a. Lakes.
 - b. Streams.
- II. Hydrosphere, or ocean.
 - A. Recently or contemporaneously absorbed waters.
 - B. Originally included sea water.
- III. Lithosphere, or crust.
 - A. Primary waters.
 - Chemically excluded waters.
 - B. Secondary waters.
 - Physically excluded waters.
- IV. Centrosphere, or interior.
 - A. Directly excluded waters.
 - B. Indirectly excluded magmatic waters.

CONFINING AGENTS

In the discussion of the confining agents in artesian systems the materials are divided for convenience into bedded and jointed rocks. In the usual discussion the impervious bed is ordinarily the only agent recognized, but the extensive fieldwork which has been conducted by a considerable body of workers for the last three years has brought to light some new agents and emphasized the importance of other seldom mentioned factors.

List of Confining Agents

- I. Bedded rocks.
 - A. Upper confining agents.
 - 1. Impervious beds.
 - 2. Stratification.
 - 3. Friction.
 - 4. Mineral crusts.
 - 5. Frost zones.

6. Confined air and gas.
7. Fresh water.
8. Salt water.
- B. Lower confining agents.
 1. Impervious beds.
 2. Stratification.
 3. Friction.
 4. Mineral crusts.
 5. Frost zones.
 6. Confined air and gas.
 7. Fresh water.
 8. Salt water.
 9. Cementation.
 10. Heat.
 11. Pressure.
- II. Jointed and fractured rocks.
 - A. Upper confining agents.
 1. Impervious hanging wall.
 2. Impervious surface coverings.
 3. Frost fillings.
 4. Vein fillings.
 5. Weathering products.
 6. Converging walls.
 7. Interrupted joints.
 8. Sea water.
 - B. Lower confining agents.
 1. Impervious footwalls.
 2. Vein fillings and cementation.
 3. Converging walls.
 4. Interrupted joints.
 5. Fresh water.
 6. Sea water.
 7. Heat.
 8. Pressure.

NATURE OF ARTESIAN CIRCULATION

Artesian circulation takes place by virtue of the variations of pressure to which the water is subjected in different parts of its reservoirs. By definition of the artesian principle, all but hydrostatic pressure is excluded as a determining cause of circulation. There are, nevertheless, a number of modifying agencies which in cases exert a powerful influence on the water-head and movements. The more important of the controlling factors may be summarized as follows:

FACTORS IN ARTESIAN CIRCULATION

Primary factors.

Gravity (hydrostatic pressure).

Modifying factors.**Factors affecting pressure.**

Barometric variations.

Temperature variations.

Density of waters.

Variations due to temperature.

Variations due to dissolved salts.

Height of adjacent water levels.

Water table.

Neighboring water bodies and supply.

Tides.

Winds.

Rock pressure.

Factors affecting movement.

Porosity.

Size of grain openings.

Temperature.

REQUISITES OF ARTESIAN FLOWS

Having outlined the more important conditions bearing on the occurrence and movements of underground waters, the controlling factors of artesian flow may be considered. In doing this it is necessary to point out certain objections to some of the commonly postulated requisites.

OBJECTIONS TO COMMON REQUISITES**CHAMBERLIN'S REQUISITE CONDITIONS OF ARTESIAN FLOWS**

Since the appearance in 1885 of the paper of T. C. Chamberlin on the "Requisite and qualifying conditions of artesian wells," a single set of requisites has, with few exceptions, been followed by writers of underground water papers. These in brief are: (1) A pervious stratum, to permit the entrance and the passage of water; (2) a water-tight bed below, to prevent the escape of water downward; (3) a like impervious bed above, to prevent the escape upward, for the water, being under pressure from the fountain head, would otherwise find relief in that direction; (4) an inclination of these beds, so that the edge at which the waters enter will be higher than the surface at the well; (5) a suitable exposure of the edge of the porous stratum, so that it may take in a sufficient supply of water; (6) an adequate rainfall, to furnish this supply; and (7) an absence of any escape for the water at a lower level than the surface at the well.

There is one very serious objection to the requisites outlined above, namely, they apply only to a single class of flows from stratified rocks, neglecting not only flows from other varieties of rock, but even other types of flow from the same rocks. There are, moreover, many exceptions to the postulated requisites, which, taken in connection with the limitations mentioned, make new and more comprehensive definitions desirable. Some of the objections to the requisites are mentioned below.

* Fifth Annual Report U. S. Geological Survey, pp. 125-173.

PERVIOUS STRATUM

A pervious stratum, although a common form of reservoir, is seldom essential to artesian flows. In addition to the porous stratum postulated by the first requisite, flows may be obtained from lamination, bedding, cleavage, and shearing planes, from solution passages and mechanically eroded reservoirs, from vesicular zones in igneous rocks, from irregular, joint, and fault fractures, and from vein and igneous contacts. Metamorphic and igneous as well as stratified rocks not only may, but actually do, yield flows in a large number of instances; in many others the water falls only a few feet short of the surface, while in practically all the wells the waters rise materially when encountered—that is, are truly artesian.

With the exception of the vesicular lavas, the sources of water mentioned are not in the nature of beds, but are in the form of actual openings—a type of passage not recognized among the older requisites.

IMPERVIOUS BED BELOW

From the nature of the passages enumerated above and given in more detail in the list on page 628 it is apparent that the second requisite—the impervious underlying stratum—loses much of its force. Such impervious beds are an adjunct to many flows in stratified rocks, but numerous other agents may serve the same purpose. In bedded rocks the following may be mentioned: Stratification, friction, mineral crusts, frost zones, confined air and gas, fresh or salt water, cementation, heat and pressure. In jointed and fractured rocks impervious footwalls, vein fillings, converging walls, interrupted joints, fresh and salt water, heat and pressure, are the most important.

IMPERVIOUS BED ABOVE

The objections to the postulation of an upper impervious bed are similar to those of the lower confining bed just enumerated. In the case of the bedded rocks the following, in addition to the postulated impervious stratum, may serve as confining agents: Stratification, friction, mineral crusts, frost zones, confined air and gas, fresh water and sea water. In the case of jointed and fractured rocks the more common additional agents are impervious hanging wall, impervious surface coverings, frost and vein fillings, weathering products, converging walls, interrupted joints, and sea water.

INCLINATION OF BEDS

Inclination of the water-bearing bed, while a common factor in artesian flow, is by no means essential. Water appears to penetrate in many cases into lenses of sandstone in rocks like those of the Carboniferous of Pennsylvania through joints and similar openings, and flows are obtained independent of any inclination of the bed affording the water. The same is true in the case of the horizontal beds yielding flows by virtue of the opposition of the stratification planes. Joint and solution passages also afford artesian flows independent of any inclination at the point penetrated.

In both bedded and crystalline rocks the pressure must, of course, be transmitted from connecting passages or other water reservoirs at higher levels, but

the supply itself does not necessarily come from a higher level, since besides the downward moving meteoric water supplies may be furnished by the sea, by waters chemically or physically excluded from the crust, or by direct or magmatic exclusion from the centrosphere.

OUTCROP OF POROUS STRATUM

The postulated suitable exposure of the edge of the porous stratum so that it may take in a sufficient supply of water, though a common, is far from an essential factor of artesian flows. Some of the horizontal sandy beds from which the flows are obtained in Long island and Michigan never outcrop, the water penetrating directly downward through the overlying layers. Moreover, throughout extensive areas of the Silurian, Devonian, and Carboniferous rocks in Pennsylvania, West Virginia, and Ohio, in the areas underlain by Cretaceous beds in the Fort Monroe district of Virginia, and the Wilmington region of North Carolina, and in many lesser areas elsewhere, the deep artesian waters represent originally included sea waters and not waters entering through the outcrop. Other sources, including waters chemically or physically excluded from the crust, or by direct or magmatic exclusion from the centrosphere, may furnish artesian supplies independent of the condition of outcrop.

ADEQUATE RAINFALL

From the fact that water may be, and in the case of the salt waters of the Carboniferous and Coastal Plain rocks just described, as well as in the case of magmatic waters actually is, derived over extensive areas from sources other than rainfall, it is clear that the latter should not be included as an absolute requisite.

POINTS OF ESCAPE

There are very few artesian systems in which there is not more or less leakage. In the thicker and more persistent beds the leakage is often sufficient to insure circulation for long distances from the outcrop. Thus, in the Cretaceous beds beneath Charleston, South Carolina, and Savannah, Georgia, fresh water has replaced the salt at least as far as the seacoast, or a distance of over 100 miles from their outcrop. At Fort Monroe, Virginia, and Wilmington, North Carolina, on the other hand, there appears to be but little leakage and the fresh water circulation reaches to a much less distance, only salt water being obtained at the localities mentioned. The absence of leakage appears to have determined the presence of salt waters in the oil-bearing rocks of Pennsylvania and elsewhere.

In order that leakage may prevent flows, it must take place near the point at which the water horizon is tapped. It has been shown many times by Chamberlin and others that its influence is limited. The requisites should therefore postulate the absence of near-by leakage rather than the non-occurrence of leakage.

ESSENTIALS OF ARTESIAN FLOWS

The essentials of artesian flows, as recognized by the writer, are as follows:

Essentials of Artesian Flows:

- I. An adequate source of water supply.
- II. A retaining agent offering more resistance to the passage of water than the well or other opening.
- III. An adequate source of pressure.

The first requisite is not made specific as regards source, as artesian waters, as has been pointed out, are not derived from a single but from a variety of sources. In the case of the second requisite the retaining agent may be a stratum, a vein or dike-wall, a joint, fault, or other fracture plane, a water layer, or some one of a variety of other agents. The pressure, while primarily due to differences in level in the different parts of the artesian system, may be transmitted in such a variety of ways and is subject to so many modifying factors that the specification of a definite source is impracticable.

MODIFYING FACTORS

It is believed that the three factors of the preceding paragraph are all that can be considered as essential to artesian flows, all other postulated requisites being in reality modifying or accessory rather than essential factors. These secondary factors, which have been enumerated in the preceding pages, may be classed as follows:

SECONDARY FACTORS OF ARTESIAN FLOWS

- I. Hydrostatic factors (relating to pressure and movement).
 1. Factors mainly affecting pressure.
 - a. Barometric.
 - b. Temperature.
 - c. Density.
 - d. Rock pressure.
 2. Factors mainly affecting movement.
 - a. Porosity.
 - b. Size of pores or openings.
 - c. Temperature.
- II. Geologic factors (relating to reservoir).
 1. Character of reservoir.
 2. Retaining agents.
 3. Structure of reservoir.
 4. Topographic conditions.
 5. Conditions relating to supply.
 - a. Catchment conditions.
 - b. Conditions of underground feed.
 6. Conditions of leakage.

No remarks on this paper.

FIGURE 1.—CRATER OF XINANTECATL (NEVADO DE TOLUCA), MEXICO, FROM THE PICO DE FRAILE

The dome in the middle was the last phase of the latest eruption, and is analogous to the dome of mount Pelée, Martinique. It is composed of glassy andesite that welled up from the conduit, but did not form a flow.

FIGURE 2.—VOLCAN DE COLIMA, MEXICO, AUGUST 30, 1906

At left the parasitic cone formed in 1869; at right the lava stream of 1885, surmounted by that of 1903. Courtesy of the Am. Mus. of Nat. Hist. E. O. Hovey, photographer

CRATER OF XINANTECATL AND COLIMA VOLCANO

FIGURE 1.—BREADCRUST BOMB, COLIMA, MEXICO, ERUPTION OF 1903

FIGURE 2.—DEEP TRANSVERSE FURROWS OF LAVA FLOW OF 1869, COLIMA, MEXICO
Courtesy of the American Museum of Natural History. E. O. Hovey, photographer

The next paper was

VOLCANOES OF COLIMA, TOLUCA, AND POPOCATEPETL

BY EDMUND OTIS HOVEY

[*Abstract*]

The principal object of presenting the paper was to show the Society some photographs of these three volcanoes which were taken upon excursions made in connection with the meeting of the Tenth International Geological Congress in the city of Mexico, in September, 1906.

Toluca is the oldest of the three. A feature of greatest interest in the crater is the dome of vitreous andesite which welled up in the crater as the latest phase of the activity of the volcano and shows a certain resemblance to the new cone of mount Pelée. The volcano of Popocatepetl shows its composite character as a strato-volcano with great clearness in the walls of the crater, and streams of lava have been among the features of the most recent eruptions. The volcano of Colima is still sending up a vigorous column of steam from its central summit crater. From this summit crater there poured out, in the latest eruption (1903), streams of very frothy lava, which present a strange appearance on account of the porous character of the surface blocks. The same feature characterizes the streams of earlier eruptions and has led some observers to the erroneous conclusion that flows of lava have not occurred at the volcano of Colima.

Remarks were made by A. Heilprin and the author.

The next paper was

GEOLOGICAL STRUCTURE OF THE UINTA MOUNTAINS

BY S. F. EMMONS

Remarks were made by C. P. Berkey. This paper forms pages 287-302 of this volume.

The following paper was read by title:

STRATIGRAPHY AND STRUCTURE OF THE UINTA RANGE

BY F. B. WEEKS

This paper has been printed as pages 427-448 of this volume.

The next paper was

STRUCTURE OF THE FRANKLIN MOUNTAINS, TEXAS

BY G. B. RICHARDSON*

The paper was discussed by W. M. Davis.

* Introduced by C. W. Hayes.

The following paper was read by title:

PROBABLE AGE OF THE MEGUMA (GOLD-BEARING) SERIES OF NOVA SCOTIA

BY J. EDMUND WOODMAN

[Abstract]

From the earliest studies to the present, the age of this series has been referred to in terms that are either vague or dogmatic without requisite evidence, and has been placed from the Silurian to the pre-Cambrian. The most commonly stated age is Lower Cambrian, the judgment being based (1) on "fossils," (2) on lithological resemblance to other supposed Cambrian rocks. For the most part, recent references have merely stated the age, assigning no reason for the belief.

The "fossils" are believed to be inorganic and the lithological resemblances untenable as evidence.

Of the possible lines of evidence not before referred to, graphite, lime, and concretions are of no direct value. Scolithus and certain as yet undescribed trails from Halifax are serviceable only as evidence of organic forms of some undetermined kind. Unconformities show that all the great events in the history of the series are pre-Devonian except the intrusions, which are lower Devonian; but thus far fail to narrow the problem further. The composition of younger rocks shows detritus from the Meguma in the Devonian and Carboniferous, but nothing more definite as yet.

Comparative structures are held to be especially important. The whole of the peninsula of Nova Scotia is built upon the Meguma series as a base, and the rocks of all ages show some sympathy in the strike of their axes, with the very marked orogenic type exhibited by that series. There are no rocks on the mainland of Nova Scotia containing Cambrian fossils, and the next age represented by fossils is the Ordovician. These rocks show a different orogenic type from the Meguma; so different as to lead to the conclusion that the mountain-building of the latter entirely preceded the Ordovician. As the formation of the bedded veins was contemporaneous with the folding in the Meguma, this places the chief features of its history into the Cambrian or before.

The thickness of the series (nearly 30,000 feet) is regarded as far too great to be assigned to the Cambrian without any direct evidence. The intrusives of the Meguma are all acid and abyssal, except some dioritic material on the northern margin of the great western gneiss and a few dikes; and the amount of igneous rock is very great compared with the area of the sediments, one mass alone covering nearly 2,000 square miles. The granites intrude Silurian rocks in Annapolis, Digby, and Kings counties, but are older than the Horton series, mapped as Devonian, in Kings. They were intruded after all the main chapters in the history of the Meguma series had been enacted. No other rocks of the province are characterized thus, the next younger, the Ordovician, having basic hypabyssal types.

In previous literature correlation, direct or implied, has been made with various series from the Silurian down, those series themselves often of doubtful or unknown age. These correlations are regarded as unwarranted, and the

FIGURE 1.—TOWERS LEFT BY EROSION ON THE LEBANON MOUNTAINS

FIGURE 2.—ILLUSTRATION OF THE EROSION OF THE GLANDARIA LIMESTONE
EROSION FEATURES ON THE LEBANON MOUNTAINS

conclusion is reached that there are no sufficient data to permit correlation of the Meguma with any known series elsewhere; and the further conclusion that there is no positive evidence for regarding the Meguma as Cambrian, but that there is some warrant, not amounting to complete proof, for placing it in the pre-Cambrian.

It is expected that this paper will be printed in full in volume 19 of the Bulletin.

The last paper of the morning session in this section was

*ARTIFICIAL PRODUCTION OF GNEISSIC STRUCTURES BY CRYSTALLIZATION
UNDER STRESS*

BY FRED. EUGENE WRIGHT

The second section of the Society elected George P. Merrill chairman and H. M. Ami secretary, and then proceeded with the reading of papers, as follows:

ORIGIN OF THE LEAD AND ZINC ORES IN MISSOURI

BY E. R. BUCKLEY

The paper was discussed by H. C. Hovey, H. M. Ami, and the author. The next paper was

LEBANON GLACIER

BY G. FREDERICK WRIGHT

[*Abstract*]

Fifty years ago Sir Joseph Hooker reported that the cedars of Lebanon were growing upon a terminal moraine. Since that time references to the glaciation of the Lebanon mountains have been frequently made, and sweeping inferences drawn concerning the influence of the Lebanon glaciers upon the surrounding country, especially upon the water-levels in the Jordan valley. It was for the purpose of getting more accurate information upon the subject that in October, 1903, in company with Professor Alfred Day, of the Protestant College in Beirût, I set out to visit the principal grove of cedars, situated at the head of the Kadîsha river, east of Tripoli. The successful outcome of the expedition is largely due to Professor Day's comprehensive knowledge of the geology of the range, and I hereby acknowledge my great indebtedness to him.

The Lebanon mountains rise by a rapid and pretty uniform slope from the eastern shore of the Mediterranean to an average elevation of 5,000 or 6,000 feet, and opposite Beirût and Tripoli to elevations of from 8,000 to 10,000 feet, Jebel Samîn, east of Beirût, being 8,567 feet above the sea and Jebel Makhmal, east of Tripoli, 10,225 feet above the sea. This range extends for a distance

of 95 miles northward from the Litany river. In structure the Lebanon range is an anticline, embracing three series of Cretaceous rocks. The lower series, about 3,000 feet in thickness, is, however, thought by some to be Jurassic. This rock is a thick layer of limestone, sparingly supplied with fossils, among which are sponges, corals, brachiopods, and the characteristic *Cidaris glandaria*, and is generally known as the Glandaria limestone. This formation is exposed at the bottom of many of the deep gorges which furrow the western flank of the mountain, and appears in foldings in several places at an altitude of from 4,000 to 5,000 feet, where the superincumbent rocks have been removed by erosion. It is this limestone which is so extensively weathered into castellated towers and so beautifully fluted by the rains, of which we shall presently speak.

The middle series consists of sandstone, soft limestone, and clay, with occasional exposures of thin strata of poor bituminous coal, with pyrites and efflorescent salts and deposits of low grade iron ore. From its reddish color this series can be distinguished a long distance away. About half way up the mountain, near Afka, in the headwaters of the Adonis river, there is a considerable exposure of volcanic ash around what seems to have been a well defined crater connected with the volcanic eruptions accompanying the elevation of the range.

The third series is composed of hippurite limestone, and forms the summit of the range, resting there in nearly horizontal strata and attaining a thickness of from 3,000 to 5,000 feet. Upon the flanks of the mountain these strata are much contorted and have been extensively removed by erosion.

The eastern limb of the Lebanon anticline descends very precipitously into the valley of Beka'a, of Cœlesyria, which separates Lebanon from the parallel range of Anti-Lebanon, from 10 to 15 miles distant. Near Zahleh, on the east side, at an elevation of between 3,000 and 4,000 feet, occur porous sandstones and clays, with fossils of Miocene age.

In order more thoroughly to test the question of the general glaciation of the range, we traversed the region diagonally across the western flank of the mountain from Beirût to Jebel Makhmal, at whose western base the Kadîsha river has its source amid the cedars of Lebanon. The distance in a direct line is about 30 miles. The route, however, was a zigzag one, leading first up the Nahr-el-Kelb to the natural bridge at the head of the north fork, and then over the divide to the Adonis river, at Afka, and thence across the headwaters of the Jozek river to Bsherreh, in the upper portion of the Kadîsha. Throughout the larger part of this distance we were immediately underneath the summits of the range upon the east, where we should have observed ancient moraines and other signs of glaciation, if any had occurred; but there was a total absence of such signs, except possibly near Afka, where the debris accumulated at the base of the summit had somewhat the appearance of till; and again, near Akûra, where an immense amount of debris having accumulated, it was gradually sliding down to a lower level, as it was undermined by the northern branch of the Adonis river; but throughout this entire distance the sculpturing of the mountain side revealed the action of water rather than of ice. The deep and frequent gorges were wholly due to the action of water, while innumerable castellated and fantastically sculptured projections of the lower limestone conclusively demonstrated that ice had nothing to do with the general sculpturing of the mountain side.

FIGURE 1.—NATURAL BRIDGE HALF WAY UP THE MOUNTAIN
Summit of upper limestone in the background

FIGURE 2.—GORGE OF THE MIDDLE KADISHA
TOPOGRAPHIC FEATURES OF THE LEBANON MOUNTAINS

FIGURE 1.—DISTANT VIEW OF THE CEDARS, SHOWING THE MORaine SURFACE

FIGURE 2.—VIEW FROM THE UPPER PART OF THE MORaine TOWARD THE SUMMIT
Showing the notch down through which the glacier descended
CEDARS OF THE LEBANON MOUNTAINS

It was only as we approached Bsherreh, near the headwaters of the Kadisha, that true signs of glaciation appeared. Here, at an elevation of 5,300 feet, the front of a moraine extends from one side of the valley to the other, a distance of about four miles. The moraine consists of limestone fragments of all sizes from vast blocks down to fine powder, derived from the hippurite limestone forming the crest of the mountain a few miles back. In the middle of the valley the precipitous face of this moraine is several hundred feet thick, and is suffering erosion from a powerful current of water which issues as a spring from its base, this being the drainage from the whole area of the moraine back of it. Through the action of this stream a triangular area has been eroded from the moraine, ending in an acute angle, where the water emerges from beneath the surface. No scratched stones were found in the moraine, which is doubtless due to the fact that the material has been brought from only a short distance, and probably had for the most part been borne upon the surface of the ice, whither it had fallen from the overhanging ledges at the summit of the mountain. But that it was of glacial origin is certain from the promiscuous intermingling of coarse and fine fragments and from the impossibility of its having been brought into position through the action of gravity upon an ordinary talus, and still further from the topography of the area extending 4 or 5 miles back toward the head of the valley.

Upon ascending to the summit of this moraine, we find that it extends about 5 miles toward the apex of the valley and presents everywhere the knoll and bowl topography characteristic of a true terminal moraine. The grove of cedars which makes the region celebrated is growing upon the upper part of this moraine, at an elevation of 6,300 feet above the sea. Standing on the eastern edge of this grove, one sees the highest summit of the Lebanon range, some 3 or 4 miles distant, with a triangular depression, 100 or 200 feet below him, which was evidently occupied by ice during the last stages of the declining glacier, which had poured down from the small plateau on the summit through a narrow trough, clearly visible, and spread out as it had opportunity over the expanding area of the valley.

Upon ascending the summit, one finds a plateau, of several square miles, where the snows gathered during the Glacial epoch in sufficient quantity to supply the glacial ice that pushed down into the valley of the Kadisha, a distance of 8 or 9 miles, and to a level which is about one-half of the total height of the mountain above the sea.

Upon descending the mountain into Cœlesyria toward Ba'albek, we could find no signs of glacial action. In the local valley of a small stream which runs between the eastern foothills and the main range of Lebanon and Disappears in the sink of the Yamuneh, there is an immense accumulation of gravel, which in places appears in ridgelike accumulations, extending across the valley, which was evidently deposited by streams rushing down the steep side of the mountain and projecting the material brought down far out beyond the base, while occasionally there were immense boulders nearly a quarter of a mile from the base, which probably were carried out so far by their own momentum, attained in rolling down from a great height.

This moraine at the head of the Kadisha river, upon which the cedars of Lebanon are growing, has all the marks of youth which characterize the glacial deposits of America and Europe. The bowls upon the surface have not

been extensively filled up with sediment and the knolls have not been greatly diminished in height. The erosion by the stream issuing from the front of the glacier is comparatively limited in extent. More careful study of it may yet shed some light on the time that has elapsed since glacial accumulation ceased at the margin, but at present we have no data from which to make calculations.

An interesting subject of inquiry concerns the relation of this glacier to the ancient water-levels in the Jordan valley. It has been pretty generally assumed that the expansion of the Dead sea shown by the sedimentary terraces which surround the valley at an elevation of 650 feet, and, as Hull maintains, at an elevation of 1,400 feet, was the direct result of the glacial conditions in the Lebanon and Anti-Lebanon mountains. Even though no glacial streams could reach the Jordan valley from the Lebanon mountains, it is maintained that the increased precipitation and diminished evaporation of the region during that epoch would result in filling the Jordan valley with a body of water 200 miles long, 30 or 40 miles wide, and at its southern end 2,000 feet deep.

It is possible, however, to interpret the facts in a different way, and make the glaciation of the Lebanon mountains an effect of the increased area of the Dead sea rather than the cause. There is abundant evidence that the land all along the shore of the Mediterranean has risen 250 or 300 feet during the most recent geological era. Raised beaches containing shells of the same species that are now living in the Red sea and the Mediterranean are found near the pyramids in Egypt and in the vicinity of Jaffa and Lattakia, on the eastern shore of the Mediterranean. Now, if the land of this region were to be depressed 300 feet, it would admit the water of the Mediterranean into the Jordan depression through the valley of Esdraelon between Nazareth and mount Carmel. It is quite possible that the filling of the valley with water during that depression may have been the cause of the greater precipitation which produced the conditions favorable to the formation of the Lebanon glacier. If so, the various means we have of estimating the date of that depression of land may aid us in determining the approximate date of the Lebanon glacier. The general aspect of this moraine and the specific evidence bearing on the date of the enlargement of the Dead sea all point to these occurrences as comparatively modern events, estimated in tens of thousands of years rather than of hundreds of thousands of years.

Remarks were made by H. C. Hovey and the author.

ICE PRESENT DURING THE FORMATION OF GLACIAL TERRACES

BY F. P. GULLIVER

[*Abstract*]

This paper described with maps and lantern slides some glacial deposits along the Connecticut, Thames, and Quinebaug rivers which have usually been classed with the terraces formed by the down-cutting of the rivers. An example of terraces which have surely been carved by river action is found in the Westfield river west of Springfield, Massachusetts.

The deposits described along the Connecticut rivers were contrasted with those found at Westfield, and it was shown that they must have been formed

SNOW "CAVE" ON THE SUMMIT OF LEBANON
Showing stores incorporated in the strata of ice

before the ice had completely melted from the valleys. These deposits were therefore forms of aggradation and not forms produced by degradation.

Typical eskers, deltas, and kettle-holes are associated with these so-called terraces; and even where these deposits have the characteristic form of river cut terraces, cross-sections as revealed by railway or other cuts show delta structure rather than the structure of alluvial flood-plains. The delta lobes point either down stream or into side valleys, and there are frequently found unfilled portions of the main preglacial valley and of its tributaries below the level of the delta-terrace, between the delta-terrace and the rock walls of the older valley.

The discussion was participated in by F. G. Clapp, R. D. Salisbury, F. Leverett, Alden, and the author.

PLEISTOCENE GLACIAL PHENOMENA OF THE BOLIVIAN PLATEAU

BY W. G. TIGHT

The paper was discussed by H. F. Reid and C. H. Hitchcock.

PREGLACIAL DRAINAGE IN THE MISSISSIPPI VALLEY, A WORKING HYPOTHESIS

BY W. G. TIGHT

Remarks were made by F. Leverett and F. Carney.

The following paper was read by title:

GLACIAL FLOWAGE OVER NEW ENGLAND

BY J. B. WOODWORTH

COMPLEXITY OF THE GLACIAL PERIOD IN NORTHEASTERN NEW ENGLAND

BY FREDERICK C. CLAPP

Discussion by F. Leverett, C. H. Hitchcock, Alden, H. M. Ami, and the author. This paper is printed as pages 505-556 of this volume.

GLACIAL LAKE MEMPHREMAGOG

BY C. H. HITCHCOCK

[*Abstract*]

The existence of this lake was first announced at the meeting of the society in 1894.* Recent studies show that it was tributary to Glacial Lake Champlain by way of both the La Moille and Winooski valleys. When the ice filled the Champlain valley to the depth of a thousand feet the impounded water on the east side could have reached the Connecticut basin by way of White river.

* Bull. Geol. Soc. Am., vol. 6, p. 400.

Further descriptions of this rather unique body of water will be found presented in a "Report on the Champlain deposits of northern Vermont," printed as a part of the Fifth Report upon the Geology of Vermont, 1906, by Professor G. H. Perkins.

Discussion by G. F. Wright and C. H. Richardson.

PRE-WISCONSIN DRIFT IN THE FINGER LAKE REGION OF NEW YORK

BY FRANK CARNEY†

The paper was discussed by R. S. Tarr and the author.

This paper has been published in the Journal of Geology, volume xv, 1907, pages 571-585.

WAVE-CUT TERRACES IN KEUKA VALLEY OLDER THAN THE RECESSION STAGE OF WISCONSIN ICE

BY FRANK CARNEY

The paper was discussed by F. G. Clapp. This paper has been published in the American Journal of Science, volume xxiii, 1907, pages 325-335.

The section adjourned for luncheon.

The section of the Society convened again at 2.15 o'clock p m, with the first section (physical and structural geology) under the chairmanship of the Acting President.

The first paper read was

ORIGIN OF METEOR CRATER (COON BUTTE), ARIZONA

BY H. L. FAIRCHILD

Remarks were made by A. C. Lane and G. P. Merrill. The paper forms pages 493-504 of this volume.

The next paper read was

AFTON CRATERS OF SOUTHERN NEW MEXICO

BY W. T. LEE

This paper is printed as pages 211-220 of this volume.

The next paper read was

VOLCANIC NECKS OF THE MOUNT TAYLOR REGION, NEW MEXICO

BY D. W. JOHNSON*

This paper is printed as pages 303-324 of this volume.

† Introduced by H. L. Fairchild.

* Introduced by Ralph Arnold.

The following paper was then read :

*EARTH-FLOWS AT THE TIME OF THE SAN FRANCISCO EARTHQUAKE**

BY ROBERT ANDERSON

[Abstract]

Landslides of different kinds resulted in great numbers from the California earthquake of April 18, 1906. The term *earth-flow* is applied to landslides having the nature of flows of surface material partially saturated with water that were produced as a result of the earthquake by the sudden accession of water to points on the surface. Underground conduits of water suffered disturbances from the earthquake shock, and seepages occurred at the surface at new points or, in increased amounts, at old points of moisture concentration. In certain cases the water seems to have risen with a gush, as if actually squeezed from the hills. The effect of this abnormal supply of water was a saturation and loosening of the subsoil and a flowing movement of the surface debris away from areas so affected. The original loosening was aided by the vibratory movement of the shock, especially at points where some moisture had already gathered in the ground, the intensity of the shock in moist ground having been much greater than elsewhere. The water was the chief agent in causing the movement of the loosened material, as a result of the weight that it added and the semi-fluid nature that it gave to the mass. The process was usually one of sapping and undermining, the surface sod remaining dry and being broken into blocks and transported in a comparatively little disturbed position on a plastic substratum. At some of the earth-flows water continued to flow after the earthquake where it had not flowed before.

Earth-flows were of frequent occurrence in the Coast ranges, the writer finding them numerous on the San Francisco peninsula and in the Santa Cruz mountains within a score of miles of the fault, and what appeared to be flows of similar origin at a much greater distance from the epicentrum. They were formed on gentle as well as steep slopes, and both in previously dry drainage depressions and on convex hillsides. In the largest flows thousands of tons of earth and rock detritus were removed and carried hundreds of yards, leaving great cavities. In one case a hole ten feet deep was excavated over an area of nearly an acre on a five-degree slope, and the material removed was spread over two acres.

Such earth-flows, as well as the related slumps caused by earthquakes, in the case of which it is not easy to determine whether or not the slide or flow of soil has been aided by a suddenly increased water supply, are of special geologic importance for the reason that they aid in the initiation of drainage lines and the furtherance of degradation.

The following paper was read by R. B. Moore :

RADIO-ACTIVITY OF THE THERMAL WATERS OF YELLOWSTONE NATIONAL PARK

BY HERMAN SCHLUNDT AND RICHARD B. MOORE †

* This subject is treated more fully in the report of the Earthquake Commission being published by the Carnegie Institution of Washington.

† Introduced by C. W. Hayes.

Remarks were made by J. F. Kemp and A. C. Lane.

The following paper was then read by title:

OCCURRENCE OF UNUSUALLY LARGE CALCITE CRYSTALS IN NEW YORK STATE

BY D. H. NEWLAND

The next paper was

ASYMMETRIC DIFFERENTIATION IN A BATHYLITH OF ADIRONDACK SYENITE

BY H. P. CUSHING

The paper was discussed by A. C. Lane. It forms pages 477-492 of this volume.

The following paper was then read by title:

FORMATION OF LEUCITE IN IGNEOUS ROCKS

BY HENRY S. WASHINGTON

The next paper was

GENETIC RELATIONS OF SOME GRANITIC DIKES

BY ALFRED C. LANE

Contents

	Page
1. Effect of difference of conditions on crystallization.....	644
2. Application to pegmatites.....	645
3. Observations in the Huron mountains.....	645
4. Conclusions	647

1. EFFECT OF DIFFERENCE OF CONDITIONS ON CRYSTALLIZATION

One of the fundamental principles in considering the coarseness of grain of rocks is that, other things being equal, the less the difference between the conditions before and after crystallization, the coarser will be the crystallization. When the difference of conditions is small, the rate of change from the one state to the other will also be slow, and the crystallization proportionately coarse. We note this in the laboratory, where the coarse crystals are formed from the relatively cool or tepid solutions, and in the kitchen, where the conditions for making the coarsely crystallized rock candy are sharply contrasted with rapid cooling and stirring of a hot syrup required to make a "fondant," which forms the center of chocolate creams. This principle is one of the first consequences of the proposition that the grain is proportional to the slowness of cooling.*

* Annual Report Geological Survey of Michigan, 1903, p. 212.

2. APPLICATION TO PEGMATITES

As applied to the giant granites, the pegmatites, the ultra coarse-grained rocks, the principle mentioned suggests that the rock broth, or magma, from which they formed was not so different in its physical conditions, such as temperature, pressure, and gas saturation, from the surrounding rock. They might accordingly be formed by a raising of the rock to a point where any part of its constituents became fluid, or, what amounts to the same thing, where its inherent rock moisture has its solvent power markedly increased. This might be the origin of what might be called segregation pegmatites. But, on the other hand, we may consider them products of an ordinary granitic magma, occurring in the granite mass or in its contact zone.

The suggestion would then be that they should be a relatively late product of it, after its contact zone had been heated up and after the magma had lost some of its original fluidity, heat or gas, and was on the point of crystallizing—indeed, had perhaps largely crystallized already, leaving only a residual anchientectic* magma.

3. OBSERVATIONS IN THE HURON MOUNTAINS

The foregoing reflections came to me from some observations in the Huron mountains. The locations where these facts may be observed are so very numerous that they can hardly be particularized, and in fact Professor A. E. Seaman, of the Houghton College of Mines, who has vastly wider experience than I of this region, says, also, that the pegmatites cut the aplites, so far as he has seen.†

The country rock is a series of banded gneisses and hornblende-schists and amphibolites, Keewatin-Laurentian, forming anticlinal regions between the basins of Huronian rocks. The country was mapped by the first surveyors as "syenite and gneissoid granite," and has since been mapped with the Azoi, Laurentian, and Archean. At times the banding is well marked, strikingly like that of a series of sediments.‡

The amount of hornblende varies greatly. The strike of the banding is pretty constantly west northwest—that is, parallel to the axis of the synclinal of Huronian rocks that lies south.

This country rock is cut by various classes of dikes.

(1) In the first place, it is cut by the group of granitic dikes which we wish to study.

(2) They are cut by a group of uralite diabases, epidiorites, etcetera, which are probably of Huronian age.

(3) Latest are diabases, like that of Marquette, which are presumably Keweenawan.§

* Vogt's word—almost eutectic. Compare also my paper on the rôle of possible eutectics in rock magmas, *Journal of Geology*, 1904, pp. 86, 89.

† However, this paper is based mainly on notes and sketches on Huron mountain, section 39, township 52, range 20, and sections 24 and 25 adjacent; also mount Homer, section 31, township 52, range 28, near Mountain lake; also Pine lake, in section 28, township 52, range 28, especially, and section 4, township 52 north, range 27.

‡ So around Ives lake.

§ See Geological Survey of Michigan, vol. vi, part i, chapter x.

The coarsest country rock is a hornblende-gneiss, but the individual grains are rarely over 2 to 3 millimeters across. A tendency to porphyritic texture, like that of the Republic granite, is rare and not pronounced.*

In but few places does the hornblendic country rock fail entirely to be cut by red dikes composed mainly of quartz and feldspar. There is but little sign of one being in excess over the other, and they are probably nearly eutectic.

Generally the rock is riddled with them and they stand in relief, and may show glacial striæ when the rest of the rock does not. They are close welded to the country rock, so that it is not very difficult to get a piece showing both sides of the contact. Indeed, frost will sometimes spall off a section clear across a small dike including a little of the country rock on either side. Sometimes, as at the top of Huron mountain, they are so abundant that at first sight the country rock appears like a lot of basic fragments or secretions inclosed in a general mass of granite: yet it is very easy to see that through all these apparent fragments there is a constant strike and dip of the schistosity in which the red granitic matter does not partake. A more careful examination, moreover, shows the nexus of dikes which make up the general granitic effect.

Now we may for convenience sake divide these dikes into (a) those whose grain is finer than the country rock, less than 2 to 3 millimeters, often about 1 millimeter, which we will call aplites; (b) those whose grain is coarser than the country rock, often 50 millimeters or so, which we may call pegmatites. Exceedingly coarse pegmatites I did not see.

We find the following variation of grain in individual dikes:

1. Dikes aplitic in general grain, with coarser border.
2. Dikes aplitic, but with coarser pegmatite bands near, but not at, the side.
3. Dikes granitic in grain, with coarser pegmatitic border.
4. Dikes with a fine-grained margin and coarser center. This type seems rare, and in many, but not all, cases may be explained by the splitting of a fine-grained dike by a later coarse-grained one.
5. In many cases the grain seems uniform—at least to casual inspection—throughout.

It is not uncommon to find a concentration of the darker minerals, the mica or hornblende, at the center of the dike.

A very striking dike of pegmatite (figure 1) on mount Homer, 28 inches wide, shows irregular prisms of quartz growing from the margin, while the center is made up of patches of graphic granite. Figure 1 is, however, only a memory sketch from notes at the time.†

In a rough way the larger pegmatites are also coarser in grain; but while both the pegmatites and the aplites are close welded and have about the

* I noticed a trace with feldspar grains 10 millimeters long on the south side of Second Pine lake. A splendid porphyritic rock at the foot of mount Homer, and near J. M. Longyear's house, on Ives lake, I take to be a Huronian dike.

† I remember seeing a similar structure in a pegmatite dike on the north side of Sand lake, Proudfoot township, Parry sound, Canada. The dike was 32 inches broad, and quartz radiated from each side for about 4 inches, while the dike as a whole was made up of biotite, muscovite, feldspar, and quartz in good-sized crystals.

same tendency to follow straight lines or cut irregularly across the country rock, with perhaps no exception the coarser dikes cut the aplites.

4. CONCLUSIONS

The inference seems warranted that the granitic dikes, aplites and pegmatites together, represent one period of invasion of the Keewatin hornblende schists and gneisses by the granitic magma.*

In the beginning the difference of conditions of country rock and magma was considerable. Still the absence of fine-grained margins shows that the conditions of solidification of the granitic magma (pressure, temperature, mineralizers) were nearer the conditions of the country rock than those of

FIGURE 1.—Structure of a granitic Dike on Mount Homer.

The quartz grows out for 4 inches (100 millimeters) from the margin in rude prismatic forms. The rest of the way is mainly filled with patches of graphitic granite. Quartz is black; feldspar white.

the magma, or at times just about half way, and then the aplitic dikes formed.

Later the temperature of the country rock rose, or it otherwise approached the conditions of liquefaction for granite. The granitic magma also cooled. At any rate, the interval decreased† from the conditions of consolidation of the dikes to the conditions of the country rock, and so the later pegmatites are coarser. If the granitic magma did not also cool, they would become even more than the aplite uniform in grain from center to margin except, so far as it was modified by a tendency of substances existing in excess of the eutectic to crystallize out earlier, or a tendency to grow first from the walls, as in veins, the same tendency that makes nuclei hasten the crystallization of an undercooled fluid.

Such tendencies may explain the coarser grain at the margin (see figure 1) in part, but it may also be explained by supposing that the granitic magma cooled as fast as the country rock heated, so that the conditions of consolidation were about half way between them.

Criticism has been made of my studies of grain that I have not allowed enough for undercooling and varying velocity of crystallization.‡

I think it will be found that I have always recognized that other factors than rate of cooling were of importance, and my mathematical work only gives a first approximation, but I have insisted that observation shows that often it has been of leading importance, and my results have been obtained

*Compare Rosenbusch, "Elemente der Gesteinslehre," p. 220.

† v of my annual report, Geological Survey of Michigan for 1903, pp. 211 and 212.

‡ Doelter: Petrogenesis, p. 45.

by definite, if not exact, numerical measurements, showing that the grain in some cases did vary as it would if the rate of cooling were the dominant factor. It must be remembered, too, that it is more than probable that most igneous magmas contain catalyzers—gases of one kind or another that promote liquidity and restrain viscosity—and that, as regards the later formed minerals, the earlier act as centers or nuclei of crystallization, notably orthite in granite, promoting crystallization and preventing undercooling, thus making the grain more responsive to the rate of cooling. Indeed, the jars recorded by seismographs may have some effect.

To give more concrete expression to our conclusions, let us insert some hypothetical figures:

Suppose the hornblende country rock was at first at $100 +$ degrees centigrade (the close welded contacts suggest that it was not wet). The temperature of formation of the aplites was probably not above 800 degrees centigrade (the inversion point of quartz to tridymite, according to Day and Allen, is about 760 degrees at ordinary pressures). If so, the initial temperature of the aplite-forming magma was a little less than 1,500 degrees ($2 \times 800 - 100$) $+ 100$; but by the time the coarsest pegmatites of this region formed the country rock temperature may have risen to over 700 degrees and the magma temperature fallen to less than 900 degrees, if we suppose the grain to be 50 times greater in the pegmatite than in the aplite.

This is merely an illustration, for I am inclined to think that the interchange of vapor pressure, or gases, is quite as important as the interchange of heat, and the growth of quartz crystals from the margin of the pegmatites is especially due to it.

The next paper was

OPHITIC TEXTURE

BY A. C. LANE

[Abstract]

Ophitic texture is not a mere synonym of the diabasic, but a variety of the poikilitic* made by the inclosure of tabular, or lath-shaped, feldspar in augite, by which a mottled appearance is often produced, like that of some ophidians (figure 1, plate 70). This mottling, when coarse enough, appears:

(1) On smooth, dull, or matte surfaces, like those of pebbles or diamond drill cores (figure 2, plate 70).

(2) On fresh fracture very faintly, but in the sunlight the reflection on large, interrupted cleavage faces of the augite produces the effect called by Pumpelly "luster-mottling."

(3) On joints and rifts the pattern is brought out in a variety of color effects, according to the type of weathering.

(4) On weathered surfaces that are also exposed to erosion a lumpy, warty, or pock-marked appearance is produced, which gave rise to the name "varioid greenstone," as applied to ophitic melaphyres (plate 71).

* Geological Survey of Michigan, vol. vi, part 1, pp. 51, 227; Journal of Geology, 1906, p. 705.

FIGURE 1.—PHOTOGRAPH OF OPHIDIAN (GILA MONSTER)
Showing ophitic pattern. Kindness of Zoological Department of Michigan Agricultural College

FIGURE 2.—DIAMOND DRILL CORE
Ophitic mottling brought out near chlorite seam. Kindness of W. J. Penhallegon

FIGURE 3.—OPHITIC TEXTURE IN THIN-SECTION
After Irving. Monograph v, plate ix
OPHITIC TEXTURE

BLUFF BACK OF DELAWARE MINE

Taken by W. J. Penhallegon, August, 1907, to illustrate effect of weathering on ophitic texture. The white object is a slide rule six inches long and one inch wide. The exposure is about 100 feet from the base of the "greenstone"

(5) Where weathering goes on faster than erosion, the rock is often broken down into a coarse gravel, each grain of which is likely to be a single augite crystal.

(6) The thin-section (figure 3, plate 70) shows the texture plainly, and how it is produced by the fact that the augite, while readily enveloping the labradorite, crowds ahead of it the corroded remnants of the olivine (at whose expense it is formed) as well as the iron oxide.

The following paper was then read by title:

OCCURRENCE OF DIAMONDS IN NORTH AMERICA

BY GEORGE F. KUNZ

The following paper was then read:

SILVER-GOLD ORES AT SAN PEDRO DE GUANACEVI, DURANGO, MEXICO

BY FREDERICK B. PECK

The following paper was then read by title and transferred with permission of the author to Section E, American Association for the Advancement of Science, and read there December 31.

PERSPECTIVE VIEW OF THE SUBMARINE CANYON OF THE HUDSON RIVER

BY J. W. SPENCER

The following papers were read by title:

TITANIFEROUS BASALTS OF THE WESTERN MEDITERRANEAN

BY H. S. WASHINGTON

THE PALEOZOIC SECTION OF THE UPPER YUKON

BY A. H. BROOKS AND E. M. KINDLE

STRATIGRAPHIC SUCCESSION NORTH OF COOK INLET, ALASKA

BY SIDNEY PAIGE AND ADOLPH KNOPP *

This paper appears as pages 325-332 of this volume.

SEISMOLOGICAL OBSERVATIONS OF THE UNITED STATES

BY H. F. REID

The following paper was read by title and was transferred with per-

* Introduced by A. H. Brooks.

mission of the author to Section E, American Association for the Advancement of Science, where it was read December 31, 1906:

CHARLES WILLSON PEALE'S PAINTING, "THE EXHUMING OF THE FIRST AMERICAN MASTODON"

BY ARTHUR BARNEVELD BIBBINS

Charles Willson Peale, who was born in Queen Anne county, Maryland, in 1741, and who became known as "The Artist of the Revolution," was one of the pioneers—perhaps the Nestor—of American vertebrate paleontology. Although very few of his scores of paintings relate to this subject, one which has lately come to the writer's notice, with the correspondence by the Peales and others relating to the subject of the painting, clearly demonstrates the artist's keen, intelligent, and practical interest in this direction—the most advanced of his time in this country.

The canvas measures 5 by 6 feet and is known as "The Exhuming of the First American Mastodon." It is signed "C. W. Peale" and dated 1806.

The scene portrays the unearthing of the first of the two mastodons which the artist obtained near Newburg-on-the-Hudson in 1801, and which were the first approximately complete skeletons found in this country. This century-old picture has recently been brought to scientific notice, since it has come into the possession of the family of one of the artist's lineal descendants, by whose kind permission this reproduction of the painting is shown.

The canvas is well preserved and its detail clear and distinct. The artist is represented as personally supervising the excavation. Members of his family, including his wife, Rembrandt and his wife, Titian and James Peale, and several scientific worthies of the day, are in attendance, and the whole countryside is, as will be seen, astir with the unwonted activity in their midst.

Rembrandt Peale, the no less celebrated son of the elder Peale, has left a very graphic account of the labors attending the excavation. This rare little pamphlet was written in London in 1803, while the incidents were vividly in mind. A copy of it was brought to my notice at the Maryland Historical Society, in Baltimore, and I have been much gratified to find yet another copy here, in the library of the American Museum of Natural History, which has been kindly loaned me for this occasion. I will refer to a few paragraphs from it.

According to Rembrandt Peale's account, his father had received word in the spring of 1801 that in the previous autumn an old farmer, in digging into a marl pit in the vicinity of Newburg, had discovered a number of gigantic bones, which he heaped on the floor of his granary or garret.

The thigh bone, 3 feet 9 inches in length, was first discovered, and for several succeeding days 100 men were encouraged by the physicians in the neighborhood to continue their efforts with very considerable success.

By the use of "rum" in too great abundance, however, the men became unruly and impatient and nearly destroyed the skeleton, and by using oxen and chains to drag the bones forth from the clay or marl, the head, hips, and tusks were much broken and many parts left behind. So great a quantity

THE EXHUMING OF THE FIRST AMERICAN MASTODON

of water from copious springs, bursting from the bottom, rose upon the men that it required several score of hands "to lade it out with milk-pails, buckets, and bowls." After the fourth day the attempt was given up, and it was in this condition when Peale arrived upon the scene, in 1801. Peale purchased the excavated bones from the farmer for a small sum, reenforced by the judicious gift of gowns to his wife and daughter and a gun to his son.

All necessary articles for the excavation had been provided from New York—pumps, ropes, pulleys, etcetera—and after weeks of close labor, a millwright fashioned the machinery and completed a large scaffold and a wheel 20 feet in diameter, wide enough for three or four men to walk abreast in, to supply tread power. A rope around this turned a spindle, which worked a chain of buckets. The water was raised and emptied into a trough and conveyed into a basin a short distance away. A ship's pump assisted the work, and later a pair of barrels, raised by a crane, removed the mud.

Several weeks of unremitting work, during which bank after bank fell in, resulted in the discovery of but a few additional bones, while the 25 hands which Peale employed at high wages were kept at their task partly by their eagerness to uncover the head and under jaw.

Finally the work was relinquished, but again taken up 20 miles west of the Hudson, where the particular bones desired were found in another mastodon, together with others sufficient to fashion two skeletons. One was exhibited in Peale's museum, which he had recently opened in Philadelphia and where he gave lectures upon natural history. Family tradition recites that after the skeleton was mounted in the Philadelphia museum a celebration and dinner were given beneath it.

The other was exhibited by his son Rembrandt in Baltimore and became the nucleus of the "Peale museum" there, which he erected and which bears upon its front to this day the faded legend:

"Erected by Rembrandt Peale,
"Baltimore Museum, 1817-1830
"City Hall, 1830"

Both the Philadelphia and the Baltimore museums, the former started in 1802 and the latter about 1813, appear to have resulted from the Newburg finds; and, since these were the first public natural history museums in America, must we not credit the Peales as well with the nestorship of American public museums?

The scope of their interest along this line is suggested by another family tradition,* which relates that while the elder Peale and Baron von Humboldt were being entertained at a formal "three o'clock dinner" by President Monroe, the guests improved this opportunity by asking the President to try to induce Congress to establish a National museum, and that Peale returned from the interview much elated by the belief that some action would shortly be taken.

Tradition also adds that Peale was first induced to undertake the exhuming of the mastodon by Von Humboldt, who had visited this country before and been entertained by the artist.

* This is stated in C. W. Peale's diary, *not* tradition.

Of the find itself, Dr John C. Warren, the New England author of a monograph on the mastodon, said in 1852 that "there probably never has been a discovery in science which excited more intense interest." He relates that Peale's mastodons were at that time the greatest curiosities in natural history in America, and virtually credits the artist with the awakening in this country of the first genuine interest in vertebrate paleontology.

Of considerable interest in connection with this painting is a letter written by Peale, just after its completion, to the celebrated Benjamin West, whose early pupil he had been. It appears in the *Pennsylvania Magazine of History and Biography*, volume ix, 1885, pages 130-132.

I quote a single paragraph:

"PHILADELPHIA, *December 16, 1807.*

"DEAR SIR:

"Desiring to represent the scene of getting up the Mammoth bones, it being a very interesting article of the museum, last summer I undertook a picture of it. This picture contains a great number of figures, for while I was engaged in that labour my exertions excited the admiration of all the people for a considerable distance round that country; few people could conceive why I should incur so much expense to get the bones. Although I have introduced upwards of 50 figures, yet the number of spectators in fine weather amounted to hundreds. Eighteen of my figures are portraits, having taken advantage of taking most of this number of my family. I have introduced my painting this picture in order to state to you that I still doubted my ability of making a tolerable picture, and at the commencement of it I made use of a temporary easel and other apparatus for painting; however, as I advanced in the work it seemed to engross my whole attention, and I really took pleasure in painting from morn till night and even to use lamplight. I then ordered the cabinet-maker to make me a commodious mahogany easel, &c.; so instead of burning my pencils and totally quitting the art, as I thought very probably would be the case, I found it much less difficult than I imagined and have ever since regretted that I had not taken a larger canvas and devoted more time to give a higher finish to the piece. I often say that the aged ought not to be discouraged from undertaking works of improvement. . . .

"Make my most respectful compliments to Mrs. West, and believe me with great esteem, your friend,

"C. W. PEALE.

"MR. BENJAMIN WEST,
London."

The painting is at present in the custody of the Woman's College of Baltimore, to which it was loaned by Mrs. Harry White, of Massachusetts, and where its centenary unveiling will shortly occur.

RELATIONS OF THE ITHACA AND CHEMUNG FAUNAS OF WESTERN MARYLAND

BY C. K. SWARTZ

This paper was transferred with the author's consent to Section E.

American Association for the Advancement of Science, and read there at the session of December 31, 1906.

The section adjourned to the room occupied by the second section, where papers were still being read.

The second section met at 2.15 o'clock p m and reorganized by electing A. P. Coleman chairman and Collier Cobb secretary. The stratigraphic papers on the program were taken up.

The first paper was read by title:

DISCOVERY OF CAMBRIAN ROCKS IN SOUTHEASTERN CALIFORNIA

BY N. H. DARTON

The next paper was

LIMESTONES OF WESTCHESTER AND PUTNAM COUNTIES, NEW YORK

BY CHARLES P. BERKEY

The following paper was then read:

GALENA SERIES

BY FREDERICK W. SARDESON

This paper forms pages 179-194 of this volume.

The following papers were read:

AGE AND STRATIGRAPHIC RELATIONS OF THE CHATTANOOGA BLACK SHALE

BY AMADEUS W. GRABAU

THE MEDINA SANDSTONE PROBLEM

BY AMADEUS W. GRABAU

PALEOGEOGRAPHY OF THE AMERICAN DEVONIO

BY CHARLES SCHUCHERT

Illustrated by a series of seven lantern slides.

The following paper was then read by title:

CARBONIFEROUS OF THE APPALACHIAN BASIN

BY JOHN J. STEVENSON

The full paper has been printed as pages 29-178 of this volume.

The following papers were read by title:

STRUCTURE OF THE DEEP RIVER TRIASSIC

BY COLLIER COBB

RED BEDS OF OKLAHOMA AND ADJACENT STATES

BY CHARLES NEWTON GOULD

STRUCTURE AND CORRELATION OF NEWARK TRAP ROCKS OF NEW JERSEY

BY J. VOLNEY LEWIS

This paper has been printed as pages 195-210 of this volume.

SOME RESULTS FROM THE STUDY OF THE CAMBRIAN BRACHIOPODA

BY CHARLES D. WALCOTT

The paper, which was an informal explanation of the results obtained in the preparation of a monograph on the Cambrian Brachiopoda, was illustrated by eighty of the plates, combined on twenty lantern slides, presenting the leading characteristics of the Cambrian Brachiopoda.

The next paper was read by title:

CRYPTOZOAN: GENERA, SPECIES, RELATIONSHIPS

BY HENRY M. SEELY

CRUSTACEAN FAUNA OF THE SHAWANGUNK GRIT IN EASTERN NEW YORK

BY JOHN M. CLARKE

ADDITIONAL FOOTPRINTS FROM THE CARBONIFEROUS SHALES OF MASSACHUSETTS

BY J. B. WOODWORTH

The reading of papers was declared finished.

RESOLUTION AS TO DIVISION OF MEETINGS INTO SECTIONS

It was moved that the Council be requested to consider the question of dividing each meeting into three sections for the reading of papers, consideration being had of the number and character of communications presented in the several branches of geologic science.

After discussion by A. W. Grabau, F. W. Sardeson, A. F. Foerste, and J. F. Kemp, the motion was passed.

Hearty votes of thanks were passed with reference to the hospitality offered the Society and the arrangements made for its meeting by Columbia University, the American Museum of Natural History, and the local committee.

The Society then adjourned.

REGISTER OF THE NEW YORK MEETING, 1906

The following Fellows were in attendance at the meeting:

F. A. ADAMS.	A. HAGUE.
H. M. AMI.	C. W. HAYES.
R. ARNOLD.	A. HEILPRIN.
H. F. BAIN.	R. R. HICE.
R. M. BAGG.	R. T. HILL.
J. BARRELL.	C. H. HITCHCOCK.
G. H. BARTON.	W. H. HOBBS.
F. BASCOM.	J. A. HOLMES.
R. BELL.	T. C. HOPKINS.
C. P. BERKEY.	E. O. HOVEY.
A. BIBBINS.	H. C. HOVEY.
A. P. BRIGHAM.	E. HOWE.
R. W. BROCK.	E. E. HOWELL.
A. H. BROOKS.	J. P. IDDINGS.
E. R. BUCKLEY.	R. T. JACKSON.
S. CALVIN.	A. A. JULIEN.
H. D. CAMPBELL.	A. KEITH.
W. B. CLARK.	J. F. KEMP.
J. M. CLARKE.	E. H. KRAUS.
J. M. CLEMENTS.	H. B. KÜMMEL.
C. COBB.	G. F. KUNZ.
A. P. COLEMAN.	A. C. LANE.
W. O. CROSBY.	D. W. LANGTON.
W. CROSS.	W. T. LEE.
H. P. CUSHING.	F. LEVERETT.
W. M. DAVIS.	W. LIBBEY.
J. S. DILLER.	W. LINDGREN.
B. K. EMERSON.	H. D. McCASKEY.
S. F. EMMONS.	W J McGEE.
H. L. FAIRCHILD.	W. McINNES.
N. M. FENNEMAN.	W. D. MATTHEW.
A. F. FOERSTE.	F. J. H. MERRILL.
M. L. FULLER.	G. P. MERRILL.
A. C. GILL.	A. M. MILLER.
C. H. GORDON.	B. L. MILLER.
A. W. GRABAU.	W. G. MILLER.
H. E. GREGORY.	W. H. NILES.
F. P. GUILLIVER.	H. F. OSBORN.

F. B. PECK.	W. G. TIGHT.
R. A. F. PENROSE.	E. O. ULRICH.
G. H. PERKINS.	C. R. VAN HISE.
L. V. PIRSSON.	F. R. VAN HORN.
J. H. PRATT.	G. VAN INGEN.
A. H. PURDUE.	C. D. WALCOTT.
H. F. REID.	C. H. WARREN.
W. N. RICE.	H. S. WASHINGTON.
C. H. RICHARDSON.	W. H. WEED.
H. RIES.	F. B. WEEKS.
R. D. SALISBURY.	L. G. WESTGATE.
F. W. SARDESON	D. WHITE.
C. SCHUCHERT.	I. C. WHITE.
W. B. SCOTT.	R. P. WHITFIELD.
E. A. SMITH.	H. S. WILLIAMS.
G. O. SMITH.	B. WILLIS.
A. C. SPENCER.	A. W. G. WILSON.
J. W. SPENCER.	J. E. WOLFF.
J. STANLEY-BROWN.	J. B. WOODWORTH.
T. W. STANTON.	F. E. WRIGHT.
J. J. STEVENSON.	G. F. WRIGHT.
R. S. TARR.	W. S. YEATES.

Fellows-elect

R. S. BASSLER.	E. HUNTINGTON.
F. G. CLAPP.	T. A. JAGGAR, JR.
H. F. CLELAND.	D. W. JOHNSON.
J. A. DRESSER.	

Total attendance, 127.

SESSION OF THE CORDILLERAN SECTION, FRIDAY, DECEMBER 28, 1906

The eighth annual meeting of the Cordilleran Section of the Society was called to order in room 71, Stanford University, California, at 10.50 a m, December 28, 1906, by the Chairman of the Section, Professor J. C. Branner.

The minutes of the seventh annual meeting were read and approved.

The following officers were elected for the ensuing year: A. C. Lawson, Chairman; George D. Louderback, Secretary, and G. K. Gilbert, Councillor.

The scientific program was then taken up and the following papers were presented and discussed:

NOTES ON THE STRUCTURE OF THE SANTA CRUZ RANGE, CALIFORNIA

BY J. F. NEWSOM

[Abstract]

The major fault lines of the range have northwest-southeast directions varying from north 20° west to north 45° west. Various movements have occurred along the major fault lines. Movement along some of the major lines has probably been to some extent lateral.

A secondary system of faults lying at angles of from north 60° west to north 75° west also occurs. These are reversed faults and are parallel with the principal axes of folding in the range.

The folds and secondary faults were probably formed by lateral movement and consequent crushing along the major fault lines.

NOTES ON THE TOPOGRAPHY OF THE SEWARD PENINSULA, ALASKA

BY J. F. NEWSOM

[Abstract]

The accumulation of ice, with its protective influence, has been an important factor in forming the rounded dome-like topography of the Seward peninsula. The protective influence of moss is also an important factor. In many places where small spots are free from moss, but surrounded by it, hogwallows are formed. Considerable hillocks are occasionally developed in old filled pond basins by the squeezing action of the freezing sands and gravels pushing up the unfrozen materials from below.

OCCURRENCE OF MIDDLE TERTIARY MAMMAL-BEARING BEDS IN NORTH-WESTERN NEVADA

BY JOHN C. MERRIAM

This paper was published in Science, new series, volume xxvi, 1907, pages 380-382.

The meeting then adjourned for the noon recess.

The afternoon session began at 2.00 p m and the first paper was read by title:

NOTES ON SOME CALIFORNIA MINERALS

BY AUSTIN F. ROGERS

The following papers were then read and discussed:

*TRANSPORTATION OF DETRITUS BY YUBA RIVER**

BY G. K. GILBERT

[Abstract]

In the process by which running water transports detritus two factors are distinguished. The larger particles are rolled along the bottom, or else lifted

* Published by permission of the Director of the U. S. Geological Survey.

momentarily, so as to make short leaps; they constitute the *bottom load*. The smaller are lifted far from the bottom, are sustained for long periods, and are distributed through the whole body of the current; they constitute the *suspended load*. The classification of load in each particular cause depends chiefly on velocity of current; a relatively swift stream will include in its suspended load relatively coarse detritus.

Many determinations have been made of the suspended load of streams, the method including (1) the measurement of the total quantity of water, and (2) the measurement of the ratio of load to water in samples taken from the stream. The determination of the bottom load is comparatively difficult, and there is much need of information as to its relative and absolute amount. The writer has taken advantage of an unusually favorable opportunity to measure the bottom load of the Yuba river, one of the tributaries of the Sacramento river in California.

The river issues from a mountain gorge at the Smartsville narrows, and thence to its mouth, a distance of 20 miles, traverses a valley. In consequence of a special condition brought about by hydraulic mining, it is now aggrading its channel in this part of its course. The growing deposit is of coarse gravel near the narrows, and of sand near the mouth, with gradual transition. During high floods the suspended load near the mouth of the river includes silt only; near the narrows it probably includes sand also, and possibly some small gravel.

In the summer of 1904 a dam 6 feet high was built at a point 5 miles below the narrows, where the channel is 1,500 feet wide; and the basin created by this dam was filled with detritus by the floods of the following winter and spring. In the summer of 1905 an addition was made to the dam, increasing its height 8 feet, and during the construction of this addition a contour map of the channel above the dam was made by the U. S. Geological Survey. The river remained low and the water clear until the following January, when a large flood occurred, lasting about one week. After the flood had passed the channel was resurveyed, and the two maps were afterward used to determine the quantity of detritus arrested. The character of the material deposited was investigated by means of pits and borings in the summer of 1906. During the flood the river was gaged to determine its discharge, and at the time of flood maximum the water was thoroughly sampled near the river mouth to determine the percentage of suspended load.

The discussion of the data thus obtained is not yet complete, but certain results of geologic bearing may be given at this time. All of the figures are approximate and subject to correction. The detritus which passed the mouth of the canyon during the flood of January, 1906, would cover a square mile to a depth of about 17 inches. The ratios of its chief components were: Silt, 24; sand, 24; gravel, 52. In this statement whatever sand was deposited in the interstices of the gravel is accounted with the gravel; by transferring it to the sand category the ratios are changed to: Silt, 24; sand, 34; gravel, 42. Near the river mouth, where the suspended load consisted of silt and the bottom load of sand, the ratio of bottom load to suspended load was about 1:1. Near the narrows, if the silt only be accounted as suspended load, its ratio to the bottom load was as 1:3; but if, as is probable, the sand also should be classed with the suspended load, the ratio is about 1:1.

A report on the subject is in preparation and will be published by the U. S. Geological Survey.

CROCIDOLITE-BEARING ROCKS OF THE CALIFORNIA COAST RANGES

BY GEORGE D. LOUDERBACK AND WM. J. SHARWOOD

[Abstract]

Although in no great quantity in any one place, crocidolite-bearing rocks have been found to be rather widely distributed in the Coast ranges of California. They may be classed as crocidolite-bearing cherts or quartzites and crocidolitic schists. A petrographical description was given, analysis presented, and mode of occurrence described. Evidence was offered that many if not all of the rocks are derived from radiolarian cherts, and that they are formed by the introduction of soda into the rock by solution.

PRIMITIVE CHARACTERS OF AMERICAN TRIASSIC ICHTHYOSAURS

BY JOHN C. MERRIAM

[Abstract]

The following characters generally appearing in the Triassic Ichthyosauria may be considered as survivals of characteristics of ancestral shore forms. Where the term "relatively" has been used the unexpressed comparison is with the Jurassic ichthyosaurs.

1. Limbs relatively large and first two segments relatively long.
2. Hind limbs relatively large.
3. Epipodial, metapodial, and phalangeal elements often elongated and with shafted median region.
4. Number of phalanges in each digit relatively small.
5. Number of digits not increased beyond five.
6. Pelvic elements relatively heavy; inferior elements much expanded; ischium and pubis not fused.
7. Skull generally relatively short compared with the length of the trunk region.
8. Jaws relatively short.
9. Maxillaries relatively long and premaxillaries relatively short.
10. Orbits relatively small in the earliest forms.
11. Vertebral centra averaging relatively long.
12. Anterior cervical vertebræ not fused.
13. Neural spines of vertebræ relatively thick.
14. Zygapophysial facets of upper arches in the anterior portion of the vertebral column not united or brought into the same plane.
15. Caudal intercentra large and united ventrally to form large Y-shaped chevrons.

FAUNA OF THE ASPHALT BEDS EXPOSED NEAR LOS ANGELES, CALIFORNIA

BY JOHN C. MERRIAM

This paper was published in *Science*, new series, volume xxiv, 1906, pp. 248-250.

ORIGIN OF SOUTH AMERICAN BEARS

BY JOHN C. MERRIAM

[Abstract]

The spectacle bear of South America represents a type known only from the western hemisphere. Its nearest allies are found in genus *Arctotherium*, occurring in the Quaternary of North and South America. Two views have been expressed concerning the origin of this type. The most nearly related forms are probably Asiatic.

The next three papers were read by title.

*RECONNAISSANCE OF A RECENTLY DISCOVERED QUATERNARY CAVE DEPOSIT
NEAR AUBURN, CALIFORNIA*

BY E. L. FURLONG*

This paper was published in Science, new series, volume xxv, March, 1907, pages 392-394.

*PHYSIOGRAPHIC CHANGES BEARING ON THE FAUNAL RELATIONSHIPS OF
THE RUSSIAN AND SACRAMENTO RIVERS, CALIFORNIA*

BY RULIFF S. HOLWAY†

This paper was published in Science, new series, volume xxvi, September, 1907, pages 382-383.

SAN PABLO FORMATION OF MIDDLE CALIFORNIA

BY CHAS. E. WEAVER*

The last paper of the day read and discussed was

TWO MOUNTAIN RANGES OF SOUTHERN CALIFORNIA

BY W. C. MENDENHALL

[Abstract]

The San Gabriel and the San Bernardino ranges are adjacent mountain masses, separated only by Cajon pass, and holding identical relations to the valley of southern California, and to the Mojave desert, lying to the north and east of it. They also are similarly related to the principal fault lines of this part of California, each of them being bounded along its southern margin by a major fracture, and one of them, the San Gabriel, certainly being limited in a similar way along its northern base, while less definite evidence indicates that the San Bernardino range is related in the same way to the desert lowland. Little is known of the geology of the interior of the two ranges, but that little indicates that in a broad way they contain similar masses of rocks, which probably differ but little in their ability to resist the attacks of erosional agents.

* Introduced by J. C. Merriam.

† Introduced by George D. Louderback.

The similarities, however, disappear when the land forms are examined. The San Gabriel range has been completely dissected, resulting in thoroughly graded streams, sharp peaks, and knife-like ridges of discordant heights. No level areas at or near the summits, nor in the valley bottoms, exist within the mountain mass. The San Bernardino range contrasts sharply with its neighbor in these respects. Throughout its western end there is a strikingly level sky-line at an elevation of 5,000 feet or more. It contains many broad meadows, with lakes and playas, separated by smooth ridges. The topography of the central part is, in brief, topography of an old, well reduced type. About its periphery, however, topographic forms are strikingly new. Several of the streams are not reduced to grade; they meander through broad uplands in the central part of the range, then plunge over falls into steep canyons, which they follow to the valleys that border the ranges.

For these striking topographic differences there seems to be no adequate explanation in the rock types, in the relative masses of the ranges, in their relation to major drainage lines, nor in their relation to precipitation. It is concluded, therefore, that since each is a faulted, uplifted block, the San Bernardino mass, in which old topographic forms are well preserved, is much later in origin than its neighbor, the San Gabriel range, in which none of these old forms are now to be found.

The Section then adjourned for the day.

SESSION OF THE CORDILLERAN SECTION, SATURDAY, DECEMBER 29, 1906

The meeting was called to order at 9.15 a m by the Chairman, and proceeded immediately to the reading and discussion of papers presented, as follows:

RELATION OF THE LOS ANGELES-OWENS RIVER AQUEDUCT TO THE FAULT LINES OF INYO AND SAN BERNARDINO COUNTIES

BY J. C. BRANNER

NOTES ON THE GEOLOGY OF THE MOUNT HAMILTON QUADRANGLE

BY JOHN F. NEWSOM AND RODERIC CRANDALL

[Abstract]

Discussion of the geological terranes present, their distribution and relations; of the main structural features, with special note of the larger faults and direction of movement; and of the relations of structural features to the topography.

CRETACEOUS STRATIGRAPHY OF THE SANTA CLARA VALLEY REGION

BY RODERIC CRANDALL*

[Abstract]

The fossils from the Cretaceous localities are listed according to their geographic distribution. Probabilities of the presence of Horsetown at mount

* Introduced by J. F. Newsom.

Diablo are discussed. The presence of Horsetown in northern California and the absence in southern California is taken as proof of a differential land movement on the west coast during mid-Cretaceous times.

PHYSIOGRAPHIC FEATURES OF SOUTH CENTRAL OREGON

BY G. A. WARING*

[Abstract]

This paper deals with the region forming the northwestern end of the Great basin. The basin range type of structure described by Gilbert and Russell is here well developed, the main features of the relief being the fault-scarps of the tilted monoclinial blocks. These, together with their associated undrained basins, now occupied by remnants of more extensive Quaternary lakes, and the peculiar lack of well defined drainage over much of the area are the three distinctive physiographic features. Low folding has also taken place and has resulted in certain definite topographic effects. In addition to the major scarps, there are numerous minor scarps, some of which may be faults, but others of which have evidently been produced by weathering agencies. The rock masses observed within the region are representative of various igneous effusive series exclusively. Here and there bosses of older rocks of medium basicity occur, but by far the greater part of the area is floored by a series of sheets of basalt and associated tuffs, which have been affected by the structures that have given rise to the striking physiographic features of the region.

The next paper was read by title:

THE METAMORPHIC AND CRYSTALLINE ROCKS OF THE SANTA CRUZ MOUNTAINS

BY SOLON SHEDD

The following were then read and discussed:

GENERAL GEOLOGICAL FEATURES OF THE TRUCKEE REGION EAST OF THE SIERRA NEVADA

BY GEORGE D. LOUDERBACK

[Abstract]

Contents

	Page
Area described	663
Bedrock complex	663
The great unconformity.....	664
The Tertiary lavas	664
Andesites	664
Rhyolites	665
Basalts	665
Distribution of types	665
Tertiary sediments	665
Post-Tertiary orogenic movements.....	666
Lake Lahontan	668
The Truckee river.....	668

* Introduced by W. C. Mendenhall.

AREA DESCRIBED

The area described includes a belt lying north and south of the Truckee river and extending from the main east front of the Sierra Nevada just west of Verdi, Nevada, to the Truckee-Carson divide near Hazen. The chief topographic divisions, passing from west to east, are: (1) the Sierran fault front; (2) the Verdi-Long Valley depression at its base; (3) the line of elevation represented by the Carson range on the south and Peavine mountain and others to the north; (4) the depression of Spanish Springs valley, Truckee meadows, Washoe valley, etc.; (5) the Virginia range; (6) the depression of Pyramid lake, the lower Truckee valley, etc.; (7) the mountains lying between the Truckee and Humboldt regions. (For topography, see Sierraville, Reno, Carson, and Wadsworth sheets of the United States Geological Survey.)

BEDROCK COMPLEX

The Bedrock complex consists chiefly of granitic rocks, largely granites (in part granodiorites), with residual masses of more or less metamorphosed sedimentary and igneous rocks into which they were intruded. The chief sedimentary types are, perhaps, quartzites and mica-schists, including muscovite-schists. A petrographically interesting meta-dacite is found at intervals for at least 12 or 15 miles north of Reno. Limestone and slate are occasionally found, and to the south, near Mound House, gypsum.

The Bedrock complex is found in the Sierran front, in Peavine mountain, and in a number of the small ranges to the north and northeast of Reno, and in the Carson range to the south. Near Steamboat springs both sediments and intrusions have been mapped and described by Becker.*

With one or two exceptions, the Bedrock complex is exposed only west of the Virginia range. This range is made up of a great thickness of Tertiary igneous material which has so deeply buried the older rocks that they have not yet been exposed by the Truckee river, which has cut a canyon directly across the range over 2,000 feet deep. The pre-Tertiaries, however, rise to the surface in this range some miles south, in the vicinity of Virginia City. The Marble bluff at the south end of Pyramid lake is of limestone, presumably Triassic.

Judging from a comparison between the Bedrock complex here and that in the Sierran province immediately west, the Humboldt region to the east, and certain areas to the south, it is probable that the schists are Mesozoic, possibly in part Paleozoic, and the granitic intrusives post-Jurassic. There is absolutely no evidence suggesting the Archean age of the rocks of the Peavine Mountain area, although like several areas since proved Mesozoic, it was so marked by the Fortieth Parallel Survey and their mapping was recently copied on the map of Bulletin 208 of the United States Geological Survey. Until some evidence or suggestive line of reasoning indicates its presence, it is misleading to map any areas of northwestern Nevada as Archean.†

The chief folding of the Bedrock complex undoubtedly took place at the time of the extensive post-Jurassic orogenic disturbances that affected so strongly the regions just east and west of the Truckee country. The granites have

* U. S. Geological Survey, Monograph xiii.

† See also Bull. Geol. Soc. Am., vol. 15, pp. 317, 335, 340.

been stripped of their covering over such large areas that the axes of folding are not generally determinable, but evidence so far observed consistently points to major axes not departing many degrees from north and south. This is in harmony with the structural lines of the pre-Tertiaries of adjoining areas.

THE GREAT UNCONFORMITY

After the folding and intrusion of the Bedrock complex follows a long period of erosion, during which the products of disintegration and decay were carried beyond the borders of this region and a country of comparatively low relief was produced. The succeeding rocks are frequently deposited on a plane granite floor, but in several places evidence of granite hills is obtainable. This erosion interval included the Cretaceous and lower Tertiary.

THE TERTIARY LAVAS

ANDESITES

The most abundant and varied volcanics of the region are andesites, chiefly hornblende andesites, varying at one extreme to augite andesites, sometimes of basaltic texture and at the other to quartz-mica-hornblende andesites with well developed phenocrysts.

Along the Carson block and the upper Truckee the more common sequence is coarse andesitic breccia with finer ash layers lying directly on the granite and capped by hornblende andesite flows. Andesites also occur on the southern flank of Peavine mountain and in the low hills north and south of Reno; also in great abundance in the Virginia range. In these latter areas breccia is only occasionally present. In some parts of the Virginia range the andesites are amygdaloidal, and they often show the propylitic type of alteration, especially near Virginia City and in the Olinghouse Canyon district.

These various andesites are very similar to and are practically continuous with those of the Sierra Nevada and are presumably of the same age—late Miocene or early Pliocene. Although the most abundant of the volcanic series of this region and of the Sierra Nevada, evidences of craters are entirely lacking, and even the foci of eruption are not generally evident. On the upper slopes of Peavine mountain the granites are cut by a number of dikes identical lithologically with andesite flows lying on the lower slopes, and were apparently loci of eruptions in that area.

The andesites have often suffered alteration by solfataric action or mineralization. This generally occurs along faults which have been seats of deposition, the rising waters sometimes (1) impregnating a zone with sulphides and very little silica, sometimes (2) with silica and very little sulphides, sometimes (3) with both well developed. Occasionally calcite gangue is found. In the first case the mineralized zone may now appear on the surface as a belt of excessive decomposition showing the combined effects of oxidation and sulphuric acid (from the oxidized sulphides) action. The ground is soft and easily eroded, is bright red from iron oxide, or leached to various shades with white as the end result. Many of these belts carry low (at present non-commercial) values in gold and silver, but occasionally rather high. At the Wedekind mine several hundred dollars a ton in silver was found at the surface, due to secondary concentration. In general the most common original sulphide is pyrite, with occasionally galena and sometimes sphalerite.

The second type gives rise to silicified zones, which weather in prominent reefs. The silica occurs partly as a deposit in cracks or other open spaces with distinct crustification and often drusy structures, partly as a cement between the crushed fragments and particles of the rock and partly as a replacement of the rock, the original porphyritic structure often being visible in the silicified product. These three modes of deposition are commonly found in the same ledge. Many of these occur in low hills north of the Truckee meadows.

The Olinghouse Canyon mines are in fissured (commonly crushed and sheeted) zones in andesite and carry gold without much deposited gangue material. The best known deposits in these rocks are along the Comstock lode and need not here be described. They belong to the third type mentioned above.

RHYOLITES

The rhyolites (and geologically related dacites) occur to the north and northeast of Reno, and more abundantly in the Virginia range, where they vary from vitreous, sometimes perlitic, to lithoidal, often porphyritic, and occur both as lava flows and dikes. They are less abundant than the andesites. They may also be accompanied by mineralization as in the White Horse district. White Horse mountain is a stock of rhyolite, and dikes pass from it out into the surrounding andesites. Good values have been found in some of the dikes. Veins in rhyolite also occur to the south near Gold Hill.

BASALTS

The latest of the Tertiary eruptions are basalts. They are found about Steamboat springs, along the summit region of the Virginia range, and abundantly in the ranges just east of this. There is commonly evidence of some deformation before the outpouring of the basalts.

DISTRIBUTION OF TYPES

A noteworthy change takes place in the character of the predominant types of Tertiary lavas as we pass from west to east across the Truckee region. In the western part the andesites occupy the field almost to the exclusion of other types. In the central part, especially in the Virginia range, we get the greatest complex. The andesites show the greatest mineralogical varieties, and although still the dominant rock type they are accompanied by considerable quantities of rhyolite and basalt. To the east of the Virginia range the andesites practically disappear, and we have a broad area, forming the eastern edge of the Truckee region, but lying chiefly in the Humboldt region, which is characterized by a great thickness of rhyolite flows and tuffs overlain by caps of basalt.

TERTIARY SEDIMENTS

The Tertiary sediments are entirely freshwater or subaerial. The greatest part has been included under the title Truckee beds. The type locality is the Kawsoh mountains and along the east edge of the Virginia range, at the eastern edge of the Truckee region. Other areas are found in the summit region of the Virginia mountains south of the Truckee river; in the Truckee meadows, especially flanking the Carson block and extending up the river to the Sierra fault front west of Verdi; along the west side of Spanish Springs valley; in

the valley northeast of Peavine mountain and running north in Long valley to the neighborhood of Honey Lake valley. It is very probable that many of the valley areas in which the beds are not now exposed really carry them overlain by Quaternary deposits. In several such places this has been demonstrated by shallow railroad grades, trenches, and wells.

These beds lie unconformably on the granites and other members of the Bedrock series, and consist largely of poorly cemented sands and clays, the former being sometimes gray and friable, sometimes charged with limonite, and occasionally nodular. The lateral corrasion of the Truckee river a few miles above Reno causes occasional landslides, that produce cliffs several hundred feet high, exposing beautiful sections of the stratification (plate 73).

Diatomaceous earth, frequently of great purity, is a characteristic member of the series and just west of Reno is about 400 feet thick. Besides diatom remains, it contains sponge spicules and a slight amount of mineral fragments. Fish remains are found in it east of the Virginia range. Some layers are ashy and carry small fragments of pumice and crystal (apparently pyroclastic) fragments of minerals of igneous rocks.

Current-bedded gravels, tuffs, and blue tuffaceous sandstones are common, especially in the lower part of the series near Verdi. We also find fossil leaves and lignite beds, the latter formed by leaf and twig accumulations (that is, drifted material) often intimately associated with laminae or with a general admixture of volcanic ash. A shell limestone was observed in the Kawsoli mountains near Hazen.

These beds lie on the weathered surface of the andesites, but they precede the block faulting of the ranges, some areas being found in the valleys and some in the summit region of the Virginia and other ranges. They are characteristically disturbed by tilting and some folding, though except along fault planes they generally lie at moderate angles of dip, say from 10 to 40 degrees. The plant remains are of the broad leaf type (or reeds) and point to a lower altitude of the country at that time. These various relationships indicate that the beds may be upper Miocene or Pliocene.

An interesting section of the basal beds occurs a mile or two southwest of Reno. The lava dips under the sediments and carries a layer of pre-Truckee soil. The sediments show several alternations of distinct lake sediment and alluvial material or wash, each a few feet or yards thick. This seems to indicate a lake with considerable fluctuation in areal extent, and possibly therefore occupying an interior basin without outlet (at least part of the time) similar to the Quaternary Bonneville or Lahontan.

The eastern part of the Truckee region and the Humboldt region are characterized by thick rhyolitic ash accumulations. They may in part have accumulated subaerially on desert valleys, but in part were deposited in water and carry sand and pebble beds. Whether these should be associated with the Truckee beds or separated as a later series has not been determined.

POST-TERTIARY OROGENIC MOVEMENTS

At the end of the Tertiary or early in the Quaternary the acute deformation took place that gave rise to the present mountain ranges. Folding or bending movements were restricted in occurrence, the most characteristic movements being those of block faulting of the Basin Range type. Most commonly the

FIGURE 1.—PEAVINE MOUNTAIN FAULT BLOCK.

Main scarp on right hand (northeast) side. Looking northwest from Reno

FIGURE 2.—LANDSLIDE IN TRUCKEE LAKE SANDS, WITH RIVER SAPPING ITS BASE

Looking southeast from near Laughtons. Photograph by G. J. Young

FAULT BLOCK AND LAKE BEDS, TRUCKEE REGION

faulting was accompanied by tilting, one slope of the range being the more or less eroded fault scarp, the other the tilted older surface, which often shows but slight relief. The faulting is determined by topographic form, by the relation of older structures to range front, and by the attitude and distribution of the lake beds, ash beds, or lavas, as the case may be.*

The areas of uplift (or downthrow), and therefore the fault scarps, generally strike north-south or at angles not far removed from that direction. High values are represented by the fault on west side of Spanish Springs valley, which strikes north 28 degrees east, and the Peavine fault, north 45 degrees west. Where short independent ranges are produced, as best developed in the country north of the Truckee meadows, they are generally not bounded on the north or south by faults, but the block is flexed so that its surface of deformation rises in a curve from the valley on the north and descends unbroken underneath the valley to the south, like the axis of an anticlinal fold.† A longitudinal curvature of the depressed portions is probably the rule, but rarely observable. An exceptional opportunity for observation is afforded by the depression between the north end of the Carson Range block and the Peavine Mountain block. The Carson range is a large block with high scarp on the east side, and rises to 10,800 feet in mount Rose. Going north it rapidly drops to the Truckee river (4,500 to 5,000 feet). On the river slope the lowering is about 4,000 feet in 4 miles. Peavine mountain is also a fault block with a scarp on its north-east side and tilted gently to the southwest. It rises to the height of 8,270 feet, and drops off to the Truckee river to the south at the rate of about 1,000 feet to the mile. The lower part of the depression between the two blocks is occupied by lake beds lying on the andesites and the whole folded into a syncline, with axis east and west, the Truckee River valley lying along this axis. In other words, the surface of deformation of the Carson block drops to the north in a curve somewhat sharper than usual and terminates the range; it then rises again from the valley and becomes the curved surface of deformation of Peavine mountain.

The principal fault scarps recognized, together with the present elevation of range summit above surface of valley at the base of the scarp, follow:

Range.	Scarp facing.	General strike.	Feet.
Sierra west of Verdi.....	East.....	North-south.....	3,400
State Line range.....	West.....	North-south.....	2,900
Peavine mountain.....	Northeast.....	North 45 degrees west.....	3,300
Small ranges northeast of Peavine.....	East.....	{ North 5 degrees east }.....	2,000
Range west of Spanish Springs valley.....	South of east.....	{ North 12 degrees east }.....	1,500
Carson range.....	East.....	North 28 degrees east.....	5,500
Virginia range—		North-south.....	
By Truckee meadows.....	West.....	North-south.....	2,500
By lower Truckee valley.....	East.....	North 5 degrees west.....	4,000

It may be noted that, the large blocks have faults closely approximating or practically running north-south. The fault differing most from this direction is the shortest one given in the table.

Secondary faults within the general range blocks are not uncommon, giving

* Methods already described in detail in Bull. Geol. Soc. Am., vol. 15, 1904, pp. 289-346.

† This was also pointed out for the Humboldt mountains, loc. cit., p. 333.

rise to benches* or small valleys with one side a fault scarp, that scarp most commonly showing a throw in like sense to the primary fault. In the Virginia range, where there are probably two primary faults, a graben was found bounded both east and west by fault scarps in part broken into steps. In some places, instead of a scarp formed by movement along a single plane, are found a series of step faults, which distribute the upper layer along the face of the scarp, making it appear as very much thicker than normal, or giving rise on distant view to doubt as to the existence of fault at that point. Such a front is found on the Carson range not far from Reno and on parts of the Virginia range.

LAKE LAHONTAN

The lower Truckee River valley and other valleys east of the Virginia range were occupied by the Quaternary lake Lahontan and carry in their lower parts the practically horizontal and unconsolidated lake beds. The sides of these valleys are striated by the shorelines, with fresh cliffs, terraces, etcetera, and are prominently coated by the Lahontan calcareous tufas already described by Russell.†

We may note that the lake beds are found about 12 miles up the Truckee River canyon, showing that it had been cut even below its present level before lake Lahontan was formed. In the upper part of the canyon the stream flows on Tertiary volcanic bedrock.

The Truckee meadows were not reached by the Lahontan water level, but throughout that period, as at present, were occupied by the Truckee river and its floodplains. Well developed terraces have been cut in the lake beds at least up to 5,100 feet 5 miles west of Reno (700 feet above the river). Some of the terraces have but a thin veneer of river deposit, but one about 80 to 100 feet above the river is underlain by a thick deposit of coarse boulders.

THE TRUCKEE RIVER

The Truckee river leaves the main Sierra front near Verdi, passes along the axial belt of the depressed area (syncline) between the end of the Carson range and the Peavine Mountain block, and flows out onto the main Truckee meadows. Crossing these, it strikes right across the slope of the Virginia range, which rises abruptly from the valley some 2,500 to 3,500 feet, and cuts by a deep canyon almost perpendicularly across the range, emerging on the depressed block to the east, where it turns north and follows the topographic (orogenic) control to Pyramid lake. If we could fill up the Truckee canyon, the river would flood the meadows and finally seek Pyramid lake to the north-east by a shorter route.

It seems very evident that the Truckee river is an antecedent stream, in existence before the present ranges, and that the Virginia mountains have been slowly raised across its course, the river being active enough to maintain its course, and for that distance at least (25 miles) remain independent of the orogenic control which is almost universal in this region.

It is probable that during the gradual uprising of the mountains the Truckee

* See figure illustrating subsidiary faulting in the Carson range, loc. cit., plate 16, and discussion, p. 341.

† U. S. Geological Survey, Monograph xi.

FIGURE 1.—GAP WHERE TRUCKEE RIVER LEAVES THE MEADOWS AND ENTERS THE VIRGINIA RANGE

Mountains rise 2,000 to 3,000 feet above the valley. Looking east. Photograph by G. J. Young

FIGURE 2—PLEISTOCENE GRAVELS WEST OF RENO

Thirty-foot fault near center of section, its scarp destroyed by river planation. Aggradation shown here probably related to upward movement in Virginia range, which is visible in the distance

meadows were often flooded, and temporarily—perhaps for many years at a time—occupied by a lake. At present the river is flowing over volcanic bed-rock at the entrance to the canyon, while the valley is occupied by alluvium and lake beds. The ground-water moving from the west has a level which gradually approaches the surface and appears above it in hollows, producing more or less marshy ground near the Virginia Range front. We have therefore what is practically an underground lake which, save for percolation into the bedrock, must discharge over the rim at the canyon opening. During periods of considerable water supply from the west the ground-water level rises above the ground and a certain area becomes flooded. Renewed faulting along the range front would quickly produce a “permanent” flooded area.

The Truckee River channel has occasionally been occupied or displaced by Quaternary basaltic lavas. One flow occurred in the canyon through the Virginia range. It extended clear across the canyon, and the river, hemmed in by hard volcanic walls, was forced to cut through it, and has left it as flat terraces projecting from the canyon walls.

About 10 miles west of Reno successive flows came down from the flanks of Peavine mountain and occupied a channel cut in Tertiary lake beds and andesites. In this case the river found it easier to cut through the lake beds, and now flows at the base of the basalt, which rises 450 feet in a striking cliff. A long basalt flow, now left as terraces, is to be found higher up the river near Truckee.

Two sets of hot springs, probably genetically connected with the acute post-Tertiary deformation, are found on the east flank of the Carson range (Steamboat springs) and about 5 miles west of Reno (Laughtons springs). Several wells sunk along the front of the Carson range have struck hot water, indicating other hot springs that do not reach the surface, but feed direct into the ground water.

The Quaternary river deposits are faulted just west of Reno sometimes as much as 30 feet, and scarps topographically very young are frequently observed. In gathering data concerning the San Francisco earthquake of 1906 information was obtained concerning small shocks felt distinctly at several towns, but rapidly dying away in distance and so circumscribed that they pointed to recent disturbance along the Virginia mountains. The most active part of the orogenic cycle inaugurated in the early Quaternary is probably passed, but the forces do not seem to have reached yet a condition of equilibrium.

The writer is indebted to Professor G. J. Young for three of the photographs used in the illustration of this paper.

A COLLECTION OF FOSSIL FISHES FROM NORTHEASTERN BRAZIL

BY J. C. BRANNER AND D. S. JORDAN

DIATOMS AS A SOURCE OF SILICA IN SEDIMENTARY BEDS

BY J. C. BRANNER

Adjournment was then taken until afternoon.

The Section reassembled and resumed its program at 2.40 p m, the following papers being presented and discussed:

GEOLOGICAL RECONNAISSANCE IN THE SAN JOAQUIN VALLEY

BY H. R. JOHNSON*

THE LATEST PHASE OF THE COLORADO RIVER PROBLEM

BY W. C. MENDENHALL

The Section then passed a vote of thanks to Stanford University for the use of its rooms and for other courtesies shown during the meeting.

The Section then adjourned sine die.

GEORGE D. LOUDERBACK,
Secretary.

REGISTER OF THE MEETING OF THE CORDILLERAN SECTION

The following fellows were in attendance at the meeting:

J. C. BRANNER.	W. C. MENDENHALL.
G. K. GILBERT.	J. C. MERRIAM.
A. C. LAWSON.	J. F. NEWSOM.
G. D. LOUDERBACK.	OLON SHEDD.

In attendance as visitors were:

E. P. CAREY.	E. S. LARSEN, JR.
RODERIC CRANDALL.	J. PERRIN SMITH.
H. C. HARTZELL.	G. A. WARING.
L. M. HOSKINS.	G. A. WILCOX.
H. R. JOHNSON.	G. J. YOUNG.

* Introduced by W. C. Mendenhall.

ACCESSIONS TO THE LIBRARY FROM OCTOBER, 1906, TO OCTOBER, 1907

By H. P. CUSHING, *Librarian*

Contents

	Page
(A) From societies and institutions receiving the Bulletin as donation ("Exchanges")	671
(a) America	671
(b) Europe	673
(c) Asia	677
(d) Australasia	677
(e) Africa	678
(f) Hawaiian Islands	678
(B) From state geological surveys and mining bureaus	678
(C) From scientific societies and institutions	678
(a) America	678
(b) Europe	679
(c) Asia	679
(D) From Fellows of the Geological Society of America (personal publications)	679
(E) From miscellaneous sources	680

(A) FROM SOCIETIES AND INSTITUTIONS RECEIVING THE BULLETIN AS DONATION ("EXCHANGES")

(a) AMERICA

NEW YORK STATE MUSEUM,	ALBANY
3041. Annual Report, no. 57, part 2f.	
3042-3046. Bulletins 90-105.	
BOSTON SOCIETY OF NATURAL HISTORY,	BOSTON
2922. Proceedings, vol. 33, nos. 1-9.	
MUSEO NACIONAL DE BUENOS AIRES,	BUENOS AIRES
3158-3159. Anales, serie 3, tomo vi-viii.	
CHICAGO ACADEMY OF SCIENCES,	CHICAGO
1913. Bulletin no. 4, part 2.	
3157. Bulletin no. 6.	
FIELD MUSEUM OF NATURAL HISTORY,	CHICAGO
2715. Geological series, vol. iii, no. 5.	
3066. Report series, vol. iii, no. 1.	
CINCINNATI SOCIETY OF NATURAL HISTORY,	CINCINNATI
COLORADO SCIENTIFIC SOCIETY,	DENVER
2782. Proceedings, vol. viii, pp. 257-314.	

NOVA SCOTIAN INSTITUTE OF SCIENCE,	HALIFAX
2798. Proceedings and Transactions, vol. xi. part 2.	
MUSEO DE LA PLATA.	LA PLATA
3019. Revista, tomo xi.	
3020. Anales, Seccion Paleontologica, V.	
3021. Anales, Seccion Botanica, I.	
CUERPO DE MINAS DEL PERU,	LIMA
2939. Boletin, nos. 35-41.	
3010. Boletin, nos. 42-47.	
INSTITUTO GEOLOGICO DE MEXICO,	MEXICO
2829. Boletin, num. 22.	
3154. Boletin, num. 24.	
SOCIEDAD GEOLOGICA MEXICANA,	MEXICO
NATURAL HISTORY SOCIETY OF MONTREAL,	MONTREAL
AMERICAN GEOGRAPHICAL SOCIETY,	NEW YORK
2859. Bulletin, vol. xxxviii, nos. 5-12.	
3050. Bulletin, vol. xxxix, nos. 1-8.	
AMERICAN MUSEUM OF NATURAL HISTORY,	NEW YORK
3055. Bulletin, vol. xxii.	
NEW YORK ACADEMY OF SCIENCES,	NEW YORK
3014. Annals, vol. xvii, part 1.	
AMERICAN INSTITUTE OF MINING ENGINEERS,	NEW YORK
3170. Transactions, vol. xxxvii, 1906.	
GEOLOGICAL SURVEY OF CANADA,	OTTAWA
2979. Annual Report, new series, vol. xiv, 1901.	
2995. Annual Report, new series, vol. xv, 1902-03.	
3014. The Cruise of the Neptune.	
3031. Summary Reports, 1905-06.	
3032. Section of Mines, Annual Report, 1904.	
3033. Separate reports on mining districts, 1904.	
2462. Geologic sheets, 59-65, 74-76, 82-83, Nova Scotia.	
877. Paleozoic Fossils, vol. iii, parts 1-3.	
3142. Mesozoic Fossils, vol. i, parts 3-5.	
256. Contributions to Canadian Paleontology, vol. i, part 5.	
953. Contributions to Canadian Paleontology, vol. ii, part 2.	
3145. Contributions to Canadian Paleontology, vol. iii, part 1.	
3141. Contributions to Canadian Paleontology, vol. iv, parts 1-2.	
3143. Micropaleontology of the Cambro-Silurian, parts 1-2.	
3144. Fossil Plants of the Erian, etc., part 2.	
3146. Figures and Descriptions of Canadian Organic Remains, decades
1, 3-4.	

ROYAL SOCIETY OF CANADA,	OTTAWA
3171. Proceedings and Transactions, second series. vol. xii, part 1.	
ACADEMY OF NATURAL SCIENCES,	PHILADELPHIA
Proceedings, vol. lviii, parts 1-3, 1906.	
AMERICAN PHILOSOPHICAL SOCIETY,	PHILADELPHIA
2354. Proceedings, vol. xiv, nos. 182-184.	
2647. Transactions, new series, vol. xxi, part 3.	
3013. The Franklin Bicentennial Celebration.	
MUSEO NACIONAL DE RIO DE JANEIRO,	RIO DE JANEIRO
3151. Archivos, vol. xiii.	
CALIFORNIA ACADEMY OF SCIENCES,	SAN FRANCISCO
GEOLOGICAL SURVEY OF NEWFOUNDLAND.	SAINT JOHNS
ACADEMY OF SCIENCE,	SAINT LOUIS
2903. Transactions, vol. xvii, nos. 1-9.	
COMISSÃO GEOGRAPHICA E GEOLOGICO,	SAO PAULO
2914. Boletim 12-14.	
3048. Boletim 17-21.	
NATIONAL GEOGRAPHIC SOCIETY,	WASHINGTON
2845. National Geographic Magazine, vol. xvii, nos. 9-12.	
3023. National Geographic Magazine, vol. xviii, nos. 1-10.	
LIBRARY OF CONGRESS,	WASHINGTON
SMITHSONIAN INSTITUTION,	WASHINGTON
3049. Annual Report, 1905.	
UNITED STATES GEOLOGICAL SURVEY,	WASHINGTON
2905-2906. Professional Papers 46-47, 49-50.	
3004-3009. Water Supply Papers 155-184.	
3179-3180. Water Supply Papers 185-194.	
3017. Twenty-seventh Annual Report.	
3060. Monograph i.	
3061-3063. Bulletins 275-287.	
3175-3178. Bulletins 288-303.	
3059. Mineral Resources, 1905.	
3022. Professional Papers 51-52.	
UNITED STATES NATIONAL MUSEUM,	WASHINGTON

(b) *EUROPE*

DEUTSCHE GEOLOGISCHE GESELLSCHAFT,	BERLIN
2861. Zeitschrift, band lvii, heft 4.	
2994. Zeitschrift, band lviii, heft 1-3.	
KÖNIGLICH PREUSSISCHEN GEOLOGISCHEN
LANDESANSTALT UND BERGAKADEMIE,	BERLIN

- GEOGRAPHISCHE GESELLSCHAFT, BERNE
 SCHWEIZ. GEOLOGISCHE KOMMISSION, BERNE
 2801. Spezialkarten 31, 34-36, and karte vii, with text.
 R. ACCADEMIA DELLE SCIENZE DELL' INSTITUTO DI BOLOGNA, BOLOGNA
 NATURHIST. VEREIN DES PREUSSISCHEN RHEINLANDE, WESTFALENS UND DES REG.-BEZIRKS OSNABRÜCK, BONN
 2838. Sitzungsberichte und Verhandlungen, 1905, hafte 2.
 2839. Sitzungsberichte und Verhandlungen, 1906, hafte 1-2.
 3071-3126. Verhandlungen. Jahrgang 1-4; 6. supplemental heft; 7-8; 10-13; 14, parts 3-4; 15-17; 19-29; 30, hafte 2; 31-50; 51, hafte 2; 52.
 3127. Autoren-und Sachregister zu Band 1-40.
 3128-3134. Verhandlungen, Jahrgang 53-59.
 ACADÉMIE ROYALE DES SCIENCES DES LETTRES, ET DES BEAUX-ARTS DE BELGIQUE, BRUSSELS
 2934. Bulletin de la Classe des Sciences, 1906.
 3069. Annuaire. 1907.
 SOCIÉTÉ BELGE DE GÉOLOGIE, DE PALÉONTOLOGIE, ET D'HYDROLOGIE, BRUSSELS
 2766. Bulletin, tome xix, fasc. 3-5, 1905.
 2996. Bulletin, tome xxi, 1906.
 BIUROULI GEOLOGICA, BUCHAREST
 1435. Geological Map of Roumania, sheets 27, 29, and 34.
 MAGYARHONI FÖLDTANI TARSULAT, BUDAPEST
 2893. Földtani közlöny, xxxvi kötét, 1-12 fuset, 1905.
 NORGES GEOLOGISKE UNDERSØGELSE, CHRISTIANIA
 DANMARKS GEOLOGISKE UNDERSØGELSE, COPENHAGEN
 DET KONGELIGE DANSKE VIDENSKABERNES SELSKAB, COPENHAGEN
 2887. Oversigt i Aaret, Förhandlingar 1906, nr. 4-6.
 3135. Oversigt i Aaret Förhandlingar 1907, nr. 1, 2.
 NATURWISSENSCHAFTLICHE GESELLSCHAFT ISIS, DRESDEN
 2885. Sitzungsberichte und Abhandlungen, Jahrgang 1906.
 ROYAL SOCIETY OF EDINBURGH, EDINBURGH
 2913. Proceedings, vol. 771.
 3052-3053. Transactions, vol. xli, part 3, and xlv, part 1.
 2510. Proceedings, vol. xxvi, nos. 1-2.
 NATURFORSCHENDE GESELLSCHAFT, FREIBURG I. B.
 2943. Berichte, band xvi, 1906.

- KSL. LEOP. CAROL. DEUTSCHEN AKADEMIE DER
NATURFORSCHER, HALLE
3172-3173. Nova Acta, bande 85-86.
3174. Leopoldina, heft 42.
- COMMISSION GÉOLOGIQUE DE FINLANDE, HELSINGFORS
2841. Bulletins nos. 17-18.
3137. Bulletin no. 20.
- SOCIÉTÉ DE GÉOGRAPHIE DE FINLANDE, HELSINGFORS
3034-3037. Bulletins, Fennia, nos. 19-22.
- SCHWEIZISCHE GEOLOGISCHE GESELLSCHAFT, LAUSANNE
GEOLOGISCH REICHS-MUSEUM, LEIDEN
2074. Sammlungen, neue folge, band i, heft 10. 4to.
- K. SÄCHSISCHE GESELLSCHAFT DER WISSENSCHAFTEN, LEIPSIK
2931. Berichte, Jahrgang 1906, heft 1-8.
2189. Abhandlungen, math. phys. Classe, band xxix, nos. 7-8.
- SOCIÉTÉ GÉOLOGIQUE DE BELGIQUE, LIEGE
2444. Annales, tome xxx, livr. 3, 1903.
2880. Annales, tome xxxiii, livr. 3, 1906.
- SOCIÉTÉ GÉOLOGIQUE DU NORD, LILLE
2890. Annales, tome xxxiv, 1905.
- COMISSÃO DOS TRABALHOS GEOLOGICOS DE PORTUGAL, LISBON
2942. Communicacoes, tomo vi, fasc. 1.
- BRITISH MUSEUM (NATURAL HISTORY), LONDON
GEOLOGICAL SOCIETY, LONDON
2862. Quarterly Journal, vol. lxii. part 4.
3047. Quarterly Journal, vol. lxiii, part 1.
- GEOLOGICAL SURVEY, LONDON
3014. Summary of Progress, 1905.
2981. Memoir. Water Supply of the East Riding of Yorkshire.
3015. Memoir. The Oil Shales of the Lothians.
3054. Memoir, Geology of Falmouth and Cambourne.
- GEOLOGISTS' ASSOCIATION, LONDON
2705. Proceedings, vol. xix, parts 9-10.
- COMISION DEL MAPA GEOLOGICA DE ESPAÑA, MADRID
SOCIETA ITALIANA DI SCIENZE NATURALI, MILAN
2892. Atti, vol. xlv, 1906.
2990. List of Correspondents and General Index to September, 1906.

SOCIÉTÉ IMPÉRIALE DES NATURALISTES DE MOSCOU,

MOSCOW

3001. Bulletin, Année 1905.

2427. Bulletin, Année 1903, nos. 1-3.

K. BAYERISCHE AKADEMIE DER WISSENSCHAFTEN,

MUNICH

2847. Sitzungsberichte, math. phys. Classe, 1905, heft 3.

3018. Sitzungsberichte, math. phys. Classe, 1906, heft 1-2.

ANNALES DES MINES,

PARIS

2997. Annales, 6e série, tome x, livr. 7-12, 1906.

3058. Annales, 6e série, tome v, livr. 1-6, 1907.

CARTE GÉOLOGIQUE DE LA FRANCE,

PARIS

2822. Bulletin, vol. xvi, nos. 106-110.

SOCIÉTÉ GÉOLOGIQUE DE FRANCE,

PARIS

2545. Bulletin, 4e série, tome iv, fasc. 7.

2821. Bulletin, 4e série, tome v, fasc. 6-7.

3070. Bulletin, 4e série, tome vi, fasc. 1.

REALE COMITATO GEOLOGICO D'ITALIA,

ROME

2923. Bolletino, vol. xxxvii, nos. 1-4, 1906.

SOCIETÀ GEOLOGICA ITALIANA,

ROME

3147. Bolletino, vol. xxv, 1906.

ACADÉMIE IMPÉRIALE DES SCIENCES,

SAINT PETERSBURG

3067. Bulletin, 1907.

COMITÉ GÉOLOGIQUE DE LA RUSSIE,

SAINT PETERSBURG

2263. Région aurifère de l'Amour, livr. 5.

2264. Région aurifère de Lena, livr. 3.

2775, 2341. Mémoires, nouvelle série, no. 16.

3039, 2343. Mémoires, nouvelle série, no. 20.

3040. Mémoires, nouvelle série, no. 3.

2773. Bulletin, vol. 23, nos. 7-10.

3139. Carte géologique. Région aurifère de l'Amour, feuille i.

3140. Carte géologique. Région aurifère de la Zela, feuille iii. 2-3.

RÜSSISCH-KAISERLICHE MINERALOGISCHE

GESELLSCHAFT.

SAINT PETERSBURG

2935. Verhandlungen, zweite serie, band xliii.

3038. Materialien zur Geologie Russlands, band xxiii, lief. 1.

GEOLOGISKA BYRÅN.

STOCKHOLM

2278. Sveriges Geol. Undersökning, series Aa, no. 120 and map.

2711, 2992. Sveriges Geol. Undersökning, series Aa, nos. 125-126, 130-133. with maps.

2712. Sveriges Geol. Undersökning, series A, Blad 5 and map.

2993. Sveriges Geol. Undersökning, series C, nos. 197-200.

GEOLOGISKA FÖRENINGENS,

STOCKHOLM

2879. Förhandlingar, band xxxviii, hafte 4-7.
2838. Förhandlingar, band xxix, hafte 1.

NEUES JAHRBUCH FÜR MINERALOGIE.

STUTT GART

2944. Neues Jahrbuch, 1906, band ii.
3064. Neues Jahrbuch, 1907, band i.
2867. Centralblatt, 1906, nos. 14-24.
3051. Centralblatt, 1907, nos. 1-19.

KAISERLICH-KÖNIGLICHE GEOLOGISCHE REICHAN-
STALT,

VIENNA

2902. Jahrbuch, band lvi.

KAISERLICH-KÖNIGLICHES NATURHISTORISCHES
HOFMUSEUM,

VIENNA

2932. Annalen, band xx, 1905.

GEOLOGISCHES INSTITUT DER K. K. UNIVERSITÄT.

VIENNA

- 3025-3029. Beiträge zur Paleontologie und Geologie Österreich-Ungarns und
des Orients, band xiv-xviii.
3030. Same, band xix, heft 1-4.

(c) ASIA

GEOLOGICAL SURVEY OF INDIA.

CALCUTTA

2927. Records, vol. xxxiv, 1906.

IMPERIAL GEOLOGICAL SURVEY,

TOKYO

BUREAU OF SCIENCE,

MANILA

(d) AUSTRALASIA

GEOLOGICAL DEPARTMENT OF SOUTH AUSTRALIA.

ADELAIDE

1456. Contributions to Paleontology, nos. 17-22.
2445. Northern Territory, Reports for 1905.
2527. Review of Mining Operations during 1905-1906.

GEOLOGICAL SURVEY OF QUEENSLAND,

BRISBANE

- 2630, 3011. Publications nos. 201, 203, 205-206.

DEPARTMENT OF MINES OF VICTORIA,

MELBOURNE

2344. Records of the Geological Survey, vol. i, part 4.
3148. Bulletins nos. 19-22.
3152. Memoirs nos. 4-5.
2778. Annual Report of the Secretary of Mines for 1905.
3012. Special Report, Economic Minerals and Rocks.

- GEOLOGICAL DEPARTMENT OF WESTERN AUSTRALIA, PERTH
 2929. Bulletins 21-22.
 2138. Annual Report for 1905.
 2945. Bulletins 23-25.

- GEOLOGICAL SURVEY OF NEW SOUTH WALES, SYDNEY
 2908. Annual Report of the Department of Mines for 1906.
 2979. Memoirs, Paleontology, no. 5, vol. ii, part 1.

- ROYAL SOCIETY OF NEW SOUTH WALES, SYDNEY
 3057. Journal and Proceedings, vol. xxxix, 1905.

- GEOLOGICAL SURVEY OF NEW ZEALAND, WELLINGTON
 2989. Bulletin. new series, nos. 1-2.

(c) AFRICA

- GEOLOGICAL COMMISSION, CAPE TOWN
 3016. Tenth Annual Report. 1905.
 2492. Annals of the South African Museum, vol. iv, part 7.

- GEOLOGICAL SOCIETY OF SOUTH AFRICA, JOHANNESBURG
 2912. Transactions, vol. iv, pp. 1-128, and Proceedings.

- GEOLOGICAL SURVEY OF THE TRANSVAAL, PRETORIA

(B) FROM STATE GEOLOGICAL SURVEYS AND MINING BUREAUS

- GEOLOGICAL SURVEY OF NORTH CAROLINA, CHAPEL HILL
 2982. Vol. i. Corundum and the Peridotites.
 2983-2986. Bulletins 1, 3, 5-11, 13-19.
 2987. Economic Papers 1, 3, 4, 6-10.

- GEOLOGICAL SURVEY OF OHIO, COLUMBUS
 2947-2948. Bulletins, fourth series, nos. 4, 5, and 8.
 3003. Bulletin no. 6, fourth series.

- DEPARTMENT OF THE INTERIOR, OTTAWA
 2788. Guelph sheet, Ontario.

- GEOLOGICAL SURVEY OF NEW JERSEY, TRENTON
 2999. Annual Report for 1905.

(C) FROM SCIENTIFIC SOCIETIES AND INSTITUTIONS

(a) AMERICA

- BROOKLYN INSTITUTE OF ARTS AND SCIENCES, BROOKLYN
 2553. Science Bulletin, vol. i, nos. 9-10.

COLORADO COLLEGE,

COLORADO SPRINGS

2555. Colorado College Studies, Science series, vol. xi, nos. 47-53.

INTERNATIONAL GEOLOGICAL CONGRESS,

MEXICO

3000. Guide Geologique au Mexique.

(b) *EUROPE*

SCHLESISCHE GESELLSCHAFT FÜR VATERLÄNDISCHE
CULTUR,

BRESLAU

2998. Eighty-third Jahresbericht, 1905.

OBSERVATOIRE ROYAL DE BELGIQUE,

BRUSSELS

3245. Annuaire Astronomique, 1907.

3246. Annales, Nouvelle serie, Physique du Globe, tome iii, fasc. 2, 1907.

DANSK GEOLOGISK FORENING,

COPENHAGEN

2783. Meddelelser, nr. 11-12.

ACADEMIE POLYTECHNICA,

PORTO

3181-3182. Annaes Scientificos, vol. i, nos. 3-4, and vol. ii, nos. 1-2.

UNIVERSITY OF UPSALA,

UPSALA

3183-3184. Bulletin of the Geological Institution, vols. vi-vii.

(c) *ASIA*

BUREAU OF SCIENCE,

MANILA

3185. Philippine Journal of Science, vol. i, no. 10.

TOKYO GEOGRAPHICAL SOCIETY,

TOKYO

2946. Journal of Geography, vol. xviii, 1906.

IMPERIAL UNIVERSITY OF TOKYO,

TOKYO

2991. Journal of the College of Science, vol. xxi, articles 2, 5, and 9.

(D) FROM FELLOWS OF THE GEOLOGICAL SOCIETY OF AMERICA (PERSONAL
PUBLICATIONS)

N. H. DARTON

3186. Geology of the Owl Creek Mountains.

U. S. GRANT

3187. Copper and other Mineral Resources of Prince William Sound.

3188. Structural Relations of the Wisconsin Zinc and Lead Deposits.

C. H. HITCHCOCK

3189. The Bible and Recent Science.

GEORGE P. MERRILL

3190. On a newly found Meteorite from Selma, Alabama.
Notes on the Composition and Structure of the Hendersonville,
North Carolina, Meteorite.

(E) FROM MISCELLANEOUS SOURCES

M. GUGENHAN

3191. Der Stuttgarter Talkessel von alpinen Eis ausgehöhlt.

ALBERT HEIM

3194. Über die nordöstlichen Lappen des Tessinermassives.
Die vermeintliche Gewölbeumbiegung des Nordflügels der Glarner-
doppelfalte südlich vom Klausenpass, eine Selbstkorrektur.

MICHEL MOURLON

3192. Le Service Géologique de Belgique.

LENNART VON POST

3193. Norrländska Torfmossestudier. 1.

LEONARDO RICCIARDI

3195. L'Unità delle Energie Cosmiche.

FEDERICO SACCO

3196. Les Lois fondamentales de l'Orogénie de la Terre.
3197. Les Étages et les Faunes du Bassin Tertiaire du Piémont.
3198. Fenomeni di Corrugamento negli Schisti Cristallini delle Alpi.
3199. Sur la Valeur stratigraphique des Lepidocyclina, etc.
4000. Réunion extraordinaire de la Société Géologique de France en
Italie, a Turin et a Gênes.

WIET TASSIN

4001. Note on an Occurrence of Graphitic Iron in a Meteorite.

OFFICERS AND FELLOWS OF THE GEOLOGICAL SOCIETY
OF AMERICA

OFFICERS FOR 1907

President

C. R. VAN HISE, Madison, Wis.

Vice-Presidents

J. S. DILLER, Washington, D. C.

A. P. COLEMAN, Toronto, Canada

Secretary

EDMUND OTIS HOVEY, American Museum of Natural History, New
York, N. Y.

Treasurer

WM. BULLOCK CLARK, Baltimore, Md.

Editor

J. STANLEY-BROWN, Cold Spring Harbor, Long Island, N. Y.

Librarian

H. P. CUSHING, Cleveland, Ohio

Councillors

(Term expires 1907)

H. M. AMI, Ottawa, Canada

J. F. KEMP, New York city

(Term expires 1908)

A. C. LANE, Lansing, Mich.

DAVID WHITE, Washington, D. C.

(Term expires 1909)

H. E. GREGORY, New Haven, Conn.

H. F. REID, Baltimore, Md.

FELLOWS IN DECEMBER, 1907

*Indicates Original Fellow (see article III of Constitution)

- CLEVELAND ABBE, JR., Ph. D., Mount Weather, Va. August, 1899.
- FRANK DAWSON ADAMS, Ph. D., Montreal, Canada; Professor of Geology in McGill University. December, 1889.
- GEORGE I. ADAMS, Sc. D., Corps of Mining Engineers, Lima, Peru. December, 1902.
- JOSÉ GUADALUPE AGUILERA, Director del Instituto Geologico de Mexico, City of Mexico, Mexico. August, 1896.
- TRUMAN H. ALDRICH, M. E., 1739 P St. N. W., Washington, D. C. May, 1889.
- HENRY M. AMI, A. M., Geological Survey Office, Ottawa, Canada; Assistant Paleontologist on Geological and Natural History Survey of Canada. December, 1889.
- FRANK M. ANDERSON, B. A., M. S., 2604 Ætna Street, Berkeley, Cal. California State Mining Bureau. June, 1902.
- PHILIP ARGALL, 728 Majestic Building, Denver, Colo.; Mining Engineer. August, 1896.
- RALPH ARNOLD, Ph. D., Washington, D. C.; Geologic Aid, U. S. Geological Survey. December, 1904.
- GEORGE HALL ASHLEY, M. E., Ph. D., Washington, D. C.; U. S. Geological Survey. August, 1895.
- HARRY FOSTER BAIN, M. S., Champaign, Ill.; State Geologist. December, 1895.
- RUFUS MATHER BAGG, JR., Ph. D., 1048 Riverdale St., West Springfield, Mass.; Mining Geologist. December, 1896.
- S. PRENTISS BALDWIN, 736 Prospect St., Cleveland, Ohio. August, 1895.
- SYDNEY H. BALL, A. B., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1905.
- ERWIN HINCKLEY BARBOUR, Ph. D., Lincoln, Neb.; Professor of Geology, University of Nebraska, and Acting State Geologist. December, 1896.
- ALFRED ERNEST BARLOW, B. A., M. A., D. Sc., Ottawa, Canada; Mining Geologist. December, 1906.
- JOSEPH BARRELL, Ph. D., New Haven, Conn.; Assistant Professor of Geology, Yale University. December, 1902.
- GEORGE H. BARTON, B. S., Boston, Mass.; Curator, Boston Society of Natural History. August, 1890.
- FLORENCE BASCOM, Ph. D., Bryn Mawr, Pa.; Professor of Geology, Bryn Mawr College. August, 1894.
- RAY SMITH BASSLER, B. A., M. S., Ph. D., Washington, D. C., U. S. National Museum. December, 1906.
- WILLIAM S. BAYLEY, Ph. D., Urbana, Ill.; Assistant Professor of Geology, University of Illinois. December, 1888.
- *GEORGE F. BECKER, Ph. D., Washington, D. C., U. S. Geological Survey.
- JOSHUA W. BEEDE, Ph. D., Bloomington, Ind.; Instructor in Geology, Indiana University. December, 1902.
- ROBERT BELL, C. E., M. D., LL. D., Ottawa, Canada; Acting Director of the Geological and Natural History Survey of Canada. May, 1889.
- CHARLES P. BERKEY, Ph. D., New York city; Columbia University. August, 1901.
- SAMUEL WALKER BEYER, Ph. D., Ames, Iowa; Assistant Professor in Geology, Iowa Agricultural College. December, 1896.
- ARTHUR B. BIBBINS, Ph. B., Baltimore, Md.; Instructor in Geology, Woman's College. December, 1903.
- ALBERT S. BICKMORE, Ph. D., American Museum of Natural History, New York; Curator emeritus, Department of Public Instruction. December, 1889.

- IRVING P. BISHOP, 109 Norwood Ave., Buffalo, N. Y.; Professor of Natural Science, State Normal and Training School. December, 1899.
- JOHN M. BOUTWELL, M. S., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1905.
- JOHN ADAMS BOWNOCKER, D. Sc., Columbus, Ohio.; Professor of Inorganic Geology, Ohio State University. December, 1904.
- *JOHN C. BRANNER, Ph. D., Stanford University, Cal.; Professor of Geology in Leland Stanford, Jr., University.
- ALBERT PERRY BRIGHAM, A. B., A. M., Hamilton, N. Y.; Professor of Geology and Natural History, Colgate University. December, 1893.
- REGINALD W. BROCK, M. A., Ottawa, Canada; Geologist, Geological and Natural History Survey of Canada; Professor of Geology, School of Mining, Kingston. December, 1904.
- ALFRED HULSE BROOKS, B. S., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. August, 1899.
- AMOS P. BROWN, Ph. D., Philadelphia, Pa.; Professor of Mineralogy and Geology, University of Pennsylvania. December, 1905.
- ERNEST ROBERTSON BUCKLEY, Ph. D., Rolla, Mo.; State Geologist and Director of Bureau of Geology and Mines. June, 1902.
- *SAMUEL CALVIN, Iowa City, Iowa; Professor of Geology and Zoology in the State University of Iowa.
- HENRY DONALD CAMPBELL, Ph. D., Lexington, Va.; Professor of Geology and Biology in Washington and Lee University. May, 1889.
- MARIUS R. CAMPBELL, U. S. Geological Survey, Washington, D. C. August, 1892.
- FRANKLIN R. CARPENTER, Ph. D., 1420 Josephine St., Denver, Colo.; Mining Engineer. May, 1889.
- ERMINE C. CASE, Ph. D., Department of Geology, University of Michigan, Ann Arbor, Mich. December, 1901.
- *T. C. CHAMBERLIN, LL. D., Chicago, Ill.; Head Professor of Geology, University of Chicago.
- CLARENCE RAYMOND CLAGHORN, B. S., M. E., Tacoma, Wash. August, 1891.
- FREDERICK G. CLAPP, S. B., Washington, D. C.; Geologic Aid, U. S. Geological Survey. December, 1905.
- *WILLIAM BULLOCK CLARK, Ph. D., Baltimore, Md.; Professor of Geology in Johns Hopkins University; State Geologist.
- JOHN MASON CLARKE, A. M., Albany, N. Y.; State Paleontologist. December, 1897.
- HERDMAN F. CLELAND, Ph. D., Williamstown, Mass.; Professor of Geology, Williams College. December, 1905.
- J. MORGAN CLEMENTS, Ph. D., 15 William St., New York city. December, 1894.
- COLLIER COBB, A. B., A. M., Chapel Hill, N. C.; Professor of Geology in University of North Carolina. December, 1894.
- ARTHUR P. COLEMAN, Ph. D., Toronto, Canada; Professor of Geology, Toronto University, and Geologist of Bureau of Mines of Ontario. December, 1896.
- GEORGE L. COLLIE, Ph. D., Beloit, Wis.; Professor of Geology in Beloit College. December, 1897.
- ARTHUR J. COLLIER, A. M., S. B., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. June, 1902.
- *THEODORE B. COMSTOCK, Sc. D., Los Angeles, Cal.; Mining Engineer.
- EUGENE COSTE, B. ès-Sc., E. M., Toronto, Canada. December, 1906.
- *FRANCIS W. CRAGIN, Ph. D., Colorado Springs, Colo.; Professor of Geology in Colorado College.
- ALJA ROBINSON CROOK, Ph. D., Springfield, Ill.; State Museum of Natural History. December, 1898.
- *WILLIAM O. CROSBY, B. S., Boston Society of Natural History, Boston, Mass.; Assistant Professor of Mineralogy and Lithology in Massachusetts Institute of Technology.

- WHITMAN CROSS, Ph. D., U. S. Geological Survey, Washington, D. C. May, 1889.
- GARRY E. CULVER, A. M., 1104 Wisconsin St., Stevens Point, Wis. December, 1891.
- EDGAR R. CUMINGS, Ph. D., Bloomington, Ind.; Assistant Professor of Geology, Indiana University. August, 1901.
- *HENRY P. CUSHING, M. S., Adelbert College, Cleveland, Ohio; Professor of Geology, Western Reserve University.
- REGINALD A. DALY, Ph. D., Ottawa, Canada; Geologist for Canada on International Boundary Commission. December, 1905.
- *NELSON H. DARTON, United States Geological Survey, Washington, D. C.
- *WILLIAM M. DAVIS, S. B., M. E., Cambridge, Mass.; Sturgis-Hooper Professor of Geology in Harvard University.
- DAVID T. DAY, Ph. D., U. S. Geological Survey, Washington, D. C. August, 1891.
- ORVILLE A. DERBY, M. S., Sao Paulo, Brazil; No. 80 Rua Visconde do Rio Branco. December, 1890.
- *JOSEPH S. DILLER, B. S., U. S. Geological Survey, Washington, D. C.
- EDWARD V. D'INVILLIERS, E. M., 506 Walnut St., Philadelphia, Pa. Dec., 1888.
- RICHARD E. DODGE, A. M., Teachers' College, West 120th St., New York city; Professor of Geography in the Teachers' College. August, 1897.
- NOAH FIELDS DRAKE, Ph. D., Tientsin, China; Professor of Geology in Imperial Tientsin University. December, 1898.
- JOHN ALEXANDER DRESSER, B. A., M. A., Montreal, Canada; Principal of Prince Albert School. December, 1906.
- CHARLES R. DRYER, M. A., M. D., Terre Haute, Ind.; Professor of Geography, Indiana State Normal School. August, 1897.
- *EDWIN T. DUMBLE, 1306 Main St., Houston, Texas.
- ARTHUR S. EAKLE, Ph. D., Berkeley, Cal.; Instructor in Mineralogy, University of California. December, 1899.
- CHARLES R. EASTMAN, A. M., Ph. D., Cambridge, Mass.; In Charge of Vetebrate Paleontology, Museum of Comparative Zoology, Harvard University. December, 1895.
- EDWIN C. ECKEL, B. S., C. E., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1905.
- ARTHUR H. ELEFTMAN, Ph. D., 706 Globe Building, Minneapolis, Minn. December, 1898.
- *BENJAMIN K. EMERSON, Ph. D., Amherst, Mass.; Professor in Amherst College.
- *SAMUEL F. EMMONS, A. M., E. M., U. S. Geological Survey, Washington, D. C.
- JOHN EYERMAN, F. Z. S., Oakhurst, Easton, Pa. August, 1891.
- HAROLD W. FAIRBANKS, B. S., Berkeley, Cal.; Geologist State Mining Bureau. August, 1892.
- *HERMAN L. FAIRCHILD, B. S., Rochester, N. Y.; Professor of Geology in University of Rochester.
- J. C. FALES, Danville, Ky.; Professor in Centre College. December, 1888.
- OLIVER C. FARRINGTON, Ph. D., Chicago, Ill.; In charge of Department of Geology, Field Columbian Museum. December, 1895.
- NEVIN M. FENNEMAN, Ph. D., Madison, Wis.; Professor of Geology, University of Wisconsin. December, 1904.
- AUGUST F. FOERSTE, Ph. D., 417 Grand Ave., Dayton, Ohio; Teacher of Sciences. December, 1899.
- WILLIAM M. FONTAINE, A. M., University of Virginia, Va.; Professor of Natural History and Geology in University of Virginia. December, 1888.
- *PERSIFOR FRAZER, D. ès-Sc. Nat., 1082 Drexel Building, Philadelphia, Pa.; Professor of Chemistry in Horticultural Society of Pennsylvania.
- *HOMER T. FULLER, Ph. D., Fredonia, N. Y.

- MYBON LESLIE FULLER, S. B., U. S. Geological Survey, Washington, D. C. December, 1898.
- HENRY STEWART GANE, Ph. D., Santa Barbara, Cal. December, 1896.
- HENRY GANNETT, S. B., A. Met. B., U. S. Geological Survey, Washington, D. C. December, 1891.
- RUSSELL D. GEORGE, A. B., A. M., Boulder, Colo.; Professor of Geology, University of Colorado. December, 1906.
- *GROVE K. GILBERT, A. M., LL. D., U. S. Geological Survey, Washington, D. C.
- ADAM CAPEN GILL, Ph. D., Ithaca, N. Y.; Assistant Professor of Mineralogy and Petrography in Cornell University. December, 1888.
- L. C. GLENN, Ph. D., Nashville, Tenn.; Professor of Geology in Vanderbilt University. June, 1900.
- CHARLES H. GORDON, Ph. D., University of Tennessee, Knoxville, Tenn. August, 1893.
- CHARLES NEWTON GOULD, A. M., Norman, Okla.; Professor of Geology, University of Oklahoma. December, 1904.
- AMADEUS W. GRABAU, S. M., S. D., New York city; Professor of Paleontology, Columbia University. December, 1898.
- ULYSSES SHERMAN GRANT, Ph. D., Evanston, Ill.; Professor of Geology, Northwestern University. December, 1890.
- HERBERT E. GREGORY, Ph. D., New Haven, Conn.; Silliman Professor of Geology, Yale University. August, 1901.
- GEORGE P. GRIMSLEY, Ph. D., Morgantown, W. Va.; Assistant State Geologist, Geological Survey of West Virginia. August, 1895.
- LEON S. GRISWOLD, A. B., Rolla, Missouri. August, 1902.
- FREDERIC P. GULLIVER, Ph. D., Norwichtown, Conn. August, 1895.
- ARNOLD HAGUE, Ph. B., U. S. Geological Survey, Washington, D. C. May, 1889.
- *CHRISTOPHER W. HALL, A. M., 803 University Ave., Minneapolis, Minn.; Professor of Geology and Mineralogy in University of Minnesota.
- GILBERT D. HARRIS, Ph. B., Ithaca, N. Y.; Assistant Professor of Paleontology and Stratigraphic Geology, Cornell University. December, 1903.
- JOHN BURCHMORE HARRISON, M. A., F. I. C., F. G. S., Georgetown, British Guiana; Government Geologist. June, 1902.
- JOHN B. HASTINGS, M. E., 1480 High St., Denver, Colo. May, 1889.
- *ERASMUS HAWORTH, Ph. D., Lawrence, Kans.; Professor of Geology, University of Kansas.
- C. WILLARD HAYES, Ph. D., U. S. Geological Survey, Washington, D. C. May, 1889.
- RICHARD R. HICE, B. S., Beaver, Pa. December, 1903.
- *EUGENE W. HILGARD, Ph. D., LL. D.; Berkeley, Cal.; Professor of Agriculture in University of California.
- FRANK A. HILL, Roanoke, Va. May, 1889.
- *ROBERT T. HILL, B. S., 25 Broad St., New York city.
- RICHARD C. HILLS, Mining Engineer, Denver, Colo. August, 1894.
- *CHARLES H. HITCHCOCK, Ph. D., LL. D., Hanover, N. H.; Professor of Geology in Dartmouth College.
- WILLIAM HERBERT HOBBS, Ph. D., Ann Arbor, Mich.; Professor of Geology, University of Michigan; Assistant Geologist, U. S. Geological Survey. August, 1891.
- *LEVI HOLBROOK, A. M., P. O. Box 536, New York city.
- ARTHUR HOLLICK, Ph. B., Bronx Park, New York; Assistant Curator, Department of Fossil Botany, New York Botanical Garden. August, 1893.
- *JOSEPH A. HOLMES, 6017 Cabanne Ave., Saint Louis, Mo.; State Geologist of North Carolina; In charge of investigation of fuels and structural materials, U. S. Geological Survey.

THOMAS C. HOPKINS, Ph. D., Syracuse, N. Y.; Professor of Geology, Syracuse University. December, 1894.

*EDMUND OTIS HOVEY, Ph. D., American Museum of Natural History, New York city; Associate Curator of Geology.

*HORACE C. HOVEY, D. D., Newburyport, Mass.

ERNEST HOWE, Ph. D., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1903.

*EDWIN E. HOWELL, A. M., 612 Seventeenth St. N. W., Washington, D. C.

LUCIUS L. HUBBARD, Ph. D., LL. D., Houghton, Mich. December, 1894.

ELLSWORTH HUNTINGTON, A. B., A. M., Milton, Mass. December, 1906.

JOSEPH P. IDDINGS, Ph. B., Professor of Petrographic Geology, University of Chicago, Chicago, Ill. May, 1889.

JOHN D. IRVING, Ph. D., South Bethlehem, Pa.; Professor of Geology, Lehigh University. December, 1905.

A. WENDELL JACKSON, Ph. B., 432 Saint Nicholas Ave., New York city. December, 1888.

ROBERT T. JACKSON, S. D., 9 Fayerweather St., Cambridge, Mass.; Assistant Professor in Paleontology in Harvard University. August, 1894.

THOMAS M. JACKSON, C. E., S. D., Clarksburg, W. Va. May, 1889.

THOMAS AGUSTUS JAGGAR, JR., A. B., A. M., Ph. D., Cambridge, Mass.; Professor of Geology, Massachusetts Institution of Technology, Boston. December, 1906.

MARK S. W. JEFFERSON, A. M., Ypsilanti, Mich.; Professor of Geography, Michigan State Normal College. December, 1904.

DOUGLAS WILSON JOHNSON, B. S., Ph. D., Cambridge, Mass.; Assistant Professor of Physiography, Harvard University. December, 1906.

ALEXIS A. JULIEN, Ph. D., Columbia College, New York city; Instructor in Columbia College. May, 1889.

ARTHUR KEITH, A. M., U. S. Geological Survey, Washington, D. C. May, 1889.

*JAMES F. KEMP, A. B., E. M., Columbia University, New York city; Professor of Geology.

CHARLES ROLLIN KEYES, Ph. D., 944 Fifth St., Des Moines, Iowa. August, 1890.

EDWARD M. KINDLE, Ph. D., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1905.

FRANK H. KNOWLTON, M. S., Washington, D. C.; Assistant Paleontologist, U. S. Geological Survey. May, 1889.

EDWARD HENRY KRAUS, Ph. D., Ann Arbor, Mich.; Junior Professor of Mineralogy, University of Michigan. June, 1902.

HENRY B. KUMMEL, Ph. D., Trenton, N. J.; State Geologist. December, 1895.

*GEORGE F. KUNZ, A. M. (Hon.), Ph. D. (Hon.), care of Tiffany & Co., Fifth avenue, at 37th street, New York city.

GEORGE EDGAR LADD, Ph. D., Rolla, Mo.; Director School of Mines. August, 1891.

J. C. K. LAFLAMME, M. A., D. D., Quebec, Canada; Professor of Mineralogy and Geology in University Laval, Quebec. August, 1890.

ALFRED C. LANE, Ph. D., Lansing, Mich.; State Geologist of Michigan. December, 1889.

DANIEL W. LANGTON, Ph. D., Fuller Building, New York city; Mining Engineer. December, 1889.

ANDREW C. LAWSON, Ph. D., Berkeley, Cal.; Professor of Geology and Mineralogy in the University of California. May, 1889.

WILLIS THOMAS LEE, M. S., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1903.

CHARLES K. LEITH, Ph. D., Madison, Wis.; Professor of Geology, University of Wisconsin; Assistant Geologist, U. S. Geological Survey. December, 1902.

- ARTHUR G. LEONARD, Ph. D., Grand Forks, N. Dak.; Professor of Geology and State Geologist, State University of North Dakota. December, 1901.
- FRANK LEVERETT, B. S., Ann Arbor, Mich.; Geologist, U. S. Geological Survey. August, 1890.
- JOSEPH VOLNEY LEWIS, B. E., S. B., New Brunswick, N. J.; Professor of Geology, Rutgers College. December, 1906.
- WILLIAM LIBBEY, Sc. D., Princeton, N. J.; Professor of Physical Geography in Princeton University. August, 1899.
- WALDEMAR LINDGREN, M. E., U. S. Geological Survey, Washington, D. C. August, 1890.
- GEORGE DAVIS LOUDERBACK, Ph. D., Associate Professor of Geology, University of California, Berkeley, Cal. June, 1902.
- ROBERT H. LOUGHBRIDGE, Ph. D., Berkeley, Cal.; Assistant Professor of Agricultural Chemistry in University of California. May, 1889.
- ALBERT P. LOW, B. S., Ottawa, Canada; Geologist, Geological Survey of Canada. December, 1905.
- THOMAS H. MACBRIDE, A. M., Iowa City, Iowa; Professor of Botany in the State University of Iowa. May, 1889.
- HIRAM DEYER McCASKEY, B. S., U. S. Geological Survey, Washington, D. C. December, 1904.
- RICHARD G. McCONNELL, A. B., Geological Survey Office, Ottawa, Canada; Geologist on Geological and Natural History Survey of Canada. May, 1889.
- JAMES RIEMAN MACFARLANE, A. B., 100 Diamond St., Pittsburg, Pa. August 1891.
- *W J MCGEE, LL. D., Director Public Museum, Saint Louis, Mo.
- WILLIAM MCINNIS, A. B., Geological Survey Office, Ottawa, Canada; Geologist, Geological and Natural History Survey of Canada. May, 1889.
- PETER MCKELLAR, Fort William, Ontario, Canada. August, 1890.
- CURTIS F. MARBUT, A. M., State University, Columbia, Mo.; Instructor in Geology and Assistant on Missouri Geological Survey. August, 1897.
- VERNON F. MARSTERS, A. M., Lima, Peru. August, 1892.
- GEORGE CURTIS MARTIN, Ph. D., Washington, D. C., U. S. Geological Survey. June, 1902.
- EDWARD B. MATHEWS, Ph. D., Baltimore, Md.; Instructor in Petrography in Johns Hopkins University. August, 1895.
- W. D. MATTHEW, Ph. D., New York city; Associate Curator in Vertebrate Paleontology, American Museum of Natural History. December, 1903.
- P. H. MELL, M. E., Ph. D., Clemson College, S. C.; President of Clemson College. December, 1888.
- WARREN C. MENDENHALL, B. S., 1108 Braly Building, Los Angeles, Cal.; Geologist, U. S. Geological Survey. June, 1902.
- JOHN C. MERRIAM, Ph. D., Berkeley, Cal.; Instructor in Paleontology in University of California. August, 1895.
- *FREDERICK J. H. MERRILL, Ph. D., New Rochelle, N. Y.; Consulting Geologist.
- GEORGE P. MERRILL, Ph. D., U. S. National Museum, Washington, D. C.; Curator of Department of Lithology and Physical Geology. December, 1888.
- ARTHUR M. MILLER, A. M., Lexington, Ky.; Professor of Geology, State University of Kentucky. December, 1897.
- BENJAMIN L. MILLER, Ph. D., South Bethlehem, Pa.; Professor of Geology, Lehigh University. December, 1904.
- WILLET G. MILLER, M. A., Toronto, Canada; Provincial Geologist of Ontario. December, 1902.
- HENRY MONTGOMERY, Ph. D., Toronto, Canada; Curator of Museum, University of Toronto. December, 1904.

*FRANK L. NASON, A. B., West Haven, Conn.

DAVID HALE NEWLAND, B. A., Albany, N. Y.; Assistant State Geologist. December, 1906.

JOHN F. NEWSOM, Ph. D., Stanford University, Cal.; Associate Professor of Mining. December, 1899.

WILLIAM H. NILES, Ph. B., M. A., Boston, Mass.; Professor, Emeritus, of Geology, Massachusetts Institute of Technology; Professor of Geology, Wellesley College. August, 1891.

WILLIAM H. NORTON, M. A., Mount Vernon, Iowa; Professor of Geology in Cornell College. December, 1895.

CHARLES J. NORWOOD, Lexington, Ky.; Professor of Mining, State College of Kentucky. August, 1894.

IDA HELEN OGILVIE, A. B., Ph. D., New York city; Tutor in Geology, Barnard College, Columbia University. December, 1906.

CLEOPHAS C. O'HARRA, Ph. D., Rapid City, S. Dak.; Professor of Mineralogy and Geology, South Dakota School of Mines. December, 1904.

EZEQUIEL ORDONEZ, 2 a General Priue, Mexico, D. F., Mex. August, 1896.

*AMOS O. OSBORN, Waterville, Oneida county, N. Y.

HENRY F. OSBORN, Sc. D., Columbia University, New York city; Professor of Zoology, Columbia University. August, 1894.

CHARLES PALACHE, B. S., University Museum, Cambridge, Mass.; Instructor in Mineralogy, Harvard University. August, 1897.

WILLIAM A. PARKS, B. A., Ph. D., Toronto, Canada; Associate Professor of Geology, University of Toronto. December, 1906.

*HORACE B. PATTON, Ph. D., Golden, Colo.; Professor of Geology and Mineralogy in Colorado School of Mines.

FREDERICK B. PECK, Ph. D., Easton, Pa.; Professor of Geology and Mineralogy, Lafayette College. August, 1901.

RICHARD A. F. PENROSE, JR., Ph. D., 460 Bullitt Building, Philadelphia, Pa. May, 1889.

GEORGE H. PERKINS, Ph. D., Burlington, Vt.; State Geologist; Professor of Geology, University of Vermont. June, 1902.

JOSEPH H. PERRY, 276 Highland St., Worcester, Mass. December, 1888.

LOUIS V. PIRSSON, Ph. D., New Haven, Conn.; Professor of Physical Geology, Sheffield Scientific School of Yale University. August, 1894.

*JULIUS POHLMAN, M. D., University of Buffalo, Buffalo, N. Y.

JOSEPH HYDE PRATT, Ph. D., Chapel Hill, N. C.; Mineralogist, North Carolina Geological Survey. December, 1898.

*CHARLES S. PROSSER, M. S., Columbus, Ohio; Professor of Geology in Ohio State University.

*RAPHAEL PUMPELLY, U. S. Geological Survey, Dublin, N. H.

ALBERT HOMER PURDUE, B. A., Fayetteville, Ark.; Professor of Geology, University of Arkansas. December, 1904.

FREDERICK LESLIE RANSOME, Ph. D., Washington, D. C.; Geologist, U. S. Geological Survey. August, 1895.

HARRY FIELDING REID, Ph. D., Baltimore, Md.; Professor of Geological Physics, Johns Hopkins University. December, 1892.

WILLIAM NORTH RICE, Ph. D., LL. D., Middletown, Conn.; Professor of Geology in Wesleyan University. August, 1890.

CHARLES H. RICHARDSON, Ph. D., Syracuse, N. Y.; Assistant Professor of Geology and Mineralogy, Syracuse University. December, 1899.

HEINRICH RIES, Ph. D., Cornell University, Ithaca, N. Y.; Professor of Economic Geology. December, 1893.

RUDOLPH RUEDEMANN, Ph. D., Albany, N. Y.; Assistant State Paleontologist. December, 1905.

- ORESTES H. ST. JOHN, Raton, N. Mex. May, 1889.
- *ROLLIN D. SALISBURY, A. M., Chicago, Ill.; Professor of General and Geographic Geology in University of Chicago.
- FREDERICK W. SARDESON, Ph. D., Assistant Professor of Geology, University of Minnesota, Minneapolis, Minn. December, 1892.
- FRANK C. SCHRADER, M. S., A. M., U. S. Geological Survey, Washington, D. C. August, 1901.
- CHARLES SCHUCHERT, New Haven, Conn.; Professor of Paleontology, Yale University. August, 1895.
- WILLIAM B. SCOTT, Ph. D., 56 Bayard Ave., Princeton, N. J.; Blair Professor of Geology in College of New Jersey. August, 1892.
- ARTHUR EDMUND SEAMAN, B. S., Houghton, Mich.; Professor of Mineralogy and Geology, Michigan College of Mines. December, 1904.
- HENRY M. SEELY, M. D., Middlebury, Vt.; Professor of Geology in Middlebury College. May, 1899.
- ELIAS H. SELLARDS, Ph. D., Tallahassee, Fla.; State Geologist. December, 1905.
- GEORGE BURBANK SHATTUCK, Ph. D., Poughkeepsie, N. Y.; Professor of Geology in Vassar College. August, 1899.
- SOLON SHEDD, A. B., Pullman, Wash.; Professor of Geology and Mineralogy, Washington Agricultural College. December, 1904.
- EDWARD M. SHEPARD, Sc. D., Springfield, Mo.; Professor of Geology, Drury College. August, 1901.
- WILL H. SHERZER, M. S., Ypsilanti, Mich.; Professor in State Normal School. December, 1890.
- BOHUMIL SHIMEK, C. E., M. S., Iowa City, Iowa; Professor of Physiological Botany, University of Iowa. December, 1904.
- *FREDERICK W. SIMONDS, Ph. D., Austin, Texas; Professor of Geology in University of Texas.
- WILLIAM JOHN SINCLAIR, B. S., Ph. D., Princeton, N. J.; Instructor in Princeton University. December, 1906.
- *EUGENE A. SMITH, Ph. D., University, Tuscaloosa county, Ala.; State Geologist and Professor of Chemistry and Geology in University of Alabama.
- FRANK CLEMES SMITH, E. M., Richland Center, Wis.; Mining Engineer. December, 1898.
- GEORGE OTIS SMITH, Ph. D., Washington, D. C.; Director, U. S. Geological Survey. August, 1897.
- WILLIAM S. T. SMITH, Ph. D., 749 N. Lake St., Reno, Nev.; Associate Professor of Geology and Mineralogy, University of Nevada. June, 1902.
- *JOHN C. SMOCK, Ph. D., Trenton, N. J.
- CHARLES H. SMYTH, JR., Ph. D., Clinton, N. Y.; Professor of Geology in Hamilton College. August, 1892.
- HENRY L. SMYTH, A. B., Cambridge, Mass.; Professor of Mining and Metallurgy in Harvard University. August, 1894.
- ARTHUR COE SPENCER, B. S., Ph. D., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1896.
- *J. W. SPENCER, Ph. D., 2019 Hillyer Place, Washington, D. C.
- JOSIAH E. SPURR, A. B., A. M., U. S. Geological Survey, Washington, D. C. December, 1894.
- JOSEPH STANLEY-BROWN, Cold Spring Harbor, Long Island, N. Y. August, 1892.
- TIMOTHY WILLIAM STANTON, B. S., U. S. National Museum, Washington, D. C.; Assistant Paleontologist, U. S. Geological Survey. August, 1891.
- *JOHN J. STEVENSON, Ph. D., LL. D., New York University; Professor of Geology in the New York University.

- WILLIAM J. SUTTON, B. S., E. M., Victoria, B. C.; Geologist to E. and N. Railway Co. August, 1901.
- JOSEPH A. TAFF, B. S., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. August, 1895.
- JAMES E. TALMAGE, Ph. D., Salt Lake City, Utah; Professor of Geology in University of Utah. December, 1897.
- RALPH S. TARR, Cornell University, Ithaca, N. Y.; Professor of Dynamic Geology and Physical Geography. August, 1890.
- FRANK B. TAYLOR, Fort Wayne, Ind. December, 1895.
- WILLIAM G. TIGHT, M. S., Albuquerque, N. Mex.; President and Professor of Geology, University of New Mexico. August, 1897.
- *JAMES E. TODD, A. M., 1000 Illinois St., Lawrence, Kas.; Assistant Geologist, U. S. Geological Survey.
- *HENRY W. TURNER, B. S., 508 California St., San Francisco, Cal.
- JOSEPH B. TYRRELL, M. A., B. Sc., 9 Toronto St., Toronto, Canada. May, 1880.
- JOHAN A. UDDEN, A. M., Rock Island, Ill.; Professor of Geology and Natural History in Augustana College. August, 1897.
- EDWARD O. ULBICH, D. Sc., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1903.
- *WARREN UPHAM, A. M., Librarian Minnesota Historical Society, Saint Paul, Minn.
- *CHARLES R. VAN HISE, M. S., Ph. D., Madison, Wis.; President University of Wisconsin; Geologist, U. S. Geological Survey.
- FRANK ROBERTSON VAN HORN, Ph. D., Cleveland, Ohio; Professor of Geology and Mineralogy, Case School of Applied Science. December, 1898.
- GILBERT VAN INGEN, Princeton, N. J.; Curator of Invertebrate Paleontology and Assistant in Geology, Princeton University. December, 1904.
- THOMAS WAYLAND VAUGHN, B. S., A. M., Washington, D. C.; Assistant Geologist, U. S. Geological Survey. August, 1896.
- ARTHUR CLIFFORD VEACH, Washington, D. C.; Geologist, U. S. Geological Survey. December, 1906.
- *ANTHONY W. VODGES, San Diego, Cal.; Captain Fifth Artillery, U. S. Army.
- *M. EDWARD WADSWORTH, Ph. D., State College, Pa.; State Geologist, Pa.; Dean of the School of Mines and Metallurgy and Professor of Mining and Geology in the Pennsylvania State College, and Dean of the School of Mines and Professor of Mining and Geology in the Western University of Pennsylvania, Pittsburg, Pa.
- *CHARLES D. WALCOTT, LL. D., Washington, D. C.; Secretary Smithsonian Institution.
- THOMAS L. WALKER, Ph. D., Toronto, Canada; Professor of Mineralogy and Petrography, University of Toronto. December, 1903.
- CHARLES H. WARREN, Ph. D., Boston, Mass.; Instructor in Geology, Massachusetts Institute of Technology. December, 1901.
- HENRY STEPHENS WASHINGTON, Ph. D., Locust, Monmouth Co., N. J.; August, 1896.
- THOMAS L. WATSON, Ph. D., Charlottesville, Va.; Professor of Geology in University of Virginia. June, 1900.
- WALTER H. WEED, E. M., Norwalk, Conn. May, 1889.
- FRED. BOUGHTON WEEKS, Washington, D. C.; Assistant Geologist, U. S. Geological Survey. December, 1903.
- SAMUEL WEIDMAN, Ph. D., Madison, Wis.; Geologist, Wisconsin Geological and Natural History Survey. December, 1903.
- STUART WELLER, B. S., Chicago, Ill.; Assistant Professor of Paleontologic Geology, University of Chicago. June, 1900.
- LEWIS G. WESTGATE, Ph. D., Delaware, Ohio; Professor of Geology, Ohio Wesleyan University.

- THOMAS C. WESTON, 591 Saint John St., Quebec, Canada. August, 1893.
- DAVID WHITE, B. S., U. S. National Museum, Washington, D. C.; Geologist, U. S. Geological Survey, Washington, D. C. May, 1889.
- *ISRAEL C. WHITE, Ph. D., Morgantown, W. Va.
- *ROBERT P. WHITFIELD, A. M., American Museum of Natural History, 77th St. and 8th Ave., New York city; Curator of Geology and Paleontology.
- FRANK A. WILDER, Ph. D., North Holston, Smyth Co., Va. December, 1905.
- *EDWARD H. WILLIAMS, JR., A. C., E. M., Andover, Mass.
- *HENRY S. WILLIAMS, Ph. D., Ithaca, N. Y.; Professor of Geology and Head of Geological Department, Cornell University.
- IRA A. WILLIAMS, M. Sc., Ames, Iowa; Teacher Iowa State College. December, 1905.
- BAILEY WILLIS, U. S. Geological Survey, Washington, D. C. December, 1889.
- SAMUEL W. WILLISTON, Ph. D., M. D., Chicago, Ill.; Professor of Paleontology, University of Chicago. December, 1889.
- ARTHUR B. WILLMOTT, M. A., Sault Ste. Marie, Ontario, Canada. December, 1899.
- ALFRED W. G. WILSON, Ph. D., 197 Park Ave., Montreal, Ontario, Canada; Mining Geologist. June, 1902.
- ALEXANDER N. WINCHELL, Doct. U. Paris, Madison, Wis.; Professor of Geology and Mineralogy, University of Wisconsin. August, 1901.
- *HORACE VAUGHN WINCHELL, Saint Paul, Minn.; Geologist for the Great Northern Railway Company.
- *NEWTON H. WINCHELL, A. M., Minneapolis, Minn.
- *ARTHUR WINSLOW, B. S., 84 State St., Boston, Mass.
- JOHN E. WOLFF, Ph. D., Harvard University, Cambridge, Mass.; Professor of Petrography and Mineralogy in Harvard University and Curator of the Mineralogical Museum. December, 1889.
- JOSEPH E. WOODMAN, S. D., Halifax, N. S.; Assistant Professor of Geology and Mineralogy, Dalhousie University. December, 1905.
- ROBERT S. WOODWARD, C. E., Washington, D. C.; President of the Carnegie Institution of Washington. May, 1889.
- JAY B. WOODWORTH, B. S., 24 Langdon St., Cambridge, Mass.; Assistant Professor of Geology, Harvard University. December, 1895.
- FREDERIC E. WRIGHT, Ph. D., U. S. Geological Survey, Washington, D. C. December, 1903.
- *G. FREDERICK WRIGHT, D. D., Oberlin, Ohio; Professor in Oberlin Theological Seminary.
- WILLIAM S. YEATES, A. B., A. M., Atlanta, Ga.; State Geologist of Georgia. August, 1894.
- GEORGE A. YOUNG, Ph. D., Ottawa, Canada; Geologist, Geological Survey of Canada. December, 1905.

FELLOWS DECEASED

*Indicates Original Fellow (see article III of Constitution)

- *CHARLES A. ASHBURNER, M. S., C. E. Died December 24, 1889.
- CHARLES E. BEECHER, Ph. D. Died February 14, 1904.
- AMOS BOWMAN. Died June 18, 1894.
- *J. H. CHAPIN, Ph. D. Died March 14, 1892.
- *EDWARD W. CLAYPOLE, D. Sc. Died August 17, 1901.
- GEORGE H. COOK, Ph. D., LL. D. Died September 22, 1889.
- *EDWARD D. COPE, Ph. D. Died April 12, 1897.

- ANTONIO DEL CASTILLO. Died October 28, 1895.
 *JAMES D. DANA, LL. D. Died April 14, 1895.
 GEORGE M. DAWSON, D. Sc. Died March 2, 1901.
 Sir J. WILLIAM DAWSON, LL. D. Died November 19, 1899.
 *WILLIAM B. DWIGHT, Ph. B. Died August 29, 1906.
 *GEORGE H. ELDRIDGE, A. B. Died June 29, 1905.
 *ALBERT E. FOOTE. Died October 10, 1895.
 N. J. GIROUX, C. E. Died November 30, 1890.
 *JAMES HALL, LL. D. Died August 7, 1898.
 JOHN B. HATCHER, Ph. B. Died July 3, 1904.
 *ROBERT HAY. Died December 14, 1895.
 *ANGELO HEILPRIN. Died July 17, 1907.
 DAVID HONEYMAN, D. C. L. Died October 17, 1889.
 THOMAS STERRY HUNT, D. Sc., LL. D. Died February 12, 1892.
 *ALPHEUS HYATT, B. S. Died January 15, 1902.
 *JOSEPH F. JAMES, M. S. Died March 29, 1897.
 WILBUR C. KNIGHT, B. S., A. M. Died July 28, 1903.
 RALPH D. LACOE. Died February 5, 1901.
 *JOSEPH LE CONTE, M. D., LL. D. Died July 6, 1901.
 *J. PETER LESLEY, LL. D. Died June 2, 1903.
 HENRY MCCALLEY, A. M., C. E. Died November 20, 1904.
 OLIVER MARGY, LL. D. Died March 19, 1899.
 OTHNIEL C. MARSH, Ph. D., LL. D. Died March 18, 1899.
 JAMES E. MILLS, B. S. Died July 25, 1901.
 *HENRY B. NASON, M. D., Ph. D., LL. D. Died January 17, 1895.
 *PETER NEFF, M. A. Died May 11, 1903.
 *JOHN S. NEWBERRY, M. D., LL. D. Died December 7, 1892.
 *EDWARD ORTON, Ph. D., LL. D. Died October 16, 1899.
 *RICHARD OWEN, LL. D. Died March 24, 1890.
 SAMUEL L. PENFIELD. Died August 14, 1906.
 *FRANKLIN PLATT. Died July 24, 1900.
 *WILLIAM H. PETTEE, A. M. Died May 26, 1904.
 *JOHN WESLEY POWELL, LL. D. Died September 23, 1902.
 *ISRAEL C. RUSSELL, LL. D. Died May 1, 1906.
 *JAMES M. SAFFORD, M. D., LL. D. Died July 3, 1907.
 *CHARLES SCHAEFFER, M. D. Died November 23, 1903.
 *NATHANIEL S. SHALER, LL. D. Died April 10, 1906.
 CHARLES WACHSMUTH. Died February 7, 1896.
 THEODORE G. WHITE, Ph. D. Died July 7, 1901.
 *GEORGE H. WILLIAMS, Ph. D. Died July 12, 1894.
 *J. FRANCIS WILLIAMS, Ph. D. Died November 9, 1891.
 *ALEXANDER WINCHELL, LL. D. Died February 19, 1891.
 ALBERT A. WRIGHT, Ph. D. Died April 2, 1905.

Summary

Original Fellows.....	60
Electd Fellows.....	235
<hr/>	
Membership	295
Deceased Fellows.....	50