strongly ribbed, and is clay-yellow in colour, mottled with fulvous, and is darker towards the tips. The membrane is roseate brown, and has a yellow patch on the costa and another at the base. The legs are reddish brown above and olivaceous beneath, paler at the joints; the thighs are but slightly dilated. The abdomen is ovate, flattened, and extends beyond the elytra. It is of a roseate brown. On either side of it are six sutures marked with yellow. The anal segment ends in a pair of incurved lobes. The whole of the under side is lighter in colour than the upper.

Taken at Quebec.

A NEW ANAPHORID, AND A NOTE ON AN OLD ONE.

BY HARRISON G. DYAR, WASHINGTON, D. C.

Eulepiste Kearfotti, n. sp.

Gray, with a reddish ochreous tint, brighter in an obscure streak beyond cell and on submedian fold. A series of black strigæ along the costa and on fringe; a small dash beyond cell, and an oblique bar in submedian fold beyond middle. Hind wing blackish, fringe long, pale, interlined with blackish. Below, blackish, with a pale line at the base of the fringe. Expanse, 22 mm.

Two males from Mr. W. D. Kearfott's collection, "Yuma Co., Ariz. Desert."

Larger than the other species of *Eulepiste*, and differing in the genitalia. Uncus a single long spine, curving downward, opposed to a broad, concave basal plate. Side pieces strap-shaped or slightly concave, curved downward, and with a distinct spine on the lower angle.

U. S. National Museum, type No. 6734.

Pseudanaphora mora, Grote.

In 1895 Lord Walsingham examined Grote's type in the British Museum, and thought it might be the female of *P. arcanella*, Clem., overlooking the description of the true female of this species by Beutenmüller (Ent. Amer. IV., 29, 1888). I have now before me ten females and eight males of *mora* from localities in New York, Pennsylvania and the District of Columbia, a majority of them taken by Mr. F. A. Merrick, at New Brighton, Pa. (see Proc. Ent. Soc., Wash., V. 40, 1902). There is a marked sexual dimorphism, the male being nearly uniformly blackish, and the female of a light ochreous ground colour. The species is very distinct from *arcanella*.