ing to character and capacity is an elephant trained. If of tractable disposition and handsome appearance, he becomes the pride of a state procession, the joy of his faithful mahout; or he may prove of brave and stubborn spirit, and be valued as an indispensable assistant in the work of bringing others of his species from the destructive life of the forest to one of service to mankind. Possibly he turns out just an ordinary catch, a mere beast of burden, a hewer of wood, and drawer of water. Elephant-hunting is the sport of kings, but not necessarily an extravagance.

In their sagacity and self-sacrifice these animals are sometimes an example to their human conquerors. One very fine beast, whose capture we witnessed, was understood to have led his pursuers a long round in order to give his companion time to escape. He did not seek cover in the jungle, but at once made boldly for the open plain. Men skilled in the reading of woodland signs told us afterward that another animal was concealed in the long grass, from whose vicinity brave La Touche Guj drew the chase by the sacrifice of his own freedom. He received the compliment of being named after the popular Lieutenant-Governor, and well deserved the honor, as well as the appellation of Guj, which means king elephant. This noble animal is pictured enjoying a drink. All along his conduct was exemplary. The elephant that enshrined Buddha in one of his numerous incarnations could not have behaved better.

The news of the capture of a wild elephant is hailed with joy by the inhabitants of neighboring villages whose lives have been made hideous by the depredations of nomads. Even a tiny fence will serve to protect fields from a wandering herd—elephants are such suspicious creatures that so slight a defense would savor of a trap—but Indian agriculturists have no visible boundaries at all; the crops appear to melt into each other, and come to a seemingly capricious conclusion.

Messier, which lies between ν and σ Scorpii, he suddenly exclaimed to his sister—the famous Caroline Herschel—"Hier ist wahrhaftig ein Loch im Himmel" (here, truly, is a hole in the heavens). It was an ab-

THE LAST MOMENT OF FREEDOM.

solutely black vacuity, about four degrees in width, perfectly free from any stars, and especially remarkable owing to its proximity to one of the richest globular clusters in the heavens. Closely south of Herschel's dark hole just mentioned, Prof. Barnard has photo-

A PITIFUL FIGHT WITH FATE.

We returned to the comfortable camp after long days in the brilliant sunshine and invigorating air of an Indian winter, fatigued with the unusual exertion and occasional intense excitement, but heartily desiring a renewal of our novel experiences. All of us very grateful to our hospitable host the Maharajah of Balrampur, to whom we were indebted for the opportunity of participating in unique and most enjoyable sport.—Country Life (London).

HOLES IN THE HEAVENS.*

By J. E. GORE, F.R.A.S.

THERE are many dark spots in the Milky Way which seem to be openings or holes in that wonderful zone of stars. These dark spots, or "coal sacks," as they are also called, seem to have been first noticed by Pinzon in 1499. They were also described by Lacaille in 1763.

The most remarkable of these spots is the well-known coal sack near the Southern Cross. It is of roughly oval or pear-shaped form, about 8 deg. in length by 5 deg. in width, and forms a conspicuous object in the sky of the southern hemisphere. It is completely surrounded by the nebulous light of the Milky Way, which is here of considerable brilliancy. The bright stars α and β Crucis—the brightest stars of the Southern Cross—nearly touch its southeastern edge. It contains only one lucid star within its boundaries. With reference to its northern border Sir John Herschel says: "The transition from rich Milky Way to almost complete darkness is here very sudden." It is, however, by no means devoid of faint stars. On a photograph taken in 1891 by Mr. H. C. Russell, at the Sydney Observatory, numerous very small stars are visible, but there are several spots which seem to be completely void of stars, and absolutely black. One of these remarkable holes is near β Crucis, and another near α Crucis.

There are other remarkable coal sacks in the Milky Way. A long, narrow, dark spot runs from α Centauri for several degrees toward the northeast. There are several in Scorpio, one of larger size between β and ϵ Cygni, and one south of α Cygni.

Examined with a telescope, the Milky Way shows many examples of small coal sacks; and some may be seen with even a good binocular field glass. One night when Sir William Herschel was examining a part of the Milky Way closely east of the globular cluster 80

graphed a great nebulous region surrounding the stars ρ Ophiuchi and 22 Scorpii. This photograph shows several dark lanes in what seems to be at least a comparatively thin sheet of stars, and this distinguished astronomer thinks it is certain that these stars are at the same distance as the nebula, for they form part

of the sky. If this be so, and the evidence seems to point in this direction, it would follow that their distance from the earth is not so great as their faintness would lead us to imagine. In his "Cape Observations," Sir John Herschel gives a list of 49 spots in the southern hemisphere totally devoid of any perceptible star. But probably photography will reveal the presence of some faint stars in these dark spots.

Closely east of the star θ Ophiuchi is a dark chasm, which passes south and west of that star, and there are several other dark lanes and holes clearly visible on the photograph taken by Prof. Barnard at the Lick Observatory.

Another small black spot was observed by Barnard a little northwest of the star γ Sagittarii. This seems to have been previously seen by Trouvelot, who says: "C'est comme un sac à charbon en miniature, ou une ouverture de la Voie Lactée à travers laquelle la vue pénètre au delà de ce grand assemblage d'étoiles."

A little southeast of α Cephei, a photograph by Barnard shows a ring of nebulous light, with a comparatively dark interior; at least the stratum of stars filling the rings seems pierced by several holes.

The "key-hole" openings in the great nebula surrounding the variable star η Argus is a remarkable feature of that wonderful nebula. A little south of this hole there is a kidney-bean shaped opening, shown in Sir John Herschel's drawing in the "Cape Observations." This opening is visible on a photograph taken by Sir David Gill in March, 1892. The photograph confirms the accuracy of Herschel's drawing, and shows that the opening is in all probability a real hole through the surrounding nebulous matter.

In the region round the star 12 Monocerotis there is a remarkable nebula of irregular shape, somewhat resembling in its general character the great nebula in the sword of Orion. Dr. Roberts, describing a photograph he took of this nebula, says: "Some remarkable tortuous rifts meander through the nebula on the north preceding half of the nebula; their margins are sharp and well defined in the midst of dense nebula. They are as clearly cut as we see the canyons of great rivers, but their width may in reality be millions of miles, for we have no reason to assume that the nebula is nearer to the earth than the stars. It is, indeed, possible that the stars which dot the surface are nearer to us than the nebula."

About 3 deg. northeast of the star θ Canis Majoris is another nebula of irregular shape. Dr. Roberts says that the star D.M. 1848 is on the margin of a dark sinuous vacancy or rift in the nebula, through which we see into the starless vacancy of space beyond it. This opening closely resembles the "key-hole" opening in the great nebula in Argus. Dr. Roberts adds: "These vacancies are most conspicuously seen where the surrounding nebula is dense, though they are also visible in some parts where it is relatively faint. The margins of the vacancies are often sharply defined, and suggestive of the idea that in consequence of some internal strain, operating from opposite directions, the nebula was rent asunder, and the parts separated from each other."

In another nebula in Monoceros, photographed by Dr. Roberts, a little west of the triple star 15 Monocerotis, there is a remarkable vacancy or hole. Dr. Roberts calls it a dark tortuous rift, and says: "The rifts prove that the nebulae are not globular, but are like clouds with relatively small depths, and that we can see through them into the darkness of space

IN THE PATHLESS JUNGLE.

ELEPHANT-HUNTING IN UPPER INDIA.

of it. With reference to the Milky Way in general, he thinks that the stars comprising it are comparatively very small bodies, and that they consequently differ vastly in point of size, at least, from the ordinary stars

beyond." There are also very noticeable areas devoid of stars in the region surrounding the nebula.

On July 12, 1891, Prof. Max Wolf, of the Astrophysical Observatory of Heidelberg, discovered three

* Knowledge and Scientific News.

dark markings in the Milky Way, about 1½ deg. west of the star γ Aquilæ. He calls them the "triple caves," and they certainly present a very remarkable appearance on his photograph. Closely east of the same star, a photograph taken by Barnard shows some curiously shaped dark markings, which seem to be openings through the Milky Way in this region. On a photograph by Max Wolf, of the region near ε Cygni, there is a remarkable black hole and smaller ones.

The question naturally suggests itself, what is the real nature of these curious black spots? Some astronomers have suggested that they are due to masses of cooled down, or partially cooled down, nebulous matter which absorbs the light of stars behind them. The term "hole," which I have used in the present paper, implies that my own view is that they are really holes or openings through the regions of stars or nebulous matter, and in this view of the matter I am supported by the opinion of several astronomers, as some of the extracts quoted above will show. Photographs of the great coal sack near the Southern Cross prove conclusively, I think, that the darkness of this remarkable spot is due to a real paucity of stars compared with the richness of the surrounding regions, and probably the same thing is true of the other dark spots in the Milky Way. We have really no evidence of the existence of dark bodies in space. Prof. Newcomb thinks that there is probably little or no extinction due to dark bodies, and he says, "We may say with certainty that dark stars are not so numerous as to cut off any important part of the light from the stars of the Milky Way, because, if they did, the latter would not be so clearly seen as it is. Since we have reason to believe that the Milky Way comprises the more distant stars of our system, we may feel fairly confident that not much light can be cut off by dark bodies from the most distant regions to which our telescopes can penetrate. Up to this distance, we see the stars just as they are."* The companions of some of the Algol variables are usually spoken of as dark bodies, but I have shown elsewhere that we have no reason to think that they are really dark. The companion of Algol, for example, may be a star of the 5th magnitude—a comparatively bright star—and yet be quite invisible to us, as neither the telescope nor spectroscope would show any trace of its existence. The little evidence we have tends to show that the satellite of Algol is *not* a dark body. The idea of "dark bodies" and "dark stars" seems to have been based on the existence of this eclipsing satellite; but it has been recently found that a difference of brightness of two magnitudes between the components of a spectroscopic binary star—like Algol—would be quite sufficient to obliterate the spectrum of the fainter star, the spectroscope merely showing the spectrum of the brighter component. Dark bodies *may* exist in space, and probably do, but as yet we have no positive evidence of their existence. The "holes in the heavens" are, I think, real, and "dark companions" of Algol variables have probably no existence except in the imagination of some astronomical writers.

It has been stated by several writers that the existence of these holes indicates that the Milky Way has a small extension in the line of sight; or, in other words, that it forms a comparatively thin stratum of stars. But Prof. Seeliger has shown that, according to the law of probabilities, if the number of stars be the same in both cases, the probability against the occurrence of these holes is just *the same* whether the extension of the Milky Way in the line of sight be great or small.† We cannot, therefore, come to any conclusion as to the actual thickness of the Milky Way from the appearance of the dark spots. It may have a great extension in the line of sight, or it may be comparatively thin in that direction. The cause of these holes must probably be looked for in the influence of some clustering power, as Sir William Herschel termed it, which tends to draw the stars away from certain spots and accumulate them in others. The existence of globular and other clusters close to dark and comparatively starless spots seems very suggestive in this connection. If these dark spots were due to intervening dark bodies, there seems to be no reason why we should find them so often close to rich regions.

[Continued from SUPPLEMENT No. 1584, page 25379.]

VALUABLE ALLOYS.—V.

BISMUTH ALLOYS.

Cliché Metal.

This alloy is composed of tin 48 parts, lead 32.5, bismuth 9, and antimony 10.5. It is especially well adapted to dabbing rollers for printing cotton goods, and as it possesses a considerable degree of hardness, it wears well.

For filling out defective places in metallic castings, an alloy of bismuth 1 part, antimony 3, lead 8, can be advantageously used.

Alloys of Bismuth, Tin, and Lead.

The readily fusible alloys of bismuth, tin, and lead have a somewhat higher melting point than those containing cadmium; they were earlier known than these, and are used for various purposes. Newton's metal and Rose's alloys are included here.

Newton's metal consists of bismuth 8 parts, lead 5, tin 3. It melts at 202 deg. F.

Rose's alloys consist of:

	I.	II.
Bismuth	2	8
Tin	1	3
Lead	1	8

* Harper's Magazine, October, 1904.
† Astrophysical Journal, vol. 12, p. 377.

The first of these alloys melts at 200.75 deg. F., and the other at 174.2 deg. F. They were formerly used in the manufacture of the so-called safety plates inserted in the top of steam boilers. These plates were intentionally made of a readily fusible alloy, so that at a certain temperature, corresponding to a certain pressure in the interior of the boiler, they would become fluid, and allow the steam to escape through the opening thus made. They were to act as a sort of safety valve, to prevent the explosion of the boiler with too high a pressure of steam. But however correct the principle may appear, it was found in practice that the boilers would frequently explode without the plates having melted; and they are at the present time hardly used at all. Chemical and physical tests have shown that by long-continued heating of the plates new alloys are formed, whose melting points are much higher than those of the original compositions. The following table gives the compositions of some alloys which are said to melt, if the pressure of the steam exceeds that indicated:

Bismuth.	Lead.	Tin.	Melting Point, Degrees F.	Corresponding Pressure of Steam in Atmospheres.
8	5	3	212.0	1
8	8	4	235.9	1½
8	8	8	253.9	2
8	10	8	260.0	2½
8	12	8	270.3	3
8	16	14	284.5	3½
8	22	12	300.6	4
8	32	24	308.8	5
8	32	26	320.3	6
8	32	28	331.7	7
8	30	24	341.6	8

Bismuth Alloys for Delicate Castings.

For delicate castings, and for taking impressions from dies, medals, etc., various bismuth alloys are in use, whose composition corresponds to the following figures:

	I.	II.	III.	IV.
Bismuth	6	5	2	8
Tin	3	2	1	3
Lead	13	3	1	5

These alloys have the property, very favorable in making sharply outlined castings, that they expand strongly on cooling, and so fill out the finest elevation and depressions of the mold.

Bismuth Alloy for Cementing Glass.

Most of the cements in ordinary use are dissolved, or at least softened, by petroleum. An alloy of lead 3 parts, tin 2, bismuth 2.5, melting at 212 deg. F., is not affected by petroleum, and is therefore useful for cementing lamps made of metal and glass combined.

SILVER ALLOYS.

Silver and Aluminium Alloy.

Aluminium and silver form beautiful white alloys, which are considerably harder than pure aluminium, and take a very high polish. They have the advantage over copper alloys of being unchanged by exposure to the air, and retaining their white color. It has been proposed to alloy coins with aluminium instead of copper, as this would make them much more durable, and they would keep their bright color; but the results of experiments made on a large scale did not warrant the change.

The properties of aluminium and silver alloys vary considerably according to the percentage of aluminium. An alloy of 100 parts of aluminium and 5 parts of silver is very similar to pure aluminium, but is harder and takes a fine polish. One hundred and sixty-nine parts of aluminium and 5 of silver makes an elastic alloy, recommended for watch springs and dessert knives. An alloy of equal parts of silver and aluminium is as hard as bronze.

Tiers-Argent.

This alloy is chiefly prepared in Paris, and used for the manufacture of various utensils. As indicated by its name (one-third silver) it consists of 33.33 parts of silver and 66.66 parts of aluminium. Its advantages over silver consist in its lower price and greater hardness; it can also be stamped and engraved more easily than the alloys of copper and silver.

Alloys of Silver and Zinc.

Silver and zinc have great affinity for each other, and alloys of these two metals are therefore easily made. The required quantity of zinc, wrapped in paper, is thrown into the melted and strongly-heated silver, the mass is thoroughly stirred with an iron rod, and at once poured out into molds. Alloys of silver and zinc can be obtained which are both ductile and flexible. An alloy consisting of 2 parts of zinc and 1 of silver closely resembles silver in color, and is quite ductile. With a larger proportion of zinc the alloys become brittle. In preparing the alloy, a somewhat larger quantity of zinc must be taken than the finished alloy is intended to contain, as a small amount always volatilizes.

Alloys of Silver, Copper, Nickel, and Zinc.

These alloys, from the metals contained in them, may be characterized as argentan or German silver with a percentage of silver. They have been used for making small coins, as in the older coins of Switzerland. Being quite hard, they have the advantage of wearing well, but soon lose their beautiful white color and take on a disagreeable shade of yellow, like poor brass. The silver contained in them can only be regained by a laborious process, which is a great drawback to their use in coinage.

The composition of the Swiss fractional coins is as follows:

	20 centimes.	10 centimes.	5 centimes.
Silver	15	10	5
Copper	50	55	60
Nickel	25	25	25
Zinc	10	10	10

Mousset's Silver Alloy.

Copper 59.06, silver 27.56, zinc 9.57, nickel 3.42. This alloy is yellowish with a reddish tinge, but white on the fractured surface. It ranks next after Argent-Ruolz, which also contains sometimes certain quantities of zinc, and in this case may be classed together with the alloy just described. The following alloys can be rolled into sheet or drawn into wire:

	I.	II.	III.
Silver	33.3	34	40.0
Copper	41.8	42	44.6
Nickel	8.6	8	4.6
Zinc	16.3	16	10.8

Silver Alloys Containing Arsenic.

Alloys which contain small quantities of arsenic are very ductile, have a beautiful white color, and were formerly used in England in the manufacture of table ware. They are not, however, suitable for this purpose, on account of the poisonous character of the arsenic. They are composed usually of 49 parts of silver, 49 of copper, and 2 of arsenic.

Alloys of Copper, Silver, and Cadmium.

Cadmium added to silver alloys gives great flexibility and ductility, without affecting the white color; these properties are valuable in the manufacture of silver-plated ware and wire. The proportions of the metals vary in these alloys. Some of the most important varieties are given below:

	I.	II.	III.	IV.	V.	VI.	VII.
Silver.....	980	950	900	860	666	647	500
Copper.....	15	15	18	20	25	50	50
Cadmium.....	5	35	82	180	309	284	450

In preparing these alloys, the great volatility of cadmium must be taken into account. It is customary to prepare first the alloy of silver and copper, and add the cadmium, which, as in the case of the alloys of silver and zinc, must be wrapped in paper. After putting it in, the mass is quickly stirred, and the alloy poured immediately into the molds. This is the surest way to prevent the volatilization of the cadmium.

Gray Silver (Japanese Silver).

An alloy is prepared in Japan which consists of equal parts of copper and silver, and which is given a beautiful gray color by boiling in a solution of alum, to which copper sulphate and verdigris are added. The so-called "mokum," also a Japanese alloy, is prepared by placing thin plates of gold, silver, copper, and the alloy just described over each other and stretching them under the hammer. The cross sections of the thin plates obtained in this way show the colors of the different metals, which gives them a peculiar striped appearance. Mokum is principally used for decorations upon gold and silver articles.

Imitation Silver Alloys.

There are a number of alloys, composed of different metals, which resemble silver, and may be briefly mentioned here.

Warne's metal is composed of tin 10 parts, nickel 7, bismuth 7, and cobalt 3. It is white, fine grained, but quite difficult to fuse.

Tonca's metal contains copper 5 parts, nickel 4, tin 1, lead 1, iron 1, zinc 1, antimony 1. It is hard, difficult to fuse, not very ductile, and cannot be recommended.

Trabuk metal contains tin 87.5, nickel 5.5, antimony 5, bismuth 5. This is similar to Warne's metal.

Tourun-Leonard's metal is composed of 500 parts of tin and 64 of bell metal.

Minargent.

This alloy, which is of a beautiful white color, contains no silver, but is made of copper, tungsten, aluminium, and nickel in the proportions of 1,000 parts of copper, 700 of nickel, 50 of tungsten, and 10 of aluminium.

GOLD ALLOYS.

Alloys of Gold and Palladium.

Alloys of gold, copper, silver, and palladium have a brownish red color and are as hard as iron. They are sometimes (although rarely) used for the bearings for the axles of the wheels of fine watches, as they cause less friction than the jewels commonly used for the purpose, and do not rust in the air. The composition used in the Swiss and English watch factories consists usually of gold 18 parts, copper 13, silver 11, and palladium 6.

Aluminium and Gold Alloy.

This alloy, called Nuremberg gold, is used for making cheap gold ware, and is excellent for this purpose, as its color is exactly that of pure gold, and does not change in the air. Articles made of Nuremberg gold need no gilding, and retain their color under the hardest usage; even the fracture of this alloy shows the pure gold color. The composition is usually 90 parts of copper, 2.5 of gold, and 7.5 of aluminium.

Colored Gold.

The alloys of gold with copper have a reddish tinge, those of gold with silver are whiter, and an alloy of gold, silver, and copper together is distinguished by a greenish tone. Manufacturers of gold ware make use of these different colors, one piece being frequently composed of several pieces of varying color. Below are given some of these alloys, with their colors: