virtues, but in a large clinic, where fifteen or twenty cases are to be operated on in an afternoon, it was to be feared that this method would be too slow. The Sluder method has little trauma and the bleeding is slight, and this method seems to offer very little advantage over Sluder's.

As to the claim that there would be less trauma than with a dull snare, Dr. Smith said he could not appreciate the difference, that any dull dissection would cause more trauma than a sharp dissection. The finger enucleation gives less bleeding but heals less readily and there is more after-pain and more sloughing, and as one departs from this extreme of dull dissection, just so much is the trauma lessened and the hemorrhage increased.

Dr. BRAUN, in closing the discussion, said that the advantages could be fully appreciated only by using the instrument. Anyone familiar with the Sluder technique could readily use this one properly. There is no difficult technique to be learned, as one can employ the instrument easily with the first case. Furthermore, all the operations in the throat should be done by the sense of vision rather than that of touch. With the Sluder technique, the finger is used in the engagement of the tonsil in the great majority of instances. With the technique described, it is not necessary to use the finger; everything can be determined by sight, and one can accurately see the line of cleavage. The lessened amount of hemorrhage is due to the fact that no sharp instruments are used, so that the blood vessels are crushed and not cut. The tonsil is pulled out of its bed and snared just in the line of cleavage, and the capsule is split. All that is crushed through is the mucous membrane and a very thin layer of connective tissue which serves to attach the tonsil to the intra-pharyngeal aponeurosis.

Dr. Braun said he made no point as to the length of time required for the operation; that if one could do a case in five minutes, not including the anesthesia, it could not be called a prolonged operation. No case requires over five minutes, including removal of adenoids. The time of operation will decrease with increasing familiarity with the technique, so that the operation may be performed in three minutes or less.

Spasmodic Asthma. R. H. BABCOCK, *Jour. A. M. A.*, Feb. 10, 1917.

Babcock believes that the disease is a sensitization due to protein absorption from some focus and in treatment all sources of such absorption must be eliminated if results are to be expected. The commonest site of foci is in the naso-pharyngeal region or the nasal accessory sinuses, though the tonsils may be the chief site.

Ed.