PULMONARY CONSUMPTION.

WORCESTERSHIRE STATISTICS.

By G. H. FOSBROKE, D.P.H., Camb.,
County Medical Officer.

IN February last the Worcestershire County Council convened a conference of persons interested in the new methods of treating consumption, to discuss the desirability of establishing a hospital in the county where the poorer classes could receive the "open-air treatment." The Conference was attended by representatives of nearly all the Corporations, District Councils, and Boards of Guardians in Worcestershire, as well as by many influential persons of both sexes.

Sir James Crichton-Browne delivered a most eloquent address on the causation and prevention of the disease, which was so much appreciated that the County Council has since caused it to be published *in extenso* and largely circulated. Dr. Paget Tomlinson gave a description of the hospital he has established with such advantage in Westmorland, and as a result of the discussion which ensued, the following resolutions were unanimously adopted, viz :

(1) That steps be taken to ascertain whether it is desirable to establish in this county a hospital for the open-air treatment of consumption among the poorer classes.

(2) That a communication be addressed to the different Poor Law and sanitary authorities in the administrative county of Worcester, asking, if such a hospital is established, what number of patients in their opinion would annually be sent from their districts.

(3) That a representative committee, consisting of representatives of the County Council, and of each of the Urban and Rural District Councils and Boards of Guardians in the administrative county, be appointed to consider the further working of the scheme.

In order that those who attended the Conference might not be wearied with figures, I was instructed, at short notice, to prepare some statistics bearing upon the question, and, accordingly, I compiled those I now present for consideration. Dr. Tatham most courteously supplied me with the death-rates for the decennial periods. The other rates were obtained from the annual reports of medical officers of health sent to the County Council.

It will be noticed that Table II. confirms the well-known fact that the persistent decline of pulmonary phthisis is a prominent

TABLE I. (a)

Showing rise and fall of Consumption (Phthisis) mortality in the Administrative County of Worcester and in England and Wales during the years 1890-99 inclusive.

(a) Compiled from Annual Reports of District Medical Officers of Health and Returns of Registrar General.

TABLE II. (b)
 Showing the rise and fall of Consumption (Phthisis) in the Registration
 County of Worcester and in England and Wales during the Decennial
 Periods 1861-70, 1871-80 and 1881-90.

Registration County of Worcester - Black line —————
 England and Wales - - - - - Dotted line - - - - -
 (b) Compiled from the Registrar General's Returns. Statistics for 1890-1900 are not yet available.

TABLE III.

Shows the relative mortality among *Males* and *Females* in England and Wales and Registration County of Worcester during the Decennial Periods 1861-70, 1871-80 and 1881-90.

feature in the vital statistics of England and Wales. Worcestershire evidently has an exceptionally low death-rate from consumption. In my annual report for 1898 I pointed out that Worcestershire, for an extended period, stands with Staffordshire and Leicestershire as one of the counties of England with least liability to the disease.

Table II. also shows that the Worcestershire death-rate has declined *pari passu* with that of England and Wales during each of the last three decennial periods. It is remarkable that the mortality of England and Wales during the ten years 1881-1890 was the same as it was in Worcestershire thirty years ago. Thus the death-rate of England and Wales during the decennial period 1881-1890 was 1,724 per 1,000,000, whereas in Worcestershire so long ago as 1861-1867 it was 1,704, and has since fallen to 1,196 in 1881-1890.

Table III. compares the relative mortalities among males and females in England with those of Worcestershire during the last three decennial periods. Here, again, the sex incidence is in favour of Worcestershire; but there seems to have been a greater decline in the county mortality among females than among males during 1861-1890.

In my annual report, previously alluded to, I stated that Worcestershire has, as compared with other counties, the lowest death-rate from phthisis among females; and in this respect it is below the neighbouring counties of Oxford and Warwick, and considerably below Shropshire. A coloured diagram, not published here, shows that the lowest death-rate recorded in the county during 1890-1899 was that of the Halesowen Rural District (0·4 per 1,000), and the highest that of the Redditch Urban District (1·5). Halesowen Rural and Stourbridge Urban Districts form the Worcestershire fringe of the Black Country, and it is noteworthy that in neither of these districts did the phthisis death-rate exceed 0·7 per 1,000 of the population during 1890-1899.

These low mortalities are interesting, and somewhat surprising when it is realized how indifferently the artizan population of these localities is housed. These districts, however, have good natural drainage, and it is not unlikely that their topographical position may account for the local immunity from phthisis.

Sir Hugh Beevor, Bart, in his Oration delivered before the Hunterian Society in 1899, speaking of the neighbouring County Borough of Dudley, said: "Dudley enjoys a comparative immunity from phthisis, which one is at first sight inclined to attribute to the smoke of the Black Country, from the heart of which it rises,

built on the top of a long slope of hill with a good fall on nearly every side."

The comparatively high mortality from phthisis in Redditch is probably due to the fact that the population is mainly engaged in the manufacture of needles. From inquiries I made in a neighbouring needle district some time back, I am inclined to think that the incidence of the disease upon needle-makers is not associated with irritation caused by inhalation of steel filings, but rather due to the fact that the artizans so often work in overheated rooms, where the liability to infection is great, and the ventilation far from what, consistent with health, it ought to be. This insufficient ventilation is not so much the fault of the manufacturers as of the workpeople themselves. In spite of the efforts of the former, the latter persist too often in closing windows and ventilators. Such being the case, it is no wonder that phthisis is easily communicated.

As my statistics were prepared for a popular gathering, no attempt was made to show the age-incidence of the disease.

Realizing in 1898 that Worcestershire was specially favoured with respect to phthisis, I communicated with Sir Hugh Beevor, who had lately been studying the question, and he replied that he was unable to suggest any definite point for consideration, as it seemed to him, to carry our knowledge further, intricate and prolonged investigation was necessary. Under such circumstances I have no explanation to offer why Worcestershire should be less prone to phthisis than other counties.

VALUATION OF JALAP TUBERS. O. SCHWESINGER (*Pharm. C. H.*, 1901, xlii. 1, through *Chem. Zeit. Rep.*, 1901, 14).—Ten grammes of the finely-powdered roots are extracted with spirits of wine for twenty-four hours at a temperature of 30° C., shaking at intervals. Fifty c.c. of the liquid are drawn off, and the solvent is evaporated. The resin is then washed with warm water as long as colouring-matter is removed, and the residue is dried. The results are concordant.—*Analyst*.

A TEST FOR PICROTOXIN. S. S. MINOVICI (*Ann. Pharm.*, 1901, vii. 1, through *Chem. Zeit. Rep.*, 1901, 52).—Anisaldehyde is a very delicate and characteristic reagent for picrotoxin, indicating its presence not only when alone, but also in the starchy mass of the fruit. A drop of sulphuric acid is added to the sample, and after two minutes, when the saffron-yellow has become distinct, one drop of a 20 per cent. solution of anisaldehyde in absolute alcohol is added. A dark-blue violet ring is produced, which soon changes to a permanent blue. The test can also be applied to dilute solutions in chloroform, alcohol, or water, but the liquids should be warmed. The limit of sensitiveness is 1 : 2,000, or even 1 : 5,000. Many other glucosides and alkaloids give obscure and unimportant colour reactions with the reagent; while convolvulin yields a cherry-red, saponin a red-brown, aconitine a pale rose, and veratrine a blue-red or indigo-blue tint.—*Analyst*.