Schreiben des Herrn E. J. White an den Herausgeber.

The following is the list of occultations of stars by the moon observed here during the period that the various parties for observing the transit of Venus were stationed in this hemisphere.

Melbourne.									
Date.	Name of Star.	Mag.	Phase.	Sid Time.	Mean Time.	Observ.	Remarks.		
October	1 B. A. C. 1882	5½	Re	2 ^h 42 ^m 37 ^s .3	14 ^h 1 ^m 42 ^s .5	M	Fair observation.		
	16 Lal. 32974	7½	Dis	0 18 15.5	10 38 45.7	M	" "		
	17 B. A. C. 6422	7	Dis	20 58 38.3	7 15 45.3	T	Good "		
	17 Lal. 35235	8½	Dis	23 23 4.2	9 39 47.5	T	" "		
	27 B. A. C. 1341	5½	Re	1 55 51.4	11 32 50.6	T	" "		
Novbr.	17 Lal. 43567	9	Dis	0 54 18.4	9 8 53.6	M	Not satisfactory, hazy. star faint.		
	17 Lal. 43596	8½	Dis	1 42 49.9	9 57 17.1	M	Very good.		
	18 B. A. C. 8102	6	Dis	4 8 44.4	12 18 51.8	M	Fair observation in hazy sky.		
	18 Lal. 45562	7	Dis	4 36 8.3	12 46 11.2	M	Very good.		
Decbr.	19 Lal. 4528	8	Dis	6 0 40.4	12 8 36.2	T	Good.		
	21 B. A. C. 1341	5½	Dis	3 49 56.2	9 50 21.6	T	"		
	23 Lal. 12661	6	Dis	4 50 59.8	10 43 23.3	M	Fair. thought star barely visible when near bright limb.		
	27 B. A. C. 3606	6	Re	8 41 55.0	14 17 57.1	T			
Januar	13 Lal. 512	6	Dis	5 21 23.9	9 51 8.3	T	Good.		
	18 Lal. 9484	7	Dis	5 48 38.4	9 58 38.8	T	"		

All the occultations have taken place at the dark limb with the exception of that of Dec. 23. The observations were made by Mes^{rs}. Moerlin and Turner, whose initials are given in the observers column. After many attempts I succeeded this morning in getting an observation of Encke's comet, it was the first clear morning that we have had for some weeks. In the equatorial of 4½ inches aperture the comet was an excessively faint nebulous object, without nucleus, compact shape, but slightly elongated. I am afraid that the presence of the moon will prevent on getting any other observation. As there has been no time for making the reductions I send you the observations as taken. A ring micrometer was used, whose internal diameter was equal to 14' 11". The chronometer was a sidereal one, whose error at 20^h 0^m sid. of the 17th was 0^m 18^s.04 fast, and gaining 2^s.40 per day. The comparison star was of the 8th magnitude, whose approximate mean place for 1875.0 was RA. 0^h 31^m 51^s,

N. P. D. 106° 55' 53". Reading of Barometer 30^m29. Fahr. Therm. 39^o5.

The comet was in the southern half of the ring, the star in the northern, but so near the centre as to give but a comparatively rough result in difference of N. P. D. Chronometer times.

Comet 20	41 49.5	* 20	44 40.8	
	42 45.0		45 38.2	
	47 19.0		50 10.0	
	48 14.0		51 7.8	
21	1 31.8	21	4 22.8	
	2 22.0		5 21.0	
	23 1.8		25 51.9	
	52.0		26 50.8	

The comet was of the last degree of faintness during this observation, owing to a slight haze, and the presence of twilight.

Melbourne Observatory. May 18, 1875.

E. J. White,

Acting Government Astronomer.

Inhalt:

Zu Nr. 2052. *Baeyer*. Ueber Fehlerbestimmung und Ausgleichung eines geometrischen Nivellements. 177. — *W. Doberck*. Elements of 36 Andromedae Σ 73. 187. — *L. Schulhof*. Elemente und Ephemeride von (147) Protogeneia. 189. — *E. J. White*. Schreiben an den Herausgeber. 191.

Kiel. 1875, August 30.

Druck von Fiencke & Schachel in Kiel.