method with my auto-inflating bag.³ (Fig. 5.) Frequent practice of Valsalva is apt to produce congestion of the tympanic and labyrinthic vessels, and should be avoided. I do not touch on the question of removal of the tympanic

FIG. 5.

membrane and ossicles, as proposed by Schwartze and performed by Sexton of New York and others; nor do I refer to mastoid complications: this discussion was not contemplated in the preparation of the present communication.

TWO CASES OF CHOLECYSTOTOMY.

By HERBERT W. PAGE, M.A., M.C. CANTAB.,
SURGEON TO ST. MARY'S HOSPITAL.

CASE 1.—Alice J—, aged thirty-seven, a married woman, with good family and personal history, was admitted into St. Mary's Hospital under Dr. Cheadle's care on May 10th, 1888. She had been jaundiced for the previous four months, and gave an account of having had repeated attacks during that period of severe pain in the hepatic region and between the shoulders. A fortnight before coming to the hospital, after an attack of pain so bad that she had had to have injections of morphia, the medical man then attending her found gall-stones in the fæces. The motions had all along been very pale. On admission she seemed in a very exhausted condition, and was intensely jaundiced. The motions were white, and the urine contained large quantities of bile. There was much general tenderness or fear of being touched over the whole abdomen, but the tenderness over the liver and region of the gall-bladder was real and acute. The abdomen was rigid, and respiration was distinctly thoracic. No tumour of any kind was to be felt. With the exception of undoubted increase in the severity of all her symptoms, there was no new feature in her condition during the next few weeks. The thing most noticeable was her surely increasing prostration, with an occasional rise of temperature to 100° or 101°, and it was felt that some operation was imperatively called for to save her life. The diagnosis of complete obstruction in the common duct seemed certain, and she was therefore transferred to my wards on June 13th, with a view to operation. Having opened the abdomen by a three-inch incision to the right of the right rectus muscle, immediately over the gall-bladder, this viscus was found lying very deeply in the abdomen, and contracted to a size not larger than that of a walnut. The gall-bladder was then incised, and exit given to a teaspoonful of clear glycerine-like fluid. Two very small gall-stones were taken out with a scoop. Now began the chief difficulties of the operation. Lying still deeper than the gall-bladder, the common duct, or a cyst representing it, could just be felt with the tip of the finger, and in it a number of small stones which could be moved freely about. By no force, however,

³ In this auto-inflating bag there are two ball-valves connecting the tubes A and B with the bag. The tube B is for the purpose of filling the bag with the patient's respired air or with any vapour he chooses to aspirate with. It is not necessary in general use to employ this. Then the nasal tube alone is used. The slightest touch on the soft rubber bag, if properly applied during the swallowing of a little saliva, will inflate the tympanum.

that I was able to apply to them, either from the outside of the duct or from within, by finger, scoop, probe, or catheter, could they be moved away from their position. The depth of all the parts from the surface rendered it quite impossible to open the duct, at any rate without much more formidable operative measures than had been contemplated, and I therefore determined very freely to manipulate these stones in the hope that in the future, if not at that moment, they might deem it well to move on into the bowel; and there seemed encouragement to this course in the fact that two small stones had been found in the fæces before the woman came to St. Mary's. No impression, however, was made upon them. The gall-bladder was sutured to the abdominal wall in the usual way, and the operation was finished with a sense of disappointment at its want of success. The patient bore the operation well and quickly rallied from it. Already on June 18th she felt a good deal better, and when the wound was dressed for the first time on the 19th we were pleased to see that the discharge was very distinctly bile-stained. Now, at any rate, there was hope that, even if a passage could not be made along the ductus communis, the bile would be prevented from passing into the blood and so imperilling her life by steadily increasing cholæmia. She would certainly be in a better position to await events. The biliary discharge increased very much in quantity, so that on June 28th the dressings had to be changed twice daily. In herself she was very much better; the temperature had never been above the normal range; her jaundice was fading away; she was free from pain; and the fæces, which had been examined as if in search for gold, occasionally were noted to be distinctly less white; no gall-stones were ever discovered in them. At the end of the third week of July the discharge much lessened and the wound showed every sign of rapid closure. Once only, on July 31st, it was noted that she was again rather yellow, and this although the biliary discharge had not ceased and the fæces unquestionably contained bile. She left the hospital on Aug 4th. She returned early in October in great alarm at what she had thought to be a return of her old pain. We came, however, to the conclusion that she was really suffering from an attack of ordinary colic—a conclusion supported by its early disappearance, and by the absence of any return of her other symptoms. True, there was much hyperæsthetic or nervous tenderness about the site of the wound, now soundly healed, but there was no rise of temperature, and the fæces were much more obviously bile-stained than when she left the hospital in August. She was accordingly discharged in a few days. I saw her next on Nov. 10th, when she looked and said she was perfectly well, although on one day in the previous week her conjunctivæ had been yellow and her urine was high-coloured. There was then no trace of jaundice, nor has there been on the several occasions when I have seen her since. I sent for her and examined the abdomen on March 9th, 1889, and found it natural in every respect. The motions were then of a much better colour, but she was sure that she was occasionally rather yellow. Of this case, therefore, I think it may be said that the operation was of signal use: first of all, in rescuing the woman from the dangers of cholæmia; and, secondly, in bringing about a return of patency in the common duct, whereby the natural channel has been restored, partially if not entirely, for the passage of bile into the alimentary canal—results which at one time seemed almost hopeless.

CASE 2.—Here, also, unforeseen difficulties, although of quite a different kind, presented themselves at the operation. A lady aged sixty-two, a patient of Dr. Morton of Kilburn, who had been seen in consultation by Dr. Broadbent, had in most respects enjoyed good health, but for years had been subject to attacks of pain in the region of the gall-bladder. Once only had she had a slight attack of jaundice, the motions had never been pale, and no tumour had ever been discovered until three or four months before I first saw her with Dr. Morton in November, 1888. Since its first discovery this tumour had invariably occupied the region of the gall-bladder, and had varied in size, being decidedly larger whenever there were attacks of pain. The swelling was the size of a large orange, reached as low as the umbilicus, and extended across the middle line. The abdominal wall was, like the rest of her body, very thin; and the tumour, which could therefore be easily defined, seemed to be freely movable. There was no great amount of tenderness. The history and symptoms thus alike pointing to the lodgment of gall-stones in the cystic duct and bladder,

and the necessity for giving her relief being no less obvious, for she was being worn out with increasing pain, the operation of cholecystotomy was undertaken on Nov. 29th, 1888. A three-inch vertical incision was made over the centre of the tumour, a little nearer the middle line than the linea semilunaris. And now arose the difficulties in the operation, for the parts had lost all precise anatomical differentiation from being firmly matted and adherent together, and it was practically impossible at any moment to say where the scalpel was cutting. Presently a small cavity was opened which gave exit to two or three drops of pus-like matter. This only served to confirm the existence of inflammatory adhesions, which, from the free mobility of the tumour, had not been previously suspected. No track could be made out from this small abscess to the gall-bladder itself, and a fine aspirator needle was therefore passed into the centre of the swelling which now remained, and which, freed from the covering of the abdominal wall, already felt much smaller than before the operation was begun. There flowed at once through the cannula a small quantity of clear fluid, and I therefore passed first of all a director, then a pair of forceps, and lastly my finger, into the cavity which had been so luckily tapped. Minute faceted gall-stones were immediately felt, and to the number of twelve were removed with the finger or scoop. Finally, one stone, the size of a large marble, and somewhat firmly fixed in a remote corner of the cavity, was got out. Almost immediately there was evidence of escape of bile, the very best sign that the obstruction had been removed. All this had clearly been accomplished without opening the peritoneum; but the operation had not been easier on that account; nay, for the reasons named, it had been decidedly more difficult. Surgeons would invariably prefer, one would think, to have in such a case to cut through tissues which have a known relation to each other, rather than into parts which have been deprived of all such relationship by inflammatory and destructive adhesions. There were no bad symptoms of any kind after the operation. There was the usual copious flow of bile, and the wound steadily healed. The drainage-tube was taken out on Dec. 11th, and on the 15th the patient was able to be moved on to the sofa. A week after this the wound was healed. I have not seen her myself since, but Dr. Morton reported on Feb. 24th that she had made an uninterrupted recovery. She then expressed herself as quite well and comfortable. She has since been out of doors.

The opportunity may here be taken, in conclusion, of saying that the patient whose case was recorded in THE LANCET (Mirror, June 25th, 1887) by my then house surgeon, Mr. Lloyd, has enjoyed good health, without any return of her former symptoms, up to the present time. That also was a case of gall-stones blocking the cystic duct, but presented no difficulties or special features. A fourth case, still in the hospital, and even more satisfactory in its result than these three, will be published shortly.

THE INDUCTION OF PREMATURE LABOUR BY MEANS OF HEGAR'S DILATORS OF SPECIALLY LARGE SIZE.

WITH NOTES OF THREE CASES.

By ARTHUR H. N. LEWERS, M.D., M.R.C.P. LOND.,
ASSISTANT OBSTETRIC PHYSICIAN TO THE LONDON HOSPITAL.

THE method most in favour at present for the induction of premature labour is the passing of a bougie several inches into the uterus, between the membranes and the uterine wall, and leaving it there till labour pains come on. Hot vaginal douches are usually given as an adjuvant from time to time while the bougie is in place. Now, it not rarely happens that there is a considerable interval between the time of insertion of the bougie and the setting-in of labour. This interval has been as much as a week in some cases that have come under my observation, both in my own practice and in that of others. It cannot be foretold with any certainty how long it will be before labour will come on after the insertion of the bougie. The practical inconveniences of this to everybody concerned are, of course, obvious. Another plan occasionally employed is to

dilate the cervix with tents. Here there is the risk of septicæmia, and this is especially so if more than one set of tents has to be used.

Looking at the satisfactory results of dilating the cervix with Hegar's dilators in gynaecological practice, it seemed to me that a similar method was worth trying for dilating the cervix in order to induce labour. The ordinary dilators used in gynaecology are not, however, of sufficient size for this purpose, and therefore Messrs. Kröhne and Sesemann have supplied me with a larger series running up to No. 40, which is $1\frac{3}{4}$ in. in diameter. It will be remembered that the ordinary series stops at No. 26. The details of the method are as follows. The patient is given a copious vaginal douche of hot iodine water; she then lies in the ordinary obstetric position, and the os uteri is brought into view with Sims' speculum. Hegar's dilators are then passed one after the other till a size is reached which seems to be the largest that will for the time being pass into the cervix. The dilators lie ready for use in a porcelain tray covered with 1 in 40 carbolic lotion, and as each one is wanted it is dipped in terebene oil (1 in 5). In a few minutes the dilator will be found to be fitting less tightly, and then it may be withdrawn and the next size introduced. As the larger sizes are reached, more time should, if necessary, be allowed between the introduction of the dilators; but it may happen, as in my first case, that the whole series as far as No. 34 (which was then the largest I had) may be passed one after the other without any difficulty being met with. During the process the vagina is from time to time irrigated with 1 in 40 carbolic lotion. When the largest bougie of the series has been passed, it is left in the cervix, the lower end of the bougie resting against the posterior vaginal wall. The presence of the dilator, *plus* the degree of dilatation already present, may be sufficient to start strong pains, and when this is so, the dilator is withdrawn and labour allowed to proceed in the usual way. If only slight pains, or no pains, result—say, four hours after the insertion of the largest size—the membranes may be ruptured. Considering that the os is dilated to the size of a circle $1\frac{3}{4}$ in. in diameter, this proceeding is not open to the objections that apply to rupturing the membranes in the first instance, when the os is undilated. In the following cases the method just described was adopted, and the results, both as regards mother and child in each case, were satisfactory.

CASE 1. *Induction on account of children at term being of unusually large size; failure of ordinary method; success of method above described.*—J. D.—, aged twenty-three, had been married three years and a half, and had had two children, the last fourteen months before admission into the London Hospital. She was admitted on June 11th, 1888, for the purpose of having labour induced. She said that both confinements had been very difficult, and that delivery had been accomplished by destroying both children with instruments. It was not clear, from her account, whether forceps only had been used, or whether craniotomy had been performed. Both children were very big; the last one was 23 in. long. The woman was below the average as to height, but there was little if any contraction of the pelvis, the measurements of which were as follows: between anterior superior iliac spines, $9\frac{1}{2}$ in.; maximum measurement between iliac crests, $10\frac{3}{4}$ in.; external conjugate, $7\frac{1}{2}$ in.; diagonal conjugate $4\frac{1}{2}$ in., or a trifle less. At the date of admission the woman was estimated to be thirty-five weeks pregnant. On June 12th, at 3.15 P.M., a vaginal douche of iodine water was given, and a celluloid bougie with a stylet was introduced into the uterus. The total length of the bougie was $12\frac{1}{2}$ in., and it was passed into the uterus so that the lower end was well within the vagina, and resting against the posterior vaginal wall. On the 14th hot douches were given every three hours after the bougie was inserted, but she had had no pains. The bougie was taken out and reinserted, the douches being continued as before. On the 18th, at 5.30 P.M., the patient had remained *in statu quo*. The bougie was therefore taken out, and Hegar's dilators were used to dilate the cervix in the manner already described. Nos. 16 to 26 of the ordinary series passed easily, and so did Nos. 28, 30, 32, and 34 of the special series. No tenaculum was used to steady the cervix, and all the dilators mentioned passed without any difficulty. At that time No. 34, which is $1\frac{1}{2}$ in. in diameter, was the largest I had. It was left in the cervix till 8.30 the same evening, when, as the patient was having labour pains regularly, it was removed by Dr. Wright, the resident accoucheur. The labour then