needs—so individualised and so hopelessly jealous among ourselves that we will not co-operate and organise our forces. Nothing could show this more clearly than the way in which the Royal Society has sacrificed its opportunities during the war period by failing to bring all its fellows together and thus make our corporate influence felt. Little wonder that science is being more and more excluded. We are talking of government by public opinion but real opinion cannot be made public. The columns of the Press are not open to any serious discussion from our side; the books we write are not understood even by ourselves. Owing to excessive specialisation we are becoming less and less able to express and protect ourselves, less interested in stating our case in any broad way; what is worse, too selfishly immersed each in his own narrow occupation to consider the general interest: and we be not careful the position won for us by the protagonists of the past will soon be lost to us.

Our failure is as window-dressers; we have not displayed our goods in the right manner. When a business is a failure, the only way to re-establish it is to reconstruct the management. That we have yet to learn this elementary truth is clear, however, from the recent appointment to the treasurership. A young man should have been found for the office, gifted not only with scientific experience but also with some breadth of outlook, some understanding of men and affairs; instead, a senior member of the Association has been chosen.

If a senior were desirable, no better choice could have been made; but I believe my old friend Principal Griffiths will not mind my saying that he is saved from inclusion in my class of elderly amiables only by the fact that throughout the war he was conspicuously active in his efforts to bring the value of scientific method before the public; he is far too soft-hearted to face the exigencies of the present situation.

The council must cease to be all but entirely an old or even an elderly man's show; responsibility must be forced upon the younger generation of active workers.

The secretaries obviously have no policy; new men must be chosen at the earliest possible date—I trust on the three years' hire system advocated in my former letter, so that the different subjects may be cared for at shorter intervals.

The real question at issue is: "How are we to get at the public?" To do this we must look beyond the meetings and in some way arrange for the preparation and issue each year of a few authoritative tracts dealing with bedrock problems in language that can be understood by all. Subjects can easily be found; under E alone enough are to hand—*Economics, Education, Energy, Entomology, Eugenics, Evolution*. Nobody believes in these, few know what the terms comprise, yet the future of our race depends on an intimate understanding and application of the doctrines they cover. A great work would be done by the Association if only, by real discussion not the dreamy, introspective twaddle of Section L, the nonsense now spoken of as education were reduced to terms of wisdom.

HENRY E. ARMSTRONG.

The Constitution of the Elements.

SINCE my last letter under the above title in NATURE of July 1, experiments have been made with a few more elements. The work has been progressively more and more difficult, for it has not been easy to find volatile compounds suitable for use, and when

found the very objectionable chemical properties of some of them have led to experimental trouble and disappointing results. Thus selenium, tin, antimony, and tellurium have so far yielded no result of any value.

Fortunately, iodine (atomic weight 126.92) gave definite and unmistakable effects. It proves to be a simple element of mass 127—a result satisfactorily confirmed by a single line at 142 corresponding to CH_3I , the vapour used in the experiments. This result has particular interest in view of the recent work of Kohlweiler (*Zeit. Phys. Chem.*, vol. xcv., 1920, p. 95), who deduces five isotopes of iodine, all of even integral atomic weights, and claims to have effected a considerable separation of these by diffusion.

Owing to the kindness of Prof. Collie and Dr. Masson in providing me with a sample of gas rich in xenon, I have been able to identify two more probable isotopes of that element and obtain trustworthy values for the atomic weights of the five already found. The provisional figures given for these turn out to be too low. The values quoted below were obtained from the position of the second-order line 64.5. They should be trustworthy to about one-fifth of a unit.

Additional evidence on argon leads to the conclusion that its isotope 36 need no longer be regarded as doubtful.

The following table gives the results to date:

Table of Elements and Isotopes.

Element	Atomic number	Atomic weight	Minimum number of isotopes	Masses of isotopes in order of their intensity
H	1	1.008	1	1.008
He	2	3.99	1	4
B	5	10.90	2	11, 10
C	6	12.00	1	12
N	7	14.01	1	14
O	8	16.00	1	16
F	9	19.00	1	19
Ne	10	20.20	2	20, 22, (21)
Si	14	28.30	2	28, 29, (30)
P	15	31.04	1	31
S	16	32.06	1	32
Cl	17	35.46	2	35, 37, (39)
A	18	39.88	2	40, 36
As	33	74.96	1	75
Br	35	79.92	2	79, 81
Kr	36	82.92	6	84, 86, 82, 83, 80, 78
I	53	126.92	1	127
X	54	130.32	5, (7)	129, 132, 131, 134, 136, (128, 130?)
Hg	80	200.60	(6)	(197-200), 202, 204

(Numbers in brackets are provisional only.)

F. W. ASTON.

Cavendish Laboratory, November 30.

Solar Variation and the Weather.

IN NATURE of July 29 last (p. 678) appears an article by Dr. C. G. Abbot on solar variation and the weather, in which reference is made to the use of solar data by the Argentine Weather Service. Drs. Nansen and Helland-Hansen have also found some interesting correlations between the variations of solar radiation as measured by Dr. Abbot and variations of temperature and pressure in Norway.

As other investigators will no doubt be tempted into this field, I feel that it may be of interest to give briefly a summary of our latest results and conclusions. These conclusions are based on the study of an immense amount of data from various parts of

the world, the results of which will be published in due time.

The first and most striking result is that the solar radiation exerts a cumulative effect on the atmosphere so that prolonged periods of high or low solar radiation have a much greater response in atmospheric action than shorter periods of greater intensity. Thus a deviation of 5 per cent. from the average radiation of short duration might not produce so great an effect as a long period, with a mean deviation of 2 per cent.

The second result is that as the sun changes from one hemisphere to another the effect on the weather changes, so that in the hemisphere where the sun is nearly vertical the pressure falls over the land surface and rises over the water surface, while the opposite effect is found in the other hemisphere. This effect I take to be a proof that a considerable part of the increased solar radiation reaches the earth's surface and intensifies the normal effect of absorption of solar radiation by the land. There are, however, clear indications that a considerable part of the increased solar radiation is absorbed by the upper air and gives origin to atmospheric waves which, if they are not the determining cause, at least are very influential in determining the discontinuities referred to recently in *NATURE* by Prof. Bjerknes.

These atmospheric waves are of a complex nature, and for this reason the correlations with solar changes are best seen by separating the solar and weather changes into shorter and longer (or slower) waves.

By taking averages of ten days we found for 1916 correlations as high as 84 per cent. with certain Argentine stations, and correlations exceeding 80 per cent. in later years. The shorter waves do not give such high correlations, apparently for several reasons, the most important, perhaps, being that there are still some errors in the solar values due to the difficulties of eliminating the effect of changing transparency of the air while the solar observations are being made.

This difficulty has undoubtedly been greatly diminished during recent months by the use of the pyranometer, which enables the observers to get measurements of the solar heat from a single bolograph. Certainly since this method was begun the correlations of the solar variations with atmospheric waves have increased. Another difficulty appears to be that while the waves start from definite centres of action, these centres shift position to some extent, so that the time interval of the effects following solar changes is variable, and this variability is of greater importance in the shorter waves.

The prolonged effects following unusually high or low values of radiation shown by Dr. Abbot's curves, reproduced in *NATURE*, I believe to result from the normal sequence of events on the surface of the sun. After a marked excess or defect of radiation there is likely to follow a similar deviation from the normal after an interval of ten to thirteen days.

This result I believe to arise from the fact that when there is a marked outbreak of heated solar gases on the edge of the sun, where absorption is normally great, the total radiation is more intensified than when the outbreak is near the centre. If the outbreak is near the edge it will be carried by rotation to the opposite edge in from about eleven to nineteen days, according as to whether the position is on the east or west edge.

In the same way cooled gases produce their greatest absorbing effects when near the edge of the sun, and there is the same tendency to repeat. These effects also tend to repeat themselves after a solar rotation,

but in that case the interval is so long that marked changes have usually occurred. However, these repetitions are sufficiently numerous to make it practically certain from the length of the rotation that the conditions causing the marked deviation from normal are in the region of the prominences, and not in that of the sun-spots, which have a period of about twenty-seven days.

These studies of the relations between solar changes and the weather have been in progress in the Argentine Meteorological Office for several years, so that when a station of the Solar Physics Observatory of the Smithsonian Institution was opened in Chile, arrangements were made between Prof. G. O. Wiggin, director of the Argentine Weather Service, and Dr. Walcott, director of the Smithsonian Institution, for the transmission of the solar radiation measurements by cable to Buenos Aires and for their use so far as possible in weather forecasting. To this work Prof. Wiggin and the assistant chief have given enthusiastic support, and through their efforts, and as a result of a growing interest in the subject in Argentina, the Legislature has recently appropriated some 25,000 pesos for the installation of a solar observatory in Argentina.

The forecasts are made for a week in advance by publishing estimated temperatures for each day, and also forecasts of expected rains. With increased experience there have been steady improvements and an increasing demand for the forecasts by commercial interests.

There are many complexities which remain to be solved, and perhaps some which can never be solved, but I believe that a distinct advance has been made in forecasting, and the progress is too far advanced to be turned backward. One of the greatest difficulties is the inversion of the solar relation with the season, and occasional inversions for other reasons not yet clearly understood.

Heretofore, most of the studies by others in regard to the relation of solar changes and atmospheric phenomena have been in reference to the sun-spots. Dr. Abbot's observations indicate clearly that there is a change of solar radiation corresponding to the eleven-year sun-spot period. The radiation increases with the increase in spottedness, but this change is of relatively small importance compared with shorter changes of greater intensity.

Sir Norman Lockyer suggested some years ago that there were weather changes due to solar changes other than the eleven-year period, and brought evidence to show that a period of about four years in the outbreak of prominences caused atmospheric changes having the same period. But solar radiation shows a variability far greater than is indicated by the observations of prominences, and the high value of the correlation between these and the weather which we have found for the Argentine and for other regions of the world leads me to believe that these changes are the chief, if not the only, cause of weather changes as distinct from the well-known diurnal and annual periods.

H. H. CLAYTON,
Chief of Forecast Division,
Oficina Meteorológica Argentina.

The Physical Meaning of Spherical Aberration.

THE nature of the distribution of light round the axial focus of a lens is a problem which has frequently been attacked mathematically. It has been quite successfully solved for points in the principal focal plane of an "aplanatic" lens when the intensity is found to