THE DUBLIN JOURNAL

OF

MEDICAL SCIENCE.

JANUARY 1, 1879.

PART I.

ORIGINAL COMMUNICATIONS.

ART. I.—*Case of Pachydermatocele.* By ARTHUR WYNNE FOOT, M.D.; Senior Physician to the Meath Hospital and County Dublin Infirmary.

THIS very remarkable case was an example of the pachydermatocele of Valentine Mott—or, as this term is, perhaps, confined rather to cutaneous overgrowths affecting the upper parts of the body, it might be more appropriately named simply elephantiasis, or Barba-does leg. By a glance at the drawings anyone can see that its right to a title founded on likeness to an elephant's foot is beyond dispute. The skin, prodigiously thickened, hangs in wide baggy folds about the leg and ankle, so that the toes can barely be detected peeping out from under them.

The subject of the case was a young married woman, aged thirty, who was sent up to my wards in the Meath Hospital, from the county Cork, 25th June, 1877. Notwithstanding the size of her left leg and thigh, she was able to stand, walk, and attend to her family and household duties. She could easily move the joints of her toes, ankle, knee, and hip, had full sensibility, and no pain in the limb, and was only inconvenienced by the weight of the massive growths which hung from the femoral region, and by the difficulty of sitting down, in consequence of the protuberance of the principal of these masses. She led a very active life in the country, having five children to look after; she rose at 7 a.m., and

was on her feet till 9 p.m. The principal reason for which the doctor who had seen her at home advised her to come to Dublin was the sake of the rest in bed which could be enforced in hospital, and which, when carried out only for a few days, he noticed to be beneficial. This woman had never been out of the county Cork until her visit to Dublin. Neither of her parents, nor any of their relations, ever had anything similar. She herself always enjoyed good health, and had a strong constitution. She possessed a good appetite, and slept well, and had had a child four months before she came to Dublin, which she nursed up to three days before her admission.

The disease had existed for nine years when she came under my observation. Her history of its origin was that a week after having got a severe wetting she felt a sharp pain in the groin, which ran down the front of the limb to her toes. The pain was attended with a swelling of the limb, which continued after the pain (which lasted for a few days) had gone away.

Since that time the limb had never regained its proper size. At the time the pain was felt she did not observe any enlarged glands in the groin, but has often done so since then, and always in the left groin. She was married twelve months after the commencement of the disease, at the age of twenty-one, and had had seven children, including twins once. Five of her seven children are living. The twins, which were seven-months' children, both died soon after birth. She had no difficulty in any of her confinements, and nursed all her children but one. During each pregnancy the limb used to get larger, and become reduced in size immediately after delivery. It swelled much more than usual when she was carrying the twins. *From the commencement of the disease up to the date of her marriage the whole limb was continually swelled, and used to pit on pressure, but, as it did not interfere with her movements, she did not mind it at all. During the last two or three years, she said, "things like small boils" appeared more than once, generally just before confinements, when her leg was most swollen. These "boils" used to first discharge matter, and afterwards water. They generally dried up in less than a week. The discharge from these "boils" used to reduce the size of the limb, but at the same time weakened her. On one occasion she got a scratch from a brier on the lower part of the leg, which gave exit to a quantity of water, and temporarily lessened the size of the limb. On three or four occasions she had had erysipelas; the last bad attack was three

Engraved by John Pollock, Boston.

From a Photograph.

DR. A. W. FOOT.—CASE OF PACHYDERMATOCELE.

DR. A. W. FOOT.--CASE OF PACHYDERMATOCELE.

years before admission, and lasted for three or four weeks. At such times she could not move or lift the leg.

The following measurements were made 26th June, 1877:—

	LEFT.	RIGHT.
	Inches.	Inches.
Base of toes - -	8 $\frac{1}{8}$	8 $\frac{1}{2}$
Middle of foot -	11	8 $\frac{7}{8}$
Above ankle joint -	19 $\frac{3}{4}$	8 $\frac{1}{2}$
Lower third of leg -	22 $\frac{1}{4}$	9 $\frac{1}{2}$
Calf of leg - -	26 $\frac{3}{4}$	12 $\frac{1}{4}$
Below knee - -	25 $\frac{1}{4}$	11 $\frac{3}{4}$
At knee - -	30 $\frac{1}{2}$	13 $\frac{1}{4}$
Middle-third of thigh	49 $\frac{1}{2}$	14 $\frac{1}{2}$
Upper third of thigh	23 $\frac{1}{2}$	17 $\frac{1}{4}$

Notwithstanding the enormous bulk of the left limb, she carried it off very well, and when she was up and dressed the deformity made surprisingly little show. She walked about, and up and down stairs, without a stick or assistance of any kind. In standing, "the small leg" (right) fitted into a groove in the large one. When she lay on her right side (her usual position) the right thigh was completely hidden by the cutaneous overgrowth connected with the inner surface of the left thigh, while the masses springing from the outer aspect of the left thigh raised the bedclothes in a high mound, and those connected with the posterior femoral region lay stretched out on the bed far behind her. The gluteal region was quite unaffected, as also were the organs of generation. Her principal difficulty was in sitting down, owing to the inconvenient protuberance of the mass attached to the posterior femoral region. Her most comfortable position was on a low stool, in which attitude the large posterior mass rested on the ground, from contact with which the skin there was rough and scabrous. Defecation and micturition, as she could not sit down on any utensil, were managed by holding out of the bed-post and discharging into a vessel placed behind her. In her long railway journey to Dublin she lay along the seat of the carriage the whole way.

The skin of the diseased leg was in general smooth, except where it came in constant contact with her clothes or the ground in sitting, and here it was scabrous and warty, and covered with thorny projections, which made it look like an ox's tongue; the holes of the sweat follicles were very wide and stretched far apart, and the

intervening areas glazed. There were many deep sulci, or creases, where the lines of flexion existed; in such parts the skin was soft and moist, and a very heavy rancid smell was exhaled from these fissures, though she kept these places washed as well as she could.

As neither the patient nor her husband would hear of any operative interference with the limb, she was retained in hospital more as a curiosity, and for the sake of observation, than for any other object; but, meanwhile, rest in bed, elevation of the limb, and compression with elastic bandages, were constantly employed, while diuretic medicines were exhibited. The effect of this treatment was soon apparent in a great relaxation of the principal mass which was attached to the posterior femoral region, which, from having been almost as firm as a camel's hump, became a flabby pendulous appendage—the dermatolysis of English writers. As she was assured that this would fill up and get firm again when she got up and discontinued the treatment, she consented, after much hesitation, to have it removed; and we were in hopes, if the operation were successful, and that a firmly-contracted cicatrix were substituted for loose folds of relaxed integument, that at another time a second large portion of the redundant integument might be removed. During the performance of the operation the venous hæmorrhage was unexpectedly profuse, owing to the division of the almost plexiform series of large veins which ramified in the subcutaneous layer, and this loss of blood, combined with an unusual degree of chloroform-sickness, and much depression from shock, acting on a very nervous temperament, contributed to bring about a fatal issue on the fourth day after the operation.

The portion of integument removed weighed many pounds; it presented evident hypertrophy of the cuticle, the corium, and the subcutaneous cellular tissue. The numerous blood-vessels, chiefly veins, gave the deeper parts of the structure the appearance of almost a cavernous metamorphosis, their lumina resembling the channels excavated in worm-eaten wood.

In reference to the name of this disease, which, though so easily recognised, has been hampered with a multiplicity of titles, there is no doubt of the synonymy of Pachydermia,^a Pachydermatocèle,^b Dermatolysis, Elephantiasis (Arabum), Spargosis,^c Barba-does leg. There is a good drawing of the pendulous form of the

^a Wagner. *Man. Gen. Path.* 1876, p. 383.

^b *Med. Chi. Trans.* Vol. XXXVII., 1854.

^c E. Wilson. *Lects. on Dermatology.* 1874-5, p. 173.

disease, called Pachydermatocele by Professor Valentine Mott, in a paper by Mr. Stokes,^a and another in a paper on two cases of Dermolysis, by Dr. G. Fritsche.^b

It is noteworthy that the patient here referred to attributed the origin of the disease to a severe wetting. The Chinese believe this disease, which is common at Ningpo, to be occasioned by wading in the mountain streams chilled by ice. In 340 cases of elephant leg, Mr. Waring found the average size round the ankle to be $11\frac{7}{12}$ inches; in this patient it was $19\frac{3}{4}$ inches. The disease is more common in males than females. According to Mr. Waring's observations, 75 per cent. of cases were males and 25 females.

ART. II.—*On the Consolidation of Internal Aneurisms.* By JOLLIFFE TUFNELL, M.R.I.A.; Consulting Surgeon to the City of Dublin Hospital; &c.

IN a comparatively recent number of this Journal^c the following opinion has been advanced in reference to the cure of aneurisms—viz., “That the layers of lymph deposited in the sac of an aneurism are formed by the walls of the sac itself, and are not a deposit from the blood.”

Sorry as I am to be obliged to differ from so distinguished a surgeon as the writer of this article, still I cannot admit the correctness of any one of the arguments which he has advanced in support of his views; and I must maintain and hold to the fact (as demonstrated in every pathological museum) that the consolidation and cure of an aneurism is not the result of any secretion from the sac itself, but a purely mechanical deposit—rapid when the sac is occluded by clots, and slow and progressive when consolidated by fibrin deposited in successive layers.

I affirm that in neither case is inflammation present, and that in no instance does lymph (the product of inflammation) act as the agent, or constitute the medium through which cure and recovery are brought about.

In the case of an aneurism treated by ligature, where the circulation is suddenly and almost entirely cut off, and rapid consolidation

^a Dub. Journal of Med. Science. Vol. LXI., p. 1.

^b Clin. Soc. Trans. Vol. VI., p. 163.

^c Vol. LXIV., No. 70, October 1, 1877. On Aneurismal Sacs. By William Colles, Surgeon to the Queen in Ireland; Regius Professor of Surgery in the University of Dublin; Surgeon to Dr. Steevens' Hospital, &c., &c.