some hours, and the poison have already reached the intestinal canal, and no alvine evacuations have occurred, 50 or 60 grammes of castor-oil may be administered, and assisted with a lavement of water. When it is supposed that the major part of the arsenious acid contained in the digestive canal is neutralised and expelled by the vomiting and by the stools, the attendant should recur to the use of emollient and diuretic liquids in abundance, in order to expel through the kidneys the portion of the poison which may be absorbed and dispersed through the tissues. These liquids, if taken at the first period of poisoning, would have the disadvantage of dissolving the arsenious acid, and thus favouring its absorption. Blood-letting should be employed whenever there is evident reaction, as well on account of the results furnished in these experiments as that its utility in certain cases has been admitted for every age. Tonics and excitants ought to be carefully prescribed, because they are useless, and may be hurtful.

22. We are finally led to believe that medical men will derive great assistance from the aqueous and diuretic medication, employed in *good time*, in the numerous cases of poisoning where the vegetable or mineral poisonous substances may have been absorbed; for then the animal economy is unburthened through the urinary passages of a great part, or of the whole of the poisons absorbed.

TALIPES.

By Mr. J. BRAID.

[From the *Edinburgh Journal*.]

I THINK there can be no doubt of club-foot being hereditary, as I have repeatedly met with two or three in a family; in one instance four, and in another, six club-feet in one family.

Different instructions as to what tendons should be divided in each variety are given by surgeons, and I find an American author, Dr. Chace, is advocating mechanical extension alone. His cases comprise no difficulties, and would have been cured in less time, and with less pain, by the cutting treatment.

Having operated upon two hundred and forty-six club-feet, embracing every variety and every age, from fifty-three years to two days old; and having, in the course of this practice, tried many modes of operation, the results of which I have watched with strict attention, I trust I may not be deemed presumptuous in offering opinions different from others who have written upon the subject, but whose opportunities of observation may not have been so ample as my own, however competent they might have been in other respects.

It appears to me, that it is not enough to

say that any given method should be adopted, because cases have been cured by that plan. The plans must vary according to the cases.

In a case of pure talipes equinus with rigid contraction, it will generally be found that not only is the tendo Achillis and plantaris implicated, but also the flexor longus pollicis pedis and flexor longus digitorum, tibialis posticus and peronæus longus and brevis. I am aware that after dividing the tendo Achillis alone, forcible extension will be adequate to bring the foot into its natural position; but I am certain that it will be after much longer time has been expended, than if all the rigid muscles had been divided at once. Moreover, I confidently affirm that if all are divided, and the foot brought to the natural position within four or five days, sufficiently firm reunion will have taken place to enable the patient to extend the foot. If any of these muscles are relaxed, it would be superfluous to divide them; but as they are generally proportionally contracted, by dividing them, the patient is saved much suffering during the extension, and the danger of relapse is lessened.

The wounds being closed with adhesive plaster, the foot and leg should be bandaged as high as the knee, to retain the foot in its original malposition for a few days, and absolute rest and quiet should be enjoined. Extension should begin in two days, and in three or four the foot may be brought to a right angle with the leg.

For the same reasons in varus, I divide at once every tendon which can possibly retain the foot in its malposition; viz., Achillis, tibialis anticus, tibialis posticus, flexor longus pollicis pedis, and flexor longus digitorum, and abductor pollicis, if required; and should there be much contraction of the sole of the foot, the plantar fascia as well as the short flexors. The wounds being closed, and the foot maintained in its original position, in two days extension may be commenced, so as to bring down the heel, and evert and flatten the foot at the same time. I consider the treatment by *successive* divisions of the tendons as objectionable. 1st, it retards the cure; 2nd, it causes too much new tendon to be inserted between the divided ends of the tibialis anticus, and thus weakens the power of bending the foot; and, 3rd, it renders necessary a second division of the tibialis posticus, flexor longus pollicis pedis, and flexor longus digitorum.

In valgus the peronæus longus and brevis generally require division, and sometimes also the tendo Achillis, peronæus tertius, and extensor longus digitorum, and proprius pollicis pedis. In two days such splints as will maintain necessary extension should be applied.

In calcaneus talipes the tibialis anticus, extensor proprius pollicis pedis, extensor longus digitorum pedis, and peronæus tertius, generally require to be divided. Divide

either all or as many as seem to bear on maintaining the malposition.

Having restored the foot to its normal position, it is of importance to ascertain whether the legs are of equal length. In most cases of single talipes, the affected leg is shorter. This must be compensated by a high-heeled shoe, otherwise the patient will recur to the original malposition.

For adults I use a straight blade one inch and a quarter long, and about the twelfth of an inch broad. The point is cut off at an angle of 60°. This is less apt to tranfix the tendon, and leave any portion undivided, than the sharp-pointed blades. For children a smaller blade will suffice.

I divide the tendo Achillis about an inch and a half from its insertion in the adult, and proportionally less in younger patients, and prefer cutting from without inwards. The tibialis anticus near its insertion in the internal cuneiform bone; the tibialis posticus near its insertion in the scaphoid bone; the flexor longus pollicis pedis, and flexor longus digitorum pedis, at the point of decussation in the sole of the foot, from within outwards; and from this same wound the plantar fascia may also be cut. The peronæus longus and brevis are best divided together above the malleolus, and separately below it near their points of insertion; the extensor longus digitorum pedis before the points of separation into tendons; the extensor proprius pollicis a little more inwardly, and the peronæus tertius in the same line, or near its insertion; all these latter from within outwards, the tendons being kept on the stretch whilst divided.

The machine I use consists of a sole-piece of hard wood (say deal) about three-eighths of an inch thick, of the length of the foot, and breadth of the sole at its greatest width, with an edge two inches and a half deep, to rest upon the inside of the foot. The two pieces are fastened together with screws, and a notch cut in the sole-piece a little anterior to the malleolus externus; and another notch in the side-piece, behind the root of the great toe, to allow the bandage the better to embrace the foot, and make it lie close to the wooden sandal. This should be covered with sheet-wadding, or a pad, and is to be bound firmly on the foot with a roller, or two or three straps and buckles. A piece of wood from two to three inches wide, and sufficiently long to reach from the foot to a little below the knee, with a notch at the bottom corner, is now to be made fast to the foot and sole-piece by means of a bandage, and placed at an acute angle with the sole, so that it may have enough leverage to elevate the foot when the upright stem is carried back to the side of the leg. An assistant is now to be requested to place a little wadding or pad at the inner edge of the leg-piece, and carry it back until it rest against the top of the tibia; the surgeon regulating the exten-

sion to what the patient can bear, by allowing the bandage to give way, more or less, as he judges necessary. He is now to pass the bandage round the foot, ankle, and leg, from the foot upwards, embracing also the leg-piece, with such a degree of tension as is necessary.

The price of such an apparatus is not more than three shillings.

ROYAL COLLEGE OF SURGEONS IN LONDON.

REGULATIONS OF THE COUNCIL

Respecting the Professional Education of Candidates for the Diploma, August 20th, 1839. Amended October 14th, 1841.

- I. Candidates will be required, in addition to a certificate of being not less than twenty-one years of age, to bring proof
 1. Of having been engaged in the acquirement of professional knowledge for not less than four years; during which period they must have studied practical pharmacy for six months, and have attended one year on the practice of physic, and three years on the practice of surgery, at a recognised hospital or hospitals in the United Kingdom;* three months being allowed for a vacation in each year.
 2. Of having studied anatomy and physiology, by attendance on lectures and demonstrations, and by dissections, during three anatomical seasons or sessions, extending from October to April inclusive.
 3. Of having attended at least two courses of lectures on the principles and practice of surgery, delivered in two distinct periods or seasons, each course comprising not less than 70 lectures:—and one course of not fewer than 70 lectures, on each of the following subjects; viz., the practice of physic—chemistry—materia medica—and midwifery with practical instruction.
- II. Members and licentiates in surgery of any legally-constituted college of surgeons in the United Kingdom, and graduates in surgery of any university requiring residence to obtain degrees, will be admitted for examination on producing their diploma, licence, or degree, together with proofs of being twenty-one years of age, and of having been occupied at least four years in the acquirement of professional knowledge.

* By a resolution of the council on the 7th of November, 1839, no provincial hospital will in future be recognised by this college which contains fewer than 100 patients, and no metropolitan hospital which contains fewer than 150 patients.