A Mirror OF HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

KING'S COLLEGE HOSPITAL.

A CASE OF CHRONIC MIDDLE-EAR SUPPURATION WITH
THROMBOSIS OF THE LATERAL SINUS IN WHICH THE
INTERNAL JUGULAR VEIN WAS NOT
LIGATURED; RECOVERY.

(Under the care of Mr. ARTHUR H. CHEATLE.)

THROMBOSIS of the lateral sinus occurring during the course of otitis media may have either of two pathological causes. On the one hand, it may be due to the passage of micro-organisms from the middle-ear through the wall of the sinus; thus a thrombus is formed, which contains the bacteria which have given rise to it; this clot readily breaks down, and therefore in these cases there is a very great probability of the occurrence of embolic pyæmia. On the other hand, the blood may clot in the lateral sinus, owing merely to a "spread of the inflammation" from the surrounding structures; in this case the coagulation is probably the effect of the transudation of toxins, and the clot so formed is firm and has little or no tendency to become disintegrated. It is not possible to distinguish clinically between these two conditions until some infective manifestation occurs, such as a pyæmic embolus; but a rigor is very suggestive and is quite sufficient to justify ligature of the internal jugular vein. A case somewhat similar to the one recorded below has been published by Röpke.¹

A boy, aged 14 years, attended at King's College Hospital on June 26th, 1899, complaining of pain round the left ear and of vomiting. 18 months previously he had had a discharge from both ears which "was stopped." On two occasions since then he had suffered from ear-ache, but there had not been any discharge. Five days before being seen at the hospital he had pain in his head and commenced to vomit. The pain, which was at first intermittent, had become continuous. The vomiting had caused great distress, and as no food could be retained he had rapidly lost flesh. Two days before admission the ear had discharged with some relief to the pain. Giddiness had been complained of for two days. A marked shivering fit occurred during the night before the visit to the hospital. The boy looked wasted and anxious and was extremely restless. He was sensible and answered questions readily. On walking he had a tendency to walk towards the right. The tongue was white, furred, and moist. The temperature was 100.2° F. and the pulse was 104. An offensive brown discharge came from both ears, a perforation being present in Shrapnell's membrane on each side. There was no bulging of the posterior superior deep meatal wall in the left ear, nor did the auricle stand out, and there was no redness or swelling behind it. Marked tenderness on pressure was present behind the mastoid process and below the ear where glandular enlargement could be felt. The heart and lungs were normal. Mr. Cargill reported that papillitis was present in both discs equally.

A few hours after admission the patient was placed under chloroform. The antrum was found to be full of granulations and cholesteatomata. On laying the antrum into the middle-ear the malleus and incus were found, both carious. The attic and middle-ear were then thoroughly cleared. On examining the walls of the antrum a discoloured and softened patch of bone was found in the posterior, leading directly to the lateral sinus. The sinus having been exposed for over an inch of its length it could be seen that it was occupied by a firm clot which was dark in colour except at the point of infection, where it was slightly yellow and evidently breaking down. As giddiness had been a marked symptom

the cerebellum both in front of and behind the sinus was thoroughly explored but with no result. The dura mater, which had been incised, having been stitched up with catgut the sinus was laid open for the whole of its exposed length and the clot was turned out. As the clot for some distance above and below the point of infection was dark-coloured and firm it was decided not to tie the internal jugular vein but to clear the thrombus out as far as possible from the wound. For this purpose a sharp spoon was first carefully passed downwards, no bleeding occurring. A gauze plug having been prepared the spoon was passed backwards towards the torcular; after some clot had been gently removed a free gush of blood took place, shooting out a healthy-looking mass of clot fully one and a half inches in length. The bleeding was easily controlled by the plug. The whole wound was then packed with iodoformed cyanide gauze. The operation occupied one and a quarter hours. An hour after the operation a rigor occurred, the temperature reaching 102.4°. From that time the progress was steady and uneventful but slow. On July 10th, as the hospital was closing, the patient was transferred to St. George's Hospital under the care of Mr. Marmaduke Sheild, who kindly looked after him until he was sufficiently well to be moved to the convalescent home.

Remarks by Mr. CHEATLE.—That ligation of the internal jugular vein is not always necessary in thrombosis of the lateral sinus is well recognised, but no guiding rule has been laid down. Young² has described a case in which recovery took place without ligation of the vein, and no doubt many surgeons have had the same experience. Bronner has related a case in which the clot as seen through the wall of the sinus looked healthy throughout, and it was therefore left unopened with a happy result. It seems that if healthy-looking clot can be seen well below the breaking-down area ligation of the vein is not necessary; but in order that such a condition may be found it must be well recognised that one rigor demands instant operation. A rigor does not necessarily mean that there is septic thrombosis of the sinus, for in a case under the care of Dr. Urban Pritchard in King's College Hospital a severe rigor occurred as a result of an acute empyema of the antrum, and no further bad symptom followed after removal of the outer wall and evacuation of the pus. But it cannot be too strongly insisted upon that there should be no waiting after a rigor has occurred during the coming of an acute or chronic middle-ear suppuration; the antrum should be at once opened and the sinus examined. An interesting point about the case related is that although the boy was perfectly well the optic neuritis was present at least three months after the operation. The giddiness which was such a marked symptom was due in all probability to disturbance of the intracranial circulation owing to the occlusion of the sinus.

GUYS' HOSPITAL.

A SUCCESSFUL CASE OF REMOVAL OF A PACKING-NEEDLE
FROM THE DESCENDING COLON; RAPID DISAPPEAR-
ANCE OF INTRA-PERITONEAL ADHESIONS.

(Under the care of Mr. N. DAVIES-COLLEY.)

By means of the Roentgen rays much assistance can be obtained in the localisation of a metallic foreign body, but a single skiagram will in many cases not be sufficient to indicate the depth from the surface. A second skiagram taken in a different direction may supply the required information, but the best method is to employ Davidson's localiser, for by its use the exact position can be readily found. Occasionally no trace of a metallic body may be found on a photograph, although the body is certainly present, and a good instance of this was described in THE LANCET of Feb. 5th, 1898, when Mr. J. H. Morgan removed a piece of stay busk from the stomach of a man; no skiagraphic shadow of the piece of metal was found because it had been lying in a position perpendicular to the sensitive plate. The rapid disappearance of peritoneal adhesions which occurred in the following case is certainly remarkable.

A married woman, aged 36 years, the mother of nine children, the last of whom had been born 15 months, was on April 12th, 1899, admitted into Guy's Hospital under the care of Mr. Davies-Colley with obstinate constipation and a painful swelling above the front half of the left iliac crest. Her

¹ Archives of Otolaryngology, vol. xxv., No. 4.

² Glasgow Medical Journal, October, 1899.