low on account of personal peculiarity :—E. F—, aged thirty-four years, mother of five children, became hemiplegic one week after her last confinement, eighteen months ago. She is a peculiarly fragile-looking woman. Late rigidity is taking place. The following resistances were observed on the extensors of the forearms :—

CASE 4.			
	Right or healthy side.		Left or diseased side.
Epidermis per square inch, before moistening	1959 ohms.		1981 ohms.
Epidermis per square inch, after moistening	125 „		190 „
Tissues beneath epidermis, electrodes being 3 in. apart, before the passage of the current . . .	2353 „		1420 „
Tissues beneath epidermis, electrodes being 3 in. apart, after the passage of the current . . .	321 „		164 „

Here we may safely conclude that the nerves regulating the bloodvessels are unaffected, and that the tissues themselves are the parts diseased.

The next division of my subject is upon the advantages derived from the employment of a galvanometer. We can place one of the electrodes upon any convenient part of the body, not necessarily in the central line, and the second over the nerve or muscle, the irritability of which we desire to know. To ascertain the current flowing at any time the only thing we have to do when using the author's galvanometer is to press down the key and read off the deflection by the scale (which can be done almost instantaneously), and to multiply the number of the division of the scale at which the beam of light rests by '025 or '050, &c., according to the shunt used. The result will then be in mille ampères. It is not even necessary to wait until the skin is soaked, because when it is dry we have merely to throw into action a larger number of cells in order to produce a current of the same strength. In the current we usually insert a fluid regulator to act as a fine adjustment. It is well to recollect that until the skin is completely soaked and the current has been passing some short time, if we measure its strength at successive intervals, we shall find it stronger and the muscular contractions more violent, because the resistance of the tissues becomes less and less, and this teaches us that we ought to measure the current by the galvanometer immediately we have regulated its strength. In health, with the exception of the muscles of the face, we find that under most circumstances from 1·5 to 2·0 mille ampères are required to produce muscular contractions with the negative pole upon closing of the current, and from 3·5 to 4·0 mille ampères by the positive pole. For the face about half these quantities. (We omit for special reasons the amounts required in the opening of the current.) In health the only common noteworthy exception to the above is that a stronger current as measured by a galvanometer is necessary by very stout people. To understand the reason for this we must fully realise the following facts:—1st, that the current is strongest just beneath the electrodes, and that it diffuses itself not only in straight lines, but in an infinite series of curves in every direction; 2nd, that these curves would be regular in every direction if the intervening structures were electrically homogeneous, but as this is not the case they follow the law of divided circuits—viz., that the strength of the current in each circuit is inversely proportionate to its resistance. Thus in the body the curves are anything but regular. The following is the main course of the current. The epidermis conducts a very small part of the current either in a straight line or in curves; of the remainder the corium and the subjacent adipose tissue transmit by far the larger quantity, allowing only an exceedingly small portion to react upon the muscle or nerve. The current that arrives at the muscles may be regarded as proved to be very weak, from the fact that when the nerve or muscle is excited directly, either by the insertion of needles or by the electrodes being placed upon them when laid bare, one cell with a very large extra resistance is sufficient to induce very violent contractions. Now, as the proportion of the current which penetrates deep enough to reach the muscle depends both upon the conducting power and upon the thickness of the strata intervening between them and the electrodes, and as a certain strength of current is absolutely necessary to cause muscular contractions, we can readily understand why it is that when the tissues covering the muscles are abnormally thick, or better conductors than usual, a stronger current than ordinary is needed to obtain a response from the muscles.

The question now arises how are we to know whether the muscles are healthy or not, in cases in which the conduction of the overlying structures is such as to prevent a current of the normal strength from exciting them. The answer is that when they do not respond to a current having a strength of from 1·5 to 2·0 mille ampères negative pole and of from 3·5 to 4·0 mille ampères positive pole, we shall find during health they still contract, having the same ratio—namely, about 3 to 7. In disease we consider the ratio of the current strengths requisite to cause positive and negative contractions respectively, as more important than the absolute strengths of the current themselves. I have also observed that in the majority of cases when the muscle regains its power, the currents needed for exciting contractions, with the negative and positive pole respectively, gradually approach the ratio 3 to 7. Although in many cases for some time at least, even if not always, will the strengths of the necessary currents be unlike those wanted before the disease commenced. For an illustration, I will quote a case where the needful strength of the current remained high, although the ratio became natural.

A. B— fell down and struck his head on the right side, immediately losing the use of his left arm. One month after, when I first saw him, he was able to use all his muscles with the exception of the deltoid. This muscle responded to 4·8 mille ampères – pole and 3 2 mille ampères + pole, ratio being 3 to 1·99. Three days afterwards, and remaining for twelve days, the current required was 6·5 mille ampères – pole and 5·0 mille ampères + pole, ratio being 3 to 2·31. There was during this time some slight improvement. One month later the requisite currents were 5·75 mille ampères – pole and 13·55 mille ampères + pole, ratio being 3 to 6·89. The patient was now so well that he did not return. It is only by the help of a galvanometer that we are able to discover with any degree of accuracy the ratio between the two currents, and certainly without it we are absolutely unable to observe daily fluctuations. Besides, very few batteries are constructed so that single cells can be set into action. Even when they can the valve of a cell varies. First, according to its E. M. F.; second, according to the number of cells already in action; third, according to the resistance of the body. The size and position also exert a very powerful, but not uniform, influence in regulating the third consideration, the only one I shall notice. Suppose that the electrodes are placed so that the body exerts a resistance of 3000 ohms, and taking a battery of Leclanché cells each having an E. M. F. of 1·5 volts, and an internal resistance of 10 ohms, then 10 cells would give a current of 4 878 mille ampères, while if the resistance of the body were 4000 ohms, 10 cells would give a current of 3 656 mille ampères only.

In disease we rarely find that the resistances of the body are alike on the two sides, so that the same cells have different values. For example, let us take Case 3, C. If 10 cells were used on the right side, 11½ (or by the ordinary method 12) cells would be needed on the left for the production of the same current. Here calculations based upon the number of cells would give an error of from 15 to 20 per cent.

In conclusion, there is not the slightest doubt that medical electricity has not made in the last decade the same strides as other branches of electricity, and this can only be attributed to the fact that medical men have neglected the instruments of precision which have done so much for the other branches of the science. Although mathematical accuracy is unobtainable in medicine, yet our motto ought to be “measurement.”

Ladbroke Grove-road.

A CASE OF EPILEPSY.

OBLIVIOUSNESS OF DANGEROUS ACTS; MEDICO-LEGAL BEARINGS; VALUE OF PERCUSSION OF THE SKULL.

BY A. ROBERTSON, M.D., F.F.P.S.G.,
PHYSICIAN TO THE TOWNS HOSPITAL AND CITY PAROCHIAL ASYLUM, GLASGOW.

THE following observations are largely based on a case which lately occurred in this hospital, of which a careful record was made by my late assistant, Dr. Williams, now of the Denbigh Infirmary. His report is as follows :—

W. R—, aged forty-five, a soldier, was admitted on Jan. 25th, 1882. Five years before admission he had a sun-stroke in India. Epileptic seizures set in shortly afterwards,

and recurred at intervals of from a month to six weeks till the time of his admission into the hospital. Under a combination of the bromide and iodide of potassium, half a drachm of the former and ten grains of the latter, thrice daily, the intervals were somewhat prolonged. On Feb. 23rd of this year he had a convulsive fit, which left him irritable and quarrelsome, but apparently quite conscious of all his actions. On the 24th, while in this mood, he suddenly attacked and knocked down an old man who was passing him in the ward, though there had not been the slightest quarrel between them. Next day he denied having struck him, and at the same time said that he had no recollection of anything that happened on the previous day. His whole bearing in saying so impressed me with the truthfulness of his statement. There was no further violence nor recurrence of the fits, but he was unsociable and somewhat obtuse. He complained of pain in the head, especially on the left side and anteriorly. The pupils were slightly dilated. On April 14th slight shivering occurred, and the axillary temperature was found to be 103° F. Headache was severe, and still most pronounced on the left side. I percussed the skull carefully. The patient stated that the taps of the finger did not cause pain on the right side or back of the head, but that they were very painful on the left side, in front of the ear, forward to the middle of the brow. Leeches were applied over the painful region, and mercurial inunction prescribed. The symptoms, however, became more serious, and he died comatose on May 16th.

Necropsy.—There was about an ounce and a half of serous fluid under the arachnoid and in the ventricles; within the dura mater, on the left side, over the entire frontal lobe, and extending downwards to the base, there was a thin organised false membrane, which seemed to have no adhesions either to the dura mater or arachnoid. The brain was firm; the specific gravity of the white substance in the frontal region was 1.047.

Remarks.—Much attention has been given by Dr. Hughlings-Jackson and others to the mental automatism which not unfrequently occurs in epilepsy, particularly after attacks of petit mal. In these cases purposive acts are performed, sometimes nearly correct in themselves, but at improper times and places, and without the consciousness of the actors. But there is a condition following such seizures in which the mental defect is even less than that just stated, where consciousness is only very slightly impaired, where there seems only a little confusion of mind, scarcely noticeable to superficial observation. The remarkable feature of this state is that the patients may maintain a conversation, apparently correct or nearly so, and perform acts, of which they have not the slightest recollection a few hours afterwards. This is illustrated by W. R.—'s case. He knocked down his fellow-patient without the least provocation, though seemingly aware of what he was saying and doing, both then and afterwards in the course of the day; yet next day he declared, and I believe truthfully, that he had no recollection not only of that act of violence, but also of all the other events of the next few hours. The question suggests itself, Had he killed the old man at the time he struck him, ought he to have been held responsible for his death? I think not; at least, not fully so. A few months ago I was engaged as medical witness (case of George Miller) in a trial for murder by an epileptic, in which the accused, like W. R.—, declared that he had no remembrance of the fatal act, nor of what happened, for at least half an hour afterwards. The main facts of the case are as follows:—G. Miller, a soldier, who had been in the Egyptian campaign, was discharged on Dec. 30th, 1882, on the breaking up of his regiment in a small town about ten miles from Glasgow. He drank freely in the course of the afternoon, at night reached the city by train, when, being unable to take care of himself in the streets, he was conveyed to the district police-office. There he was able to tell his name, &c., and walked upstairs to a cell with assistance. He was laid on a fixed wooden bed beside another drunken prisoner, who was asleep at the time, and locked up with him about 8.15 P.M. The latter was stated by those who knew him best to be very inoffensive, and not quarrelsome either in drink or out of it. They were seen by the warder five different times at short intervals, the last time being at 10.45 P.M., and always found asleep. Twenty minutes later, when again visited, the accused was sitting at the side of the cell, with his arms crossed, and his comrade in the middle of the floor, lying with his face down, dead. The head of the deceased was fearfully bruised, and some of his ribs

were broken. The conclusion arrived at was that the soldier had jumped on him or kicked him to death with his heavy boots. The gaoler stated that the accused said, on being questioned respecting his comrade's death, he knew nothing about it, and while saying so appeared cool and collected, and so correct in every way that, had nothing happened, the officer in charge would have allowed him to go free. He was taken to the bar and formally charged with the crime of murder; the lieutenant on duty testified that he was apparently quite conscious of what was said to him. Thereafter he was removed to a cell, where he spoke for a short time freely about his Egyptian experiences to the officer appointed to watch him, and then fell asleep. He slept soundly for nearly five hours. When he awoke he was told about his crime, but he appeared quite incredulous, and declared that he had not the least recollection of it or of any of the after circumstances. It was ascertained that he had been subject to attacks of petit mal, and that on one occasion he had a convulsive seizure. He had also been convicted of inflicting a severe blow on a woman, an entire stranger to him, who was passing him quietly in the street; he denied all knowledge of this assault also. He had likewise about two years previously made a determined attempt at suicide.

This history points to the fatal attack as having been of an epileptic character. No doubt the prisoner had previously been indulging in alcoholic liquors, but this does not satisfactorily account for his condition when first seen after the homicide. The seeming cool composure and indifference which he then displayed, the failure to realise his position, and the forgetfulness of the act, as well as of the events of the next half hour, all indicated a really imperfect consciousness, even though to ordinary observers he was apparently fully aware both of what he was saying and doing. At the same time the epileptic seizure, which I believe occurred on his awakening from the drunken sleep, was itself probably due to the action of the alcohol on his weak and susceptible brain. It only further remains to be said that the jury found the accused to have been of unsound mind at the time he killed his comrade in the cell, and he was sentenced to be confined in the Criminal Lunatic Asylum during Her Majesty's pleasure.

It is not improbable that the objection may occur to some minds that, after all, the chief actors in both of these cases may not have been so oblivious as they alleged and appeared to be, it being so very much their interest, and especially G. Miller's, to deny all recollection of their criminal acts. Such objectors will, I think, admit that the following case, very similar to the others so far as the mental condition is concerned, is free from that doubt. A gentleman, aged fifty, whom I see occasionally, has been subject to the minor, and much less frequently the major, attacks of epilepsy for several years, the intervals between the seizures varying from a week or two to some months. Though mentally not now quite equal to his former self, he is still active and intelligent, and, what is specially to be observed, possessed of fairly good memory. On several occasions he has had attacks of petit mal when attending to his occupation, and has afterwards found his way downstairs and through busy streets, returning quite correctly to his place of business, but entirely unconscious of all that had happened in the intervals, lasting for from five to twenty minutes. It is not known how far he may have seemed intelligent and correct to others while walking in that condition. What I have next to mention is, however, more to the point under consideration, and was told me by his wife, a highly intelligent lady. One afternoon, about four months ago, while he was seated on the sofa at home, she observed by the movement of his lips and of one of his hands, that he had taken one of his "turns." The ordinary medical attendant of the family was sent for, and arrived within a quarter of an hour. By this time the patient was apparently quite conscious, and conversed correctly with his medical friend for about five minutes. About half an hour after the latter had left the house, some reference was made by the patient's wife to the doctor's visit and his instructions, when she found that he had no recollection of his having seen him, and indeed positively insisted that the doctor had not called that day.

In W. R.—'s case the exact correspondence between the painful area on percussion of the head and the extent of the false membrane within the dura mater, so far as it could be ascertained, was very striking. Would it not have been warrantable, as my friend Dr. Macewen, of this city, suggests to me, to have trepanned the skull and removed the false membrane? In the light of the necropsy

this might have been done, and with probable advantage, as the irritation due to its presence was, I think, the immediate cause of death. In any future case of a similar kind, in which percussion clearly defines a painful region, I should be disposed to try trephining, in the event of no improvement resulting from ordinary treatment.

As the method of diagnosis by percussion of the skull is still comparatively new, I may mention that there is at present in this hospital a patient in whom it revealed a painful area over the motor region of one side of the brain. He had been long subject to convulsive seizures, mainly unilateral, and has greatly improved since the application of a series of blisters over this region.

Glasgow.

ON THE USE OF CARBOLISED SAWDUST AS A DRESSING IN ANTISEPTIC SURGERY.

By H. P. SYMONDS,
SURGEON TO THE RADCLIFFE INFIRMARY, OXFORD.

ONE of the drawbacks of the usual antiseptic dressing is the rapidity with which the discharges come through on the first day or two after operation, often necessitating the re-dressing of the case within a few hours. To prevent this, and yet not to interfere with the aseptic condition of the wound, is a distinct advantage both to the patient and the surgeon. The material I have used recently in a considerable number of cases is coarse sawdust, soaked in (1 in 10) solution of absolute phenol and spirit of wine, then allowed to dry slightly so that the spirit may evaporate, leaving the sawdust charged with carbolic acid. When used it is enclosed in a bag made of several layers of gauze, and applied outside the deep dressing, the usual external dressing being put over it. The sawdust thus takes the place of the padding of loose gauze which is generally used. Its absorbent power is very great, and it has the additional advantage of keeping up an equable pressure on the divided tissues. I find that fourteen ounces of sawdust will readily absorb about one pint of fluid.

The following cases in which this dressing has been used form a successive series, taken without any selection.

CASE 1.—Amputation through the middle of the arm for disorganisation of the elbow-joint and necrosis of the humerus. The operation was done on July 24th. The stump was dressed at the end of twenty-four hours in order to remove the drainage-tube. On July 30th the dressings were changed and sutures removed. The wound was quite healed, with good union throughout. The patient was discharged from the infirmary seven days after operation.

CASE 2.—A woman, aged sixty-one, with extensive scirrhus of the breast. On August 10th the whole breast was removed, and the axilla cleared of glands. Two drainage-tubes were put in the wound. Dressings changed on the 12th to remove the tubes, and again on the 17th, when some of the sutures were taken out. On the 19th the wound was thoroughly healed, and the patient left the infirmary.

CASE 3.—A man, aged forty-eight. Amputation of forearm for large sarcoma in the back of the hand. Operation done on August 3rd. Drainage-tube removed in twenty-four hours. Wound dressed again seven days after the operation, when complete union had taken place.

CASE 4.—A middle-aged woman, with compound dislocation of the elbow, admitted August 11th. She had a small wound on the inner side of the right elbow. Both bases were dislocated backwards, and the inner condyle chipped off. The joint was dressed antiseptically, a counter opening made, and a small drainage-tube passed through. This was removed in twenty-four hours, and the dressing not changed again for four days. No soaking through took place, and the wound remained aseptic. The highest temperature 99°4'.

CASE 5.—Removal of an adenoma from the breast of a girl aged eighteen. This case occurred in private practice. Drainage-tube removed in twenty-four hours. Dressed again on the seventh day. The incision was quite healed.

In all these cases complete primary union took place without any formation of pus. In only one did the temperature reach 100°, and that on the day after operation, after

which it became normal. I have not quoted these cases as being at all remarkable, but merely as common instances in antiseptic surgical practice in which the sawdust dressing was used. Surgeon-Major Porter, in "The Surgeon's Pocket-book," states that he has used sawdust as a dressing in suppurating offensive wounds; but I am not aware that it has been tried, when prepared in the way I have described, in antiseptic dressing. The three points in its favour are its powerful antiseptic qualities when saturated with carbolic acid, its great absorbent power, and its adaptability to any surface. I may add that the sawdust should be coarse, as I find that if it is very fine it passes through the gauze and irritates the skin.

Oxford.

ACUTE RHEUMATISM AND PHTHISIS.

By J. GORDON BLACK, M.D. LOND.

IN THE LANCET of July 7th Dr. Austin calls attention to acute rheumatism as a premonitory symptom of phthisis, and gives the histories of four cases in illustration. The following particulars of a case attended by me last year may perhaps be of interest in connexion with this subject.

On March 6th, 1882, I first saw a pale, delicate-looking young woman, aged twenty. Both father and mother were dead, the former of phthisis, and a sister was then suffering from that disease. My patient had been doing rough hard work beyond her strength in a cold damp atmosphere, for some months. At the previous Christmas she caught cold and a cough ensued. She had, however, been away from work only two days. The symptoms were on examination:—Short dry cough with pain under left axilla, sickness and vomiting of food, sleepless nights, and free perspiration. Pulse 100, weak, and pupils dilated. Tongue red in middle, furred at the sides. Temperature only slightly above, and respiration, normal. Menses regular, but discharge too pale. Some rough breathing below left clavicle; no dulness or crepitation. No increase of cardiac dulness, but sounds unduly accentuated. At this visit I was undecided as to the exact nature of the attack, fearing some lung mischief or impending febrile disorder. A diaphoretic and expectorant mixture was prescribed and the affected side ordered to be poulticed. Next day I found that the cough and sickness had entirely left her. The pain was still bad in the side, and perspiration continued. Temperature 102°, pulse 100. A soft murmur was now heard with the first heart sound, loudest over the pulmonary cartilage, slightly conducted towards the apex, but not heard over the second right cartilage. The right knee and ankle were swollen and tender. Thus the affection had developed into rheumatic fever, and the signs of lung implication suddenly disappeared. Salicine treatment happily soon gave relief to the distressing symptoms, and by the second day of its use the patient was comparatively well. The cardiac murmur became much less audible, but never went entirely away. At the end of March the patient, being rosy and plump, in complete contrast to her previous condition, was allowed to go to a farmhouse in the country.

On April 17th, a few days after her return, I saw her again, and was shocked at the serious change which had occurred. She was now in bed, complaining of excessive muscular weakness. The cough, fever, and vomiting of food had returned worse than ever. Some dulness was now noted in both infra-clavicular spaces, and at the top of the sternum. Pectoriloquy and mucous râles were heard on the left side below the clavicle, whilst on the right there was double respiration. Expectoration was copious and nummular. Pulse 120; temperature 101°. By April 27th the above symptoms had become greatly aggravated. The expectoration was tinged with blood, and there were signs of a vomica at the top of the left lung. For the next few weeks the disease progressed with the fearful rapidity of so-called "galloping consumption"; but afterwards, curiously enough, became quite chronic. The appetite had never failed, and from first to last colliquative diarrhoea was absent. Thus were the powers of nature sustained until the patient had become a veritable skeleton. Death occurred on the 24th of November.

The succession of symptoms above detailed arrested my attention as being unfamiliar, and I was therefore much interested to observe that Dr. Austin had recently attended four somewhat similar cases. Judging by my own expe-