out, in the absence of the peculiar whitish-pink, abundant expectoration. It seems to me that the difference in the two conditions may lie in this—that in acute spasmodic asthma it is the pulmonary artery, or afferent system, that suffers from spasm, and in acute suffocative pulmonary oedema, it is the pulmonary vein, or efferent system, which is affected; for in the former case there is anæmia of the lungs with no expectoration during the attack, and in the latter there is engorgement, with pints of it. In other words, in acute spasmodic asthma ("dry asthma") the lungs are in much the same condition as the fingers are when they are cold, white, and "asleep," or as in Raynaud's disease. In acute suffocative pulmonary oedema ("wet asthma") they are, perhaps, in a similar condition to the extremities in certain forms of erythromelalgia. I think, therefore, one may reasonably argue that the two diseases are similar, but that they differ in the site affected by the provocative agent, whatever it may be. Both clinical pictures may be ascribed respectively to vaso-motor spasm of the different pulmonary vascular systems.

I shall certainly rely with confidence upon chloroform as a remedy for both conditions in future, for it seems to me from the observation of my case that great damage was done by sedative drugs and also by stimulation by alcohol, and none of the vaunted remedies for asthma did my patient any good. She seemed to be drifting into a condition of imbecility, like an epileptic does, until the systematic administration of chloroform stopped the violence of the attacks. She still has attacks—about two a month (instead of three or four a week as in March last)—it is true, but as the duration of them is so short and the remedy inspires such confidence she feels sure eventually she will completely recover.

Bolton-le-Moors.

TWO CASES OF ACUTE DELIRIUM.¹

BY R. H. COLE, M.D., M.R.C.P. LOND.

CASE 1.—The patient was a single woman, aged 40 years, living with her widowed mother. For the sake of employment and to supplement her income she had lately taken some work. There was no family history of insanity, but both the patient and her mother were decidedly nervous, and a brother had died from hip disease. Until nine months before she came under my observation she had enjoyed fairly good health. She then developed a bad attack of pernicious anæmia, from which, however, she recovered at the end of three months and returned to her work. On Feb. 21st, 1900, she again began to feel ill and prostrate, and she feared a relapse of her former illness. This time, however, she was more nervous and apprehensive, complaining of vague cranial uneasiness, and she said that she felt that she was "going off her head." The blood was examined again; it was found to present signs of pernicious anæmia and she was admitted for treatment into a nursing home. The nervousness and restlessness increased and in a week's time, after a few hours' warning, she became completely delirious which, on March 4th, necessitated her removal to an asylum where she was under the care of Mr. G. E. Mould. Before the delirium developed she had three sleepless nights. The patient now presented a wild appearance; she had no powers of recognition whatever and chattered incessantly, her speech being incoherent and the same phrases being repeated over and over again and she asked questions which she proceeded to answer herself. She gesticulated and was under the influence of visual and aural hallucinations. At times she had an ecstatic expression, apparently believing herself in heaven with angels flitting about. She had to be fed artificially and her bladder and rectum required constant attention. In spite of sedatives it was difficult to procure any sleep. The pulse was weak, varying between 120 and 140, the temperature being 99° F. There were a dry cracked tongue and sordes on the lips. The patient was given abundant nutriment and stimulants but she was evidently becoming exhausted. On the 9th her temperature rose to 105° and she seemed to be in a moribund condition. She was ice-packed, a hypodermic

injection of strychnine was administered, and the stimulant (brandy) was increased. She rallied somewhat and the temperature was reduced, only to rise again on the following day to 107°. The patient was then placed in an ice-cold bath for half an hour which reduced the temperature to normal for a few hours. On the temperature rising once more she was placed on an iced water-bed and was given quinine and strychnine by the mouth and meat extracts and stimulants. She responded but feebly to these agents; the delirium had now become of a low muttering form and she picked at the bed-clothes. She died on the 11th in a state of collapse. Unfortunately no post-mortem examination was obtained.

CASE 2.—The patient was a young professional man, aged 27 years and single. An aunt had suffered from chronic insanity. With the exception of an attack of influenza a few years previously the patient had always enjoyed excellent health. He was by nature sociable, athletic, and clever, and had led a regular and temperate life. He was about to begin his life's work, having just finished his final examinations. He seemed unduly to feel the responsibility of his office, and some irritability and change of disposition were observed by his friends for a fortnight or more. On April 3rd, 1901, he first complained of not feeling well, being depressed and unable to collect his thoughts; however, he continued at his work until the 6th, with sleepless nights, when one of his friends gave him a Dover's powder and sent him to bed. On the 7th his manner was so strange that an attendant was engaged. He looked very dejected and would hardly speak; he gave expression, however, to delusions that he had neglected his work and ruined his family and he was somewhat suspicious about his diet. He could obtain no sleep without hypnotics and five days later (on the 12th) he was transferred to an asylum for special treatment. He would not eat an ordinary meal, but after persuasion drank milk on the first two days after his admission. From the 14th to the 21st he had to be fed with a tube (nasal). He was now in an exceedingly resistive mood, champing his teeth, refusing all food, and struggling during the dressing and when undressing or when going out for exercise and returning to his room; he now, however, slept well and without any hypnotics. On the 22nd he took his meals voluntarily, and on the next day he had a lucid interval which commenced suddenly and lasted three hours; during this interval he stated that he had a vague recollection of being moved to the asylum, but could scarcely remember anything more. He gradually became more restless and inattentive, and in the course of an hour he had drifted into a semi-stuporous condition, but he was quite tractable and took his food well. He would not speak except to utter emotional phrases, and any reference to his friends or relatives increased his agitation. In this state he remained for five weeks. On the night of May 28th he was sleepless and he looked ill on the next morning, his temperature being 100° F. and his pulse 120. He became very excited, struggled with his attendants, and took his food only by compulsion. During the next three days the patient was in a condition of acute delirium and evidently under the influence of terrifying hallucinations, chiefly of a visual nature. He occasionally talked, but his speech was incoherent and almost unintelligible. The pyrexia increased to 102°, and his pulse varied from 120 to 150. Sedatives were administered but he obtained very little sleep. From June 2nd to the 8th he lay on a water-bed in a very prostrate condition, with a dry tongue and sordes on his lips, and in order to ensure sufficient nourishment he was again fed by a nasal tube. Six ounces of brandy and four ounces of port wine were now given per diem and sleep was obtained by means of two drachms of paraldehyde every night. His pulse, which had always been rapid, was now intermittent and gave cause for anxiety. On the 5th he had a syncopal attack, but recovered with the assistance of a hypodermic injection of strychnine. The muscular system was in a state of hyper-tonicity and change of position was accompanied by pain. On the 9th the patient again began to take food voluntarily and seemed to be a little stronger. His temperature on this day was 100° and his pulse was 130. On the 12th the temperature returned to normal although the pulse continued to be rapid. In order to ease his back he was now allowed to sit up for an hour occasionally. Mentally he was still in a state of confusion, but he would answer simple questions and appeared to be slowly recovering. On the 15th he woke up in the middle of the night to use his night-stool, and a quarter of

¹ Abstract of a paper read before the St. Mary's Hospital Medical Society on Dec. 11th, 1901.

an hour after returning to bed he complained of feeling very ill and faint. Subcutaneous injections were again administered in order to restore the failing heart, but this time the patient died. A necropsy could not be obtained until 36 hours after death, when the brain presented a congested appearance and there was evidence of minute subpial hæmorrhages. The meninges showed no signs of inflammation. Some of the cerebro-spinal fluid was collected in a sterilised bottle, from which cultures of the diplococcus lanceolatus were obtained. Nissl sections of the brain were made later under the direction of Dr. F. W. Mott, but no pathological changes in the cortical cells could be discovered. The other organs were normal.

Acute delirium is generally regarded as the most convenient term to apply to a somewhat rare but fatal group of clinical cases of which the two narrated above are types and which seem to be worthy of record. In all, practically, there is a predisposition to mental instability, hereditary or acquired. Some observers, especially in Italy, have described a bacillus associated with this condition, and others have found various kinds of cocci, whilst in many cases no organisms can be found at all. It seems probable that the toxins arise autogenetically.

Upper Berkeley-street, W.

THE DECAY OF AUSCULTATION AND THE USE OF THE BINAURAL STETHOSCOPE.

BY H. W. SYERS, M.A., M.D. CANTAB.,

PHYSICIAN TO OUT-PATIENTS, GREAT NORTHERN CENTRAL HOSPITAL.

THE student of the present time is in danger of being overwhelmed by the multiplicity of the methods of diagnosing and treating disease. The continual outpouring of new remedies, most of which soon pass into merited oblivion, and the incessant recommendation of new (very often ancient) modes of treatment, chiefly distinguished by their grotesqueness, with all of which he is credited with possessing some sort of knowledge, render his career in the domain of therapeutics a chequered one. But with this side of the question I have now no concern, intending, as I do, to deal only with the method of diagnosis.

Twenty years ago the student was taught to rely almost entirely on the results obtained from the employment of his unaided senses, from the aspect of the patient, his attitude, and his mode of answering questions; in all these ways it was necessary for the learner to draw his own inferences, the use of special instruments being confined to that of the microscope, stethoscope, laryngoscope, and ophthalmoscope. Now, owing to the introduction of complicated blood examinations, of bacteriological methods of diagnosis, of colour reactions with the urine, and of many other proceedings of kindred nature, so much time has to be devoted to special modes of research that there is real danger of the old-fashioned but well-tried methods being relegated to the background in the student's scheme of practical work. To this craze for the employment of laboratory tests and for short cuts to diagnosis I attribute much of the present-day weakness in bedside examination which is, I regret to say, so often in evidence in union with the possession of the highest medical qualifications. These remarks are quite general and apply to no mode of examination in particular.

But, to pass from the abstract to the concrete, I would draw attention to a most urgently required reformation in a very important method of examination, an examination which is undertaken daily and hourly and which lies at the very foundation of clinical work. I refer to the examination of the chest and to the manner in which it is carried out. Until about 20 years ago the condition of the thoracic organs was always investigated by means of the wooden stethoscope, an instrument now almost banished and rarely seen either in hospitals or consulting-rooms. At the time referred to it became the fashion to employ the binaural instrument, at first in quite a tentative manner, then in combination with the use of the wooden instrument, and finally boldly and openly as the only stethoscope worthy the name. Now, were it the fact that the binaural instrument possessed advantages rendering it greatly superior to the wooden one this supersession of the latter would be merely an ordinary instance of the survival

of the fittest, and nothing would remain to be said on the use of stethoscopes. But the facts are altogether opposed to this view of the case, and I am convinced that much of the faulty diagnosis of chest disease which is so frequently seen at the present day is the result of the habitual use of the binaural stethoscope. And the reason why this important matter has not attracted attention hitherto is, I have no doubt, the undue weight which is attached to those laboratory methods of diagnosis to which I have already alluded, and which can only result in the production of an unsound, unpractical, theoretical (in the worst sense of the term) type of clinician. The errors which are the direct and immediate result of the constant use of the double stethoscope are of course most obvious in the case of those whose experience of the physical evidence of disease is limited, and is less apparent when the examiner by long practice has become familiar with the physiognomy of disease, so that he is to a great extent free from the necessity of employing an instrument in order to draw his conclusions as to the nature of the illness. But even in this case errors of omission, though not perhaps of commission, are not seldom made through the employment of the binaural stethoscope, while in the hands of the younger brethren faults of both kinds are, to my certain knowledge, only too frequently committed, due entirely to the obstinate use of this most pernicious instrument.

I will now in a few words give my personal experiences in this important matter of diagnosis by means of the ever-popular binaural stethoscope. I have convinced myself that as regards the investigation of the sounds of the heart this instrument is most unreliable, and increasing experience, not only of students' work, but of that of qualified practitioners who are in the habit of employing the double stethoscope, confirms me in the opinion that unless the use of this instrument be discontinued in no long time aortic regurgitation at all events will be a disease which cannot be certainly diagnosed. I do not, of course, refer to those cases in which the diastolic aortic murmur is so obvious that it cannot possibly be mistaken, but to that very large class of murmurs due to incompetent aortic valves in which the bruit is soft, low-pitched, and not by any means easy to hear. It is these murmurs which are constantly passed over, the second sound being described as "clear." The least reflection will convince anyone of the great danger attending the failure to discover such a murmur. In the case of a candidate for insurance much injury may be done to the office and one of which the directors would have every right to complain. It is well known that the soft, low-toned regurgitant murmurs are just those which are found in cases subsequently dying suddenly, the very softness of the murmur being a most unfavourable indication as depending upon a feeble ventricular systole. But it may be urged that in aortic disease the physical signs, apart altogether from the murmur, are so characteristic that it would be impossible for the lesion to be overlooked. I would reply to this objection that in the case of the murmur in question the left ventricle is often scarcely at all hypertrophied, there is no throbbing of the carotids, and the radial pulse may not possess the characters associated with the presence of aortic reflux. As regards the latter, indeed, the qualities of the pulse found in aortic regurgitation are by no means seldom met with in people whose aortic valves are perfectly healthy. It may also be mentioned that even if hypertrophy is present it is often masked by emphysema.

Thus, then, we are reduced to the evidence furnished by the murmur and this alone, in forming the diagnosis of a most serious and fatal form of cardiac valve lesion, and it is neither more nor less than a calamity for such a murmur to be overlooked. Yet to my certain knowledge this happens every day. Now, a murmur may be passed over from the incompetence of the examiner as well as from the unsuitableness of his instrument. In other words, it may be in some cases that a bad workman unfairly complains of his tools. But this argument, in my experience, does not apply to the question now under consideration. Most efficient and even brilliant auscultators I have found to overlook constantly this variety of murmur simply and solely because the binaural stethoscope which they to their subsequent sorrow, had been in the habit of using had misled them. The same observers when immediately afterwards employing a wooden stethoscope have invariably detected the murmur at once and have acknowledged their error, attributing it to its rightful cause—the use of an untrustworthy instrument. This is a certain proof that the slip was the