ART. IV.—ACTION OF LOBELINA.

BY ISAAC OTT, M. D.

SOME time ago I published some experiments on the action of this alkaloid of *Lobelia inflata* on lower animals. At that time I did not show that it was the active principle on which lobelia depends for its properties.

Baratlier,* in experimenting with ten to forty drops of a tincture of lobelia, obtained the following results: soon after taking it there was burning in the throat, eructation, nausea with widening of the pupil, disposition to sleep, difficulty in breathing, oppression over the chest, and in large doses tumultuous action of the heart, later sinking of the pulse, and after a few hours colic and fluid stools.

With the assistance of Dr. J. B. Heller, I made on myself the following experiment:

<i>Time.</i>	<i>Pulse.</i>	<i>Respiration.</i>	<i>Temperature by the mouth.</i>
2.23 P.M.	74	17	99
2.25 "	acetate of lobelina, 1-192 M. by the stomach.		
2.26 "	73	19	99 Tickling and dryness in the throat.
2.31 "	70	17	99
2.37 "	69	17	99½
2.41 "	68	18	99½; pupils di- lated; uncomfortable feeling in the stomach.
2.46 "	68	17	99½
2.54 "	72	18	99¼
2.56 "	1-192 M. of lobelina.		
3.04 "	74	19	99¼
3.05 "	lobelina. 1-192 M. retching ensues.		
3.08 "	76	18	99¼
3.10 "	lobelina. 1-192 M. vomiting follows.		

* Baudlin, Gifte und Gegengifte.

<i>Time.</i>	<i>Pulse.</i>	<i>Respiration.</i>	<i>Temperature by the month.</i>
3.13 P.M.	77	18	99 $\frac{1}{4}$
3.20 “	72	20	99 $\frac{1}{4}$
3.27 “	70	18	99
3.44 “	65	19	99 $\frac{1}{4}$
3.58 “	68	18	99 $\frac{1}{4}$
4.12 “	retching, vomiting, prostration, headache, and slight disposition to sleep.		

As in the lower animals, it at first increases the respiratory action and temperature. It also reduces the heart beat, which subsequently exceeds the normal, and then recedes it nine beats below the normal.

Like lobelia, it gives rise to nausea, emesis, prostration, dilatation of pupils, with disposition to sleep. Of course large doses would have a different effect.

This experiment demonstrates that the active principle of *Lobelia inflata* is lobelina, and that probably it is the only active principle.

ART. V.—MITRAL STENOSIS; CEREBRAL EMBOLISM; LEFT HEMIPLEGIA, CEREBRAL SOFTENING, DEATH.

BY M. A. WILSON, M. D., NEW YORK.

[Read October 27th, 1876, Before the Neurological Society, N. Y.]

MEDICAL literature is rapidly becoming replete with cases which may be included under the above heading, and the subjects of “Embolism” and “Thrombosis” are ones with which the profession are much more familiar at the present time, than even but a very few years ago.

It is within the recollection of the majority of physicians when all cases of sudden or unexpected death, were considered as due to “heart disease” or “apoplexy” alone; cerebral