A Mirror OF HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

WEST LONDON HOSPITAL.

A CASE OF "ANTITOXIN FEVER."

(Under the care of Dr. SEYMOUR TAYLOR.)

IN numerous instances after the injection of diphtheria antitoxin rashes have been observed, chiefly erythematous and urticarial in nature; pain and effusion in the joints have also been noticed in many cases, and in these points the interesting case described below is not unlike the results of an antitoxin injection, but at least a week seems to have elapsed between the injection and the first appearance of the symptoms and this fact cannot but throw some doubt on a causal connexion between the two. For the notes of this case we are indebted to Dr. Henry J. Davis, late house physician.

Whilst nursing a case of tracheotomy the patient, aged twenty-four years, contracted a mild faucial attack of diphtheria. She was admitted to the West London Hospital under the care of Dr. Seymour Taylor on Sept. 10th, having been previously indisposed for five days with sore throat and general malaise. The uvula was pendulous and cedematous, the tonsils were red and swollen, and their inner surfaces were covered with a white, glistening membrane which extended on to the left palatal arch; some adenitis was present. Cultivation showed the presence of diphtheria bacilli. 20 c.c. of antitoxin from the British Institute of Preventive Medicine were injected in the loin and twenty-four hours later 10 c.c. were injected in the buttock. Beyond some local pain following the lumbar subcutaneous injection no constitutional disturbance followed. The membrane rapidly disappeared, leaving some follicular distension which cleared on the third day. The temperature remained normal for seven days and the patient appeared to be making a rapid recovery. The adenitis had quite subsided by Sept. 15th. On the 18th there was some elevation of temperature coincidently with enlargement and tenderness of the infracondylar and cervical glands. The tonsils, especially the right, again became swollen and two days later a well-marked scarlatiniform rash appeared on the buttocks, thighs, and legs together with tenderness and effusion in all the principal joints. Constitutional disturbance was now considerable. There were headache, fever, drowsiness, nausea, and severe pains in the joints, back, and limbs. A papular rash appeared on the face and neck and with the fading of the erythematous rash a raised, blotchy, serpiginous eruption broke out on the trunk. The rash itched considerably. The effusion in the joints increased, the knees and ankles being exquisitely tender. The joint pains were relieved by salicylates, phenacetin, and bromides, the knees, elbows, and ankles being painted with equal parts of the liniments of aconite, belladonna, and chloroform and wrapped in cotton-wool. On the 23rd the rash had quite faded and the temperature was normal; all the symptoms rapidly disappeared, the patient made an uninterrupted recovery, and left the hospital for home on Oct. 16th. There was no albumin in the urine, peeling, cardiac complication, or paralysis throughout the illness.

Remarks by Dr. DAVIS.—The sudden onset with tonsillitis, adenitis, headache, rash, and joint pain suggested at one time the possibility of scarlet fever and acute rheumatism, but with the disappearance of the rash all the symptoms rapidly subsided. The antitoxin was obtained fresh from the Institute.

Medical Societies.

HUNTERIAN SOCIETY.

Exhibition of Cases and Specimens.

A CLINICAL meeting of this society was held at the London Institution on Dec. 8th, the President, Dr. G. E. HERMAN, being in the chair.

Dr. APPLEFORD showed a man, aged forty-nine years, who was the subject of a Curious Deformity of the Right Hand. The thumb and first finger were remarkable for their size, the latter being $5\frac{1}{4}$ in. in length and tuberculated. The second finger at the age of seven years had been removed on account of its bulk. The hand was a very useful one.

Dr. ARTHUR DAVIES showed a woman, aged twenty-six years, who had been treated seven years ago by Koch's Tuberculin, since which time the physical signs of phthisis had remained practically stationary. Owing to a contracting cavity at the left base the heart had been drawn into the post-axillary region.

Mr. POLAND exhibited a rare case of Acute Rachitic Curvature of the Tibia Backwards (just before the anterior tubercle) upon which he had performed osteotomy with success. The patient, a girl aged seventeen years, had fairly severe rickets from infancy, the ends of the long bones being still bulky and the femora much curved. The bend at the upper end of each tibia was at the lower level of the tubercle and presented an appearance not unlike dislocation of the knee. The rest of the bone in each limb appeared quite normal. At this point the operation was performed. The right tibia and fibula were carefully and gradually divided and their diaphyses forcibly bent backwards (bringing them into a straight line with their upper epiphyses) and the limb was placed on a double inclined plane. Firm consolidation took place readily in the corrected position. Radiographs of the left leg and knee were shown as illustrating the contrast between the two limbs.

Dr. KELSON showed a Rhinolith removed from a woman, aged sixty-two years, who had suffered from foul-smelling breath and some foetid discharges from the nose. The rhinolith was easily removed and all the symptoms ceased.

Mr. A. H. TUBBY showed a case of Syphilitic Curvature of both Tibiæ. The patient, a girl, aged twelve years, with a history of snuffles and a rash about two weeks after birth, was a stunted child of pallid and syphilitic aspect. The legs were remarkably curved forwards, which condition, existing since four years of age, had been gradually getting worse. The tibiæ were longer than the femora and were very tender, much hypertrophied, with rounded crests and uneven sides; their curve was entirely anterior, not lateral, thus being in distinct contrast to rachitic curvature. The condition was osteoporotic, similar to that found in Parrot's nodes. The teeth were not pegged or notched, but the palate was very highly arched.

Mr. COTMAN exhibited a boy, aged twelve years, who for about six months had had a persistent ring of erythema round his mouth caused by sucking his lips during and on waking from sleep. Remedies applied and watching him for a few nights effected a cure, but the trouble invariably returned as soon as these were discontinued.

Remarks on these cases were made by the PRESIDENT, Mr. TREGELLES FOX, and Mr. LAMPLOUGH.

BRITISH GYNÆCOLOGICAL SOCIETY.

Hæmorrhagic Endometritis.—Exhibition of Specimens.

A MEETING of this society was held on Dec. 9th, Dr. GRANVILLE BANTOCK, the Vice-President, being in the chair.

Dr. J. A. SHAW-MACKENZIE read an adjourned paper on Hæmorrhagic Endometritis. In pointing out the difficulties of differential diagnosis between chronic inflammatory conditions of the endometrium and commencing adenocarcinoma in curetted fragments he exhibited and contrasted microscopic sections of the same. He showed two uteri with microscopic sections removed by Dr. Bantock and