8. Loomis' System of Medicine.
9. Journal of Medical Research, new series, 1901, vol. I, p. 261.
10. British Medical Journal, 1902, p. 777; Saunders' Year-book, 1902, p. 412.
11. Cited by Cabot. Clinical Examination of the Blood, p. 190.
12. Genito-Urinary and Venereal Diseases, p. 435.
13. THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, September, 1898.
14. Lancet, December 22, 1900.
15. British Medical Journal, December 30, 1899.
16. Loc. cit., p. 543.
17. Journal of Pathology and Bacteriology, 1901, vol. vii, p. 214.
18. Deut. med. Woch., August 31, 1899.

ABSCESSSES OF THE LIVER FOLLOWING TYPHOID FEVER.

BY JOHN G. SHELDON, M.D.,
OF TELLURIDE, COL.

JOHN H., aged thirty-four years, came to me June 24, 1901, complaining of pain in the right side; chilly feelings and sweats, and great weakness. He said that three days before he saw me he had left his bed, where he had been confined for six weeks suffering from a disease diagnosed by his attending physician as typhoid fever. The history of this illness was typical for an ordinary case of typhoid fever, except that he felt that he was regaining strength very slowly, and that he had felt a heaviness and dull pain in the right side during the last week that he remained in bed.

The patient's previous history was negative. He suffered from measles when a child, and had smallpox when twenty years of age, but had had no other illness.

The examination showed a well-developed, poorly-nourished and emaciated middle-aged man. His skin seemed somewhat darker than normal, but he was not jaundiced. The mucous membrane of the lips and conjunctivæ was pale. There was marked tenderness in the region of the liver, and I inferred from the physical findings that the liver was diffusely enlarged. Its lower border was well defined and extended a hand's breadth below the costal arch at the outer border of the right rectus muscle. The upper border of relative hepatic dullness extended, in the right mammillary line, to the upper border of the third rib. Examination of the chest showed diminished expansion and respiratory excursion on the right side. The apex of the left lung seemed duller than normal. Harsh breathing was found in the apex of the left lung and over the lower portion of the right. A few moist râles were present in both lungs posteriorly. No friction rubs were heard. The heart was slightly enlarged transversely, and a systolic murmur was present over the pulmonic area. No thrills were found. The pulse was heating 108 times per minute; was not strong, but was regular. The temperature was 101° F. The spleen was not felt. There was no tenderness in the lower abdominal region; and no evidences of fluid in the peritoneal cavity could be found. The remainder of the examination was negative. The urine showed a trace of albumin, but was otherwise normal.

A diagnosis was not made at this time. A diagnosis of cardiac insufficiency, with passive congestion of the liver, was not warranted.

Tuberculosis and syphilis could not be positively excluded, but the symptoms led us to suspect the presence of an inflammatory condition in the liver.

The developments in the case were as follows:

June 25th. Chill in the evening, followed by sweating. Temperature, 102° F.; pulse, 120. Pain in right side severe. Blood examined for malarial organisms—none found.

26th. Morning temperature 99° F. Slight chill at 2 P.M. Temperature, 8 P.M., 100° F.; pain continuous; tenderness marked.

27th. Leucocyte count, 18,000; differential count not made. No tubercle bacilli found in the sputum.

The patient remained about the same during June 28th, 29th, and 30th. His symptoms were gradually increasing in severity, and his liver was becoming larger. Several punctures were made into the right hepatic lobe from behind, and on the evening of June 30th I introduced a needle between the eighth and ninth ribs about three inches from the spine, and at a depth of five inches, withdrew about a drachm of thick, yellow pus.

Owing to the marked enlargement of the liver anteriorly and to the absence of signs of pleural adhesions, I decided to attack the liver from in front, and on July 1st an incision, three inches long, was made parallel to the right costal margin, the centre of the incision corresponding to the outer border of the right rectus muscle. The liver was smooth, and many small yellow areas were seen on its surface. No adhesions between the peritoneum and liver were observed. The peritoneal cavity was protected with gauze, after which an aspirating needle was introduced into the liver and pus found at the depth of about one inch. This abscess was opened with a blunt artery forceps and contained about a drachm of pus. Another abscess was located to the right of the first and was drained through the same opening. The bleeding became free, but was easily controlled by a pack or by introducing the finger into the liver wound. By using the needle, uterine probe, artery forceps, and finger, I located and drained five abscesses, none of which contained more than two drachms of pus. I then introduced a rubber tube, wrapped the gauze into the wound, and packed the lumen of the tube to prevent hemorrhage. The anæsthetic which consisted of $\frac{1}{100}$ of a grain of hyoscine hydrobromate, and $\frac{1}{4}$ of a grain of morphine hypodermically, followed in one-half hour by ether, was taken nicely, and the operation, which was of short duration, was not attended with much shock.

The patient slept five or six hours after the operation, after which I removed the pack from the lumen of the tube. A considerable amount of pus and bile, with some blood, escaped. Eight hours after the operation the patient's temperature was 100° F., his pulse 106. He had very little pain—less than before the operation—and said that he felt relieved. For one week the drainage was profuse, and the liver diminished markedly in size. The temperature rarely came down to normal, but the pulse remained good, and very little pain was complained of. The tube was removed on the sixth day following the operation and a gauze drain substituted. On July 10th the patient had a chill; his pulse was 118; temperature, 102° F., and he complained of considerable pain in the right side. Examination showed no changes in the physical findings—the liver dulness extended, pos-

teriorly, as high as the eighth rib. A uterine probe was introduced deeply into the liver wound and a small amount of pus was discharged. This did not give the patient relief. The chills and fever recurred and the pain continued, being situated more posteriorly than usual.

On July 16th the liver fitness extended to the lower border of the scapula. The anterior margin of the liver was found to be about one finger's breadth below the costal arch. There was no respiratory excursion on the right side. Under local anesthesia a section of the ninth rib was removed in the posterior axillary line. The pleura was opened and not found adherent to the diaphragm. Twelve hours later pus was located at a depth of two inches from the base of the wound, and a large abscess cavity was opened by using a blunt artery forceps. A drainage tube was inserted. The patient felt relieved, the chills ceased, and the temperature and pulse became better. The patient at this time was very weak and somewhat emaciated. He got along very comfortably until August 1st, when he developed a cough. Pus still continued to discharge through the operation wound—the tube having been removed. From August 1st to August 10th the patient gradually failed. The pulse was weak and beat from 110 to 130 times per minute. The temperature was irregular, ranging from 100° to 101° F. No decided chills were complained of. The cough became more severe, and on August 8th blood was expectorated. On August 9th the patient complained of difficulty in breathing. On August 10th the dulness had extended higher on the right side than it had been previously. No movable dulness was made out. I introduced an artery forceps into the wound and opened the pleural cavity in an upward direction. The opening was dilated and a tube inserted. The improvement following this procedure was slight. On August 15th I explored the pleural cavity with the finger, broke up some adhesions and liberated a considerable amount of pus. The patient failed gradually. On August 20th he developed a diarrhoea, which lasted until August 22d, when he died.

The autopsy was done August 23d. The external examination of the body was negative, except for the presence of the two operation wounds. No icterus present.

The peritoneal cavity contained a small amount of clear, straw-colored fluid. The peritoneal surfaces were everywhere smooth, except that there were a few adhesions over the right lobe of the liver.

The pericardium contained a small quantity of clear fluid. Its surfaces were everywhere smooth. The heart was small, weighed 225 grammes. The myocardium was pale and friable. The cardiac valves were normal.

Many adhesions and a small quantity of pus were found in the right pleural cavity. The apices of both lungs were adherent. The lower portion of the left pleural cavity was normal. The pleura on the right side was markedly thickened.

The lower lobes of both lungs were hyperæmic, especially in the posterior parts. In the apex of the right lung several irregular cicatrices were found. A small cavity and a few scars were found in the apex of the left lung. The middle lobe of the right lung was firm and cut with resistance. Mucopurulent exudate could be squeezed from the bronchi. The bronchial glands were slightly pigmented. Calcareous deposits were found in two of them. There was found an opening in the diaphragm

—large enough to admit the index finger—which communicated with a cavity in the right hepatic lobe.

The liver weighed 1650 grammes. The right lobe presented a cicatrix and an irregular depression over its anterior surface. The posterior part of the right lobe was adherent to the diaphragm. On section the right lobe showed many irregular cicatrices in its anterior part. Situated in the posterior part of the right lobe of the liver was a globular cavity, 7 centimetres in diameter, communicating with the right pleural cavity through an opening in the diaphragm. This cavity contained dark-colored pus. It had no definite lining membrane. The liver tissue surrounding the abscess cavity was hyperæmic and friable.

The left hepatic lobe was enlarged in proportion to the size of the right. On its inferior surface five globular abscess cavities could be seen. These abscesses measured from 2 to 4 centimetres in diameter, and projected from the surface of the liver. On section the left lobe contained many abscesses, most of which were small. All of them had a definite lining membrane; and all of the smaller ones could be found to communicate directly with a branch of the portal vein. Pus could be squeezed from most of the portal vein radicals.

The biliary apparatus seemed normal; as did the hepatic artery. The splenic, portal, and superior mesenteric veins showed no gross abnormal lesions.

The lining of the small intestine was hyperæmic. No ulceration was found. Two Peyer's patches near the ileocaecal valve were elevated and indurated. The remainder of the patches presented a shaven-head appearance. The serosa of the small intestines was normal. The mucosa of the large bowel was hyperæmic. Otherwise, the large intestines were normal.

The appendix showed no abnormal changes. The mesenteric glands were enlarged and indurated. Softening was found in the interior of one gland, 2 centimetres in diameter.

The kidneys were slightly larger than normal. Their capsules were not adherent. On section they were granular and friable, but otherwise they presented nothing abnormal. The remainder of the examination revealed nothing of interest. The nervous system was not examined.

The succession of pathological changes leading to the patient's death were believed to be as follows:

Typhoid fever; carrying of infectious material from the intestines or mesenteric glands by the portal vein to the liver; suppurative pyelophlebitis, with abscess formation. The injury to the liver tissue done at the first operation resulted in the formation of a large abscess, differing from the smaller abscesses by being surrounded by a zone of hyperæmic liver tissue, and by being destitute of a definite lining membrane. This abscess might be classed as a secondary pyosepticemic abscess, as described by Andrew Davidson.¹ The empyema resulted from the second operation.

The diagnosis of typhoid fever was made from the clinical history and from the post-mortem findings in the intestines. A Widal reaction

was not tried. Slides made from the first operation and from the small abscesses found in the left hepatic lobe on post-mortem examination showed the presence of a small bacillus, which might have been either the typhoid or the colon bacillus. However, typhoid fever seems the most probable cause for the patient's primary sickness.

The belief that the hepatic abscesses resulted from diseased portal vein radicals was supported by the finding of a suppurative pyelophlebitis post-mortem and by the abscesses—especially the smaller ones—communicating directly with a branch of the portal vein. Infection of the liver by way of the arterial current was improbable. Although Hewlett⁷ states that typhoid bacilli are found in the blood in 83 per cent. of the cases of typhoid fever, and Courmoat⁸ believes that typhoid septicæmia is the rule, the absence of changes in the arteries and the absence of suppurating foci in other parts of the body speak strongly against arterial infection in this case.

Suppuration in the liver following typhoid fever is a rare condition. Holscher reports that 12 cases of liver abscesses occurred in the 2000 fatal cases of typhoid fever at Munich. Romberg reports 1 case of liver abscess from 88 fatal cases of typhoid. Scholtz reports 3686 cases of typhoid fever, 362 of these cases died, and in none of them was suppuration in the liver found. Da Costa⁹ has collected 22 cases of liver abscess following typhoid fever. Osler¹ makes the following statement concerning the occurrence of abscesses of the liver following typhoid fever: "Solitary abscess is exceedingly rare and occurred in but 2 cases in my series." Julius Dreschfeld⁵ says: "Acute yellow atrophy of the liver and abscess of the liver may be cited as very rare complications." I believe that it is impossible to say, with any degree of accuracy, in what proportion of fatal cases of typhoid fever liver abscess is found. During my experience as a medical student, as an interne in Cook County Hospital, and in private practice, I have knowledge of the post-mortem findings of about 200 cases of typhoid fever. The case herewith reported is the only one in which I believe that suppuration in the liver was found.

The relation of the typhoid bacillus to hepatic suppuration has not been determined. It is possible that the suppuration is due to the presence of the typhoid bacillus alone. I am unable to find the record of a case in which the typhoid bacillus alone was found in a liver abscess that followed typhoid fever. Lannois⁶ reports a case from which he isolated the typhoid bacillus, but the colon bacillus was also present. It is true that typhoid bacilli are found in pure cultures in the liver in cases of typhoid fever,¹ but in these cases no inflammatory reaction is present. As yet we have no proof that typhoid bacilli alone are capable of producing an abscess of the liver in patients suffering from typhoid fever. Bindo de Vecchi¹⁰ has shown that the liver is suscep-

tible to infection with certain organisms. The hepatic resistance may explain the inability of the typhoid bacillus to produce liver abscesses in cases of typhoid fever.

The colon bacillus is, in all probability, the organism that is most active in producing the so-called typhoid liver abscesses. In support of this theory are the following facts:

1. Colon bacilli are present in the abscesses of the liver following typhoid fever.

2. Roger⁹ has shown that the liver promotes the growth of the colon bacillus, while it inhibits the growth of the typhoid bacillus.

3. Macaigne and Kleck¹⁰ have shown that the virulence of the colon bacillus is increased by disorders of the intestinal tract.

4. Sanarelli¹¹ found that colon bacilli, isolated from typhoid stools, were more virulent than when isolated from healthy stools. He thought that the increase in virulence is due to the effect of the typhoid toxins, and devised an ingenious experiment which seems to prove this point. If it is true that the virulence of the colon bacillus is increased in a patient suffering from typhoid fever, and that the liver offers little if any resistance to invasion by the colon bacillus, it is not surprising that hepatic suppuration occasionally follows typhoid fever.

The organisms that are responsible for the production of hepatic abscesses following typhoid fever, reach the liver by way of the portal vein, the hepatic artery, or the bile-ducts. Infection through the portal vein seems to be the most common. In these cases it is probable that the organisms reach the portal vein from the intestinal walls or from the mesenteric glands. They are lodged in the portal radicals in the liver and there begin the process of abscess formation. In these cases the liver is usually diffusely enlarged and many abscesses are found. The abscesses have a well-defined wall—or lining membrane—and the smaller ones can be seen to directly communicate with the branches of the portal vein. The liver tissue surrounding the abscess, especially the smaller ones, does not show marked inflammatory reaction. Usually the findings in the liver are independent of changes in other parts of the body, and suppuration in other organs is uncommon in hepatic abscesses following pyelophlebitis. The cases reported by Lannois, Osler, Romberg, Gerhard, Trougel, Buckling, Asch, and Bernhard and the case herewith reported are examples of portal vein infection.

Abscesses of the liver—following typhoid—due to infection by way of the hepatic artery are more rare than the variety described in the foregoing. Cases, however, are reported by Louis, Chvostek, Sidlo, and Barth. In these cases the liver may or may not be enlarged. If enlargement does occur it is rarely uniform. The abscesses are usually single or few in number. They are apt to be of large size, with no

well-defined lining membrane and not connected with the portal vein radicals. Abscesses in other regions of the body is the rule and not the exception. There was an abscess in the parotid in Louis' case. Suppurative perichondritis of the larynx was found in Chvostek's case; and in Sidlos' case an abscess was found in the mastoid region. In these cases it is probable that the colon bacillus as well as the typhoid bacillus is floating in the blood stream. I make this statement because such cases are rare and typhoid septicæmia, as Hewlett and Courmont (*loc. cit.*) have shown, is common.

The third method of infection of the liver, with abscess formation, namely, by way of the bile tracts, is not common. Klebs alone reports a case of this character. Suppurative cholangitis occasionally follows typhoid fever, but I am unable to find reports of cases of well-defined liver abscesses following typhoid fever in which the organisms have been carried to the liver by the bile channels. The pus in these forms of abscess, described in the foregoing, is usually of a white or yellowish-white color. I can find records of hemorrhagic or dark-colored pus occurring only in cases where secondary pyosepticemic abscesses were present—that is, where abscesses were ruptured either by operation or by invading some cavity of the body. In these cases the abscess cavity has no "pyogenic membrane," the pus usually contains blood and is dark colored, and the liver tissue surrounding it shows marked inflammatory reaction. The abscess found in the posterior part of the right hepatic lobe in the case herewith reported was one of this character.

The symptoms of hepatic abscess following typhoid fever are: 1. The general symptoms of an infection. 2. Local manifestations pointing to involvement of the liver. Occasionally there are no symptoms complained of that would indicate hepatic involvement. Such cases are recorded by Theodore Dunin¹² and by Osler.¹³

The cases of liver abscess following typhoid that I can find records of occurred in adult males from twenty-five to thirty-seven years of age. One case occurred in a boy, aged twelve years, and another in a girl, aged eleven years. In none of the cases was there a history of previous disease involving the liver.

The first indication of involvement of the liver was manifested by pain in the hepatic region, accompanied by chills and fever. These symptoms occurred in all of the cases that I have reported, except the two mentioned in the foregoing, in which no symptoms referable to the liver were present. The pain and chills usually came on about twenty-five days after the onset of the illness. In one case¹⁴ they occurred as early as the thirteenth day; and in two cases—Lannois¹⁵ and the case herewith reported—they did not appear until the forty-second day.

Pain was present in all of the cases but two. It was usually of a

severe, sharp character, but in some cases was described as being dull. It was usually located in the region of the liver; was generally increased by inspiration, and in one case¹⁶ was aggravated by taking food. In no case was the pain intermittent or paroxysmal; and in no case did it cease until the pus was relieved from tension, either by spontaneous rupture,¹⁷ or by operation. Violent, prolonged, and repeated chills were present in most cases. Dunin¹⁸ mentions the absence of chills in his case. In no case did the chills present the regularity that might be expected in malaria.

Jaundice was by no means a constant symptom. Da Costa³ reports jaundice in 7 of the 22 cases that he has found recorded. It was present in the cases reported by Romberg,¹⁹ Kohts⁴ (Case I.), Sidlo,¹⁷ Petrina,¹⁴ Louis,²⁰ Chvostek,²¹ and Burder.²² In these cases jaundice occurred as early as the thirteenth day,¹⁴ and as late as the thirty-ninth day¹⁷ after the beginning of the illness. Jaundice occurred in cases with abscesses in other portions of the body,¹⁷ and in cases where suppuration was found only in the liver. It does not appear that jaundice is present more often in cases of pylephlebitis than in cases of infection by way of the hepatic artery.

In most cases the liver was enlarged. Usually the enlargement was diffuse, but in some cases the right lobe was affected alone. In Bauder's, Lannois', and Dunin's cases no enlargement of the liver was found clinically or at post-mortem examination. In Gerhard's case,²³ fluctuation was present over the right hepatic lobe. In all cases where the liver was enlarged, marked tenderness was present.

Ascites was present in one case of pylephlebitis,⁶ but no record of ascites is found in a similar case reported by Asch and Bernhard.¹⁴

No constant or characteristic condition was found in the spleen. Mention was made of its being enlarged in 3 cases. Thrombosis was found in the splenic veins in Lannois' case. I have not found record of splenic suppuration occurring in any case.

Suppuration in portions of the body other than the liver occurred in 7 of the cases. These may be divided into two classes: 1. Suppuration in the mesenteric glands. 2. Suppuration in other portions of the body. In the cases described by Romberg, Kohts, Louis, and the one herewith reported, suppuration was found in the mesenteric glands. Cases coming under Class II. may be divided into two groups: 1. Cases complicated by empyema. 2. Those associated with pus foci in distant regions of the body. I make this division because it is possible that a right-sided empyema might occur by direct extension from hepatic suppuration; while abscesses in distant portions of the body would suggest the carrying of the infection by the blood current.

Empyema occurred in the case described by Osler, and in the one herewith reported. Serous pleuritis was present in the cases reported

by Lannois and by Gerhard. In none of these cases did the hepatic suppuration indicate infection by the arterial supply.

Suppuration in distant portions of the body occurred in the cases described by Louis, suppuration in the parotid gland; by Chvostek, suppurative perichondritis; by Sidlo, pus in the mastoid, temple, and right axilla.

Very little can be found relating to urine, blood, and stool examinations in the reports of these cases. The findings in the case herewith described have been given.

Examination of the contents of the abscess cavities revealed pus of a light color in all cases except in those in which secondary pyosepticemic abscesses were found. In these the pus was of a dark color. In cases where bacteriological examinations were made the colon bacillus was found. Lannois found the typhoid bacillus also in his case. A micrococcus was found by Dunin, and a staphylococcus was found in Romberg's case.

The symptoms in cases of abscess of the liver following typhoid fever extend over a variable period of time. Gerhard's case died on the fifty-fifth day; Romberg's on the thirty-second day, and the one that I described on the one hundred and first day. Most of the cases died from exhaustion. Dunin's case terminated suddenly. Osler's succumbed to peritonitis. Occasionally cases recover. Sidlo's case passed pus in the stools on the eighty-fourth day and was well on the one hundred and twenty-eighth day. Kolt's case had, on the eightieth day, twenty bowel movements containing pus, and recovered on the one hundred and twentieth day. The cases that recovered were doubtless, from their histories, liver abscesses following pylephlebitis. French states that suppurative pylephlebitis may terminate in recovery. Treves²⁵ has operated on a case of suppurative pylephlebitis that recovered. I cannot find records of a case of hepatic abscess of arterial origin that got well. The prognosis in cases of hepatic suppuration following typhoid fever is always grave, and death may be expected in all cases associated with foci of suppuration in other portions of the body.

The diagnosis may be easy, difficult, or impossible. In Dunin's case the patient died suddenly without any symptoms indicating hepatic involvement. Pain in the hepatic region, chills, and enlargement and tenderness of the liver would suggest hepatic suppuration. Pylephlebitis might produce the same symptoms as hepatic abscess, and cannot be distinguished from it. Aspiration of the liver should be used as diagnostic aid.

The treatment of hepatic suppuration following typhoid fever is surgical. In all cases where a solitary abscess can be diagnosed, the pus should be evacuated by operation. Surgical interference is advised

against in cases of multiple abscesses of the liver. However, I believe that hepatic drainage would relieve the patient somewhat, and might be of value in some cases.

CASE I.—(Lannois, *Revue de Médecine*, 1895, p. 913). The patient was in bed with typhoid for five or six weeks. After the temperature had been normal for three days, he had chills and fever, and ten days later had pain in the liver and stomach. These symptoms continued. The urine contained albumin. He became emaciated and died without being jaundiced or without the liver becoming enlarged.

Autopsy. The liver weighed 1800 grammes. Twelve abscesses containing yellow pus were found in the hepatic substance. Peritonitis was present. Thrombi were found in the mesenteric and portal veins and in the spleen. Peyer's patches were found in the stages of ulceration, healing, and pigmentation. The appendix was adherent, and one ulcer was found in its lumen. The biliary apparatus was normal. Right serofibrinous pleuritis was present. Typhoid and colon bacilli were found in the hepatic abscesses.

CASE II.—(Osler, *Transactions of the Association of American Physicians*, 1897, vol. xii.). A man, aged thirty-seven years, suffered from septic symptoms. Three months previously he had had typhoid. Death resulted from peritonitis without symptoms of hepatic involvement.

Autopsy. Peritonitis. Appendix perforated. Pus in the mesentery. Empyema on the right side. The liver was large and contained many abscesses. Suppurative pyelophlebitis was present. The splenic vein was thrombosed.

CASE III.—(George S. Gerhard, *Medical News*, July, 1886). A man, aged twenty-six years, suffered from typhoid. On the twenty-first day the liver became large and tender and he suffered from diarrhoea. Chills developed and were accompanied by fever; later fluid appeared in the right pleural cavity. The patient declined rapidly. Fluctuation was found over the liver, and he died on the fifty-fifth day of his illness.

Autopsy. One large and many small abscesses were found in the liver. Fluid was present in the right pleural cavity. Peyer's patches were involved. The biliary apparatus seemed normal.

CASE IV.—(Prof. Kohts, *Berliner klinische Wochenschrift*). A boy, aged eleven years, had typhoid. His temperature was normal on the thirtieth day. Two days later he had a chill, which was followed by pain in the liver, repeated chills and fever, jaundice, and hepatic enlargement. Eighty-four days after the onset of the illness he had a sudden pain in the abdomen followed by the passage of pus through the bowels. The patient recovered after being sick one hundred and twenty days.

CASE V.—(Prof. Kohts, *loc. cit.*). A patient, aged twelve years, had the symptoms of typhoid. On the twenty-fifth day he had a chill which was followed by pain in the region of the liver. At the same time hepatic enlargement and tenderness over the entire abdomen became manifest. Death occurred on the thirty-first day of the illness.

Autopsy. The liver was enlarged and contained many abscesses. Thrombi were present in the portal vein. Suppuration was present in the mesentery. No abnormal changes were found in the biliary

apparatus; 100 grammes of fluid were found in the pericardial cavity. Peyer's patches were pigmented.

CASE VI.—(Romberg, *Berliner klinische Wochenschrift*, March 3, 1890). A man, aged thirty-seven years, suffered from an ordinary attack of typhoid. On or about the eighteenth day of his illness he had a chill, which was followed by sweating, fever, pain in the liver, jaundice, enlargement of the liver, and tenderness over the hepatic and abdominal regions. His spleen was palpable. These symptoms progressed rapidly, and he died on the thirty-fourth day of his illness.

Autopsy. The liver was enlarged and contained many abscesses. Thrombi were found in the portal vein and in the hepatic artery. Peyer's patches were involved. Pus was found in the mesentery of the cæcum. Pneumonia was present, involving the lower lobe of the right lung. A staphylococcus was found in the hepatic abscesses.

CASE VII.—(Louis, *Fievre Typhoide*, 1841, vol. i., obs. xvii.). The patient suffered from an ordinary attack of typhoid fever. During the decline of the fever he developed chills, fever, enlargement and tenderness of the liver, and jaundice. The patient died shortly after the appearance of these symptoms.

Autopsy. The liver was enlarged and contained one large abscess and six smaller ones. Peyer's patches were involved. Pus was found in the glands of the mesentery and in the parotid gland.

CASE VIII.—(Chvostek, *Med. Chirurg. Rundschau*, June, 1868, p. 1888). This was an ordinary case of typhoid fever that developed septic symptoms, with pain and tenderness in the liver, accompanied by cough and expectoration. Jaundice was present. The patient died from exhaustion.

Autopsy. The liver was found enlarged and contained two large abscesses. The intestines showed evidences of typhoid infection. An abscess was found in the lung. Pericarditis was present.

CASE IX.—(Burger, *Lancet*, October 17, 1874). This case differed clinically from the preceding one, in that no hepatic enlargement was present. The patient was jaundiced.

Autopsy. The liver was not enlarged, but many small abscesses were found in the right and left hepatic lobes. No other foci of suppuration were found. The portal vein was thrombosed. Peyer's patches showed evidences of recent involvement.

CASE X.—(Sidlo, *Der Militär Arzt*, Wien, 1875, No. 23, p. 20). This case of typhoid ran an ordinary course until the twenty-third day. Then he had chills, fever, and sweats, with pain and tenderness in the hepatic region, and with enlargement of the liver. He became jaundiced on the thirty-ninth day. Later, abscesses developed in the mastoid, temple, and right axilla. On the eighty-fourth day he developed a diarrhoea, and pus was passed through the bowels. The patient was well on the one hundred and twenty-eighth day.

CASE XI.—(Petrini, *Prag. med. Wochen.*, 1888, Nos. 41 and 43). This case of typhoid developed chills and fever, with pain and tenderness in the liver; jaundice and hepatic enlargement on the thirteenth day of his illness. He died from exhaustion.

Autopsy. The liver was large and contained many small abscesses. Thrombi were present in the portal vein. The intestines showed typhoid lesions. No foci of suppuration were found in the body outside of the liver.

CASE XII.—(Theodore Dunin, *Deutsche Arch. für klin. Med.*, 1886, Band xxxix., p. 379). A patient, aged thirty-five years, complained of typhoid symptoms and also of a thenar inflammation. The suppuration in the hand opened spontaneously on the fourteenth day of his illness. The typhoid symptoms continued. The patient had no chills; neither did he complain of symptoms that would suggest involvement of the liver. No jaundice or hepatic enlargement were present. On the eighteenth day he got out of bed (was not delirious) to go to the lavatory, and died suddenly.

Autopsy. The liver was not enlarged. It contained many small abscesses. Suppurative pyelophlebitis was present. The spleen was markedly enlarged, but no abscesses were found in it. Peyer's patches and the solitary follicles were infiltrated and necrotic. Oedema was present in the lower part of the right lung. The brain was anæmic. A micrococcus and an unidentified bacillus were found in the pus in the liver. No streptococci were present.

BIBLIOGRAPHY.

1. Andrew Davidson. *Allbutt's System of Medicine*, vol. iv. pp. 133, 134.
2. A. W. Hewlett. *Medical Record*, November 30, 1901.
3. J. M. Da Costa. *THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES*, July, 1898.
4. Osier. *Practice of Medicine*, 1901, 4th edition, p. 26.
5. Julius Dreschfeld. *Allbutt's System of Medicine*, vol. i.
6. Lannois. *Revue de Médecine*, 1895, p. 26.
7. Muir and Ritchie. *Manual of Bacteriology*, p. 333.
8. Blodo de Vecchi. *Lo Sperimentale*, An. III, F. 3.
9. Roger. *Sem. Médicale*, October 19, 1898.
10. Macaigne and Kleckl. *Arch. Gén. de Méd.*, December, 1896.
11. Sanarelli. *Ann. de l'Inst. Pasteur*, 1894, pp. 193 and 195.
12. Theodore Dunin. *Deutsche Arch. f. klin. Med.*, 1886, Bd. xxxix. p. 379.
13. Osier. *Transactions of the Association of American Physicians*, 1897, vol. xii. p. 380.
14. Petrina. *Frag. med. Wochens.*, 1881, Nos. 41-43.
15. Lannois. *Revue de Médecine*, 1895, p. 913.
16. *Ibid.*
17. Sidlo. *Der Militär Aerzt*, Wien, 1875, No. 23, p. 20.
18. Kohls. *Berliner klinische Wochenschrift*.
19. Romberg. *Ibid.*, 1890, 3-1.
20. Louis. *Fievre Typhoide*, 1841, vol. i., obs. 17.
21. Chrostek. *Med. Chirurg. Rundschau*, June, 1864, p. 168.
22. Burder. *Lancet*, October 17, 1874.
23. Gerhard. *Medical News*, July, 1886.
24. Asch and Bernhard. *Berliner klin. Wochens.*, 1882, vol. xiv. p. 772.
25. Treves. *British Medical Journal*, February, 1894.
26. Courmont. *Journ. de Physiologie et Pathologie generale*, 1902, vol. iv. p. 155.

A PRELIMINARY COMMUNICATION CONCERNING THE NATURE AND TREATMENT OF RECURRENT VOMITING IN CHILDREN.

BY DAVID L. EDSALL, M.D.,

INSTRUCTOR IN CLINICAL MEDICINE IN THE UNIVERSITY OF PENNSYLVANIA; ASSOCIATE OF THE WILLIAM PEPPER LABORATORY OF CLINICAL MEDICINE.

THE nature of the disorder that various authors have called recurrent, periodical, or cyclic vomiting has never been demonstrated, and the treatment has been correspondingly ineffectual. The commonest