Can an Animal Count?

SIR JOHN LUBBOCK, in his interesting paper on animal intelligence (*NATURE*, vol. xxxiii. pp. 46-7), virtually puts this question with reference to the dog. But the question whether a dog, or any other animal, can count will depend upon what we mean by counting. In the ordinary and correct signification of the term, counting consists in applying conventional signs to objects, events, &c., as when we say "one," "two," "three," to the striking of a clock. Clearly in this sense there is no reason to suppose that any animal can count. But there is another sense in which the term "counting" may be used—*i.e.* as designating the process of distinguishing, with respect to number, between the relative contents of two or more perceptions. While addressing an audience of 100 individuals a lecturer can immediately perceive that it does not contain 1000; and even without, in the true sense, counting them may make a tolerably close guess at their number. The accuracy of such a guess will depend upon two conditions. The first of these is the number of units to be computed, and the second is the previous practice he may have had in that kind of computation. Thus, every man is able to tell the difference between one and two, two and three, &c., up to perhaps seven and eight objects or events, without resorting to the expedient of marking off each with a separate sign. But somewhere about this point most persons require to adopt a system of numerical notation, if they desire to be accurate; and probably no one, without either special practice or some such system, could be perfectly sure whether he held eleven or twelve shillings in his hand, or whether a clock had just struck eleven or twelve. Indeed, it is just because of the rapidly-increasing difficulty of thus computing diminishing differences of ratio by immediate perception, that primitive man first lays the foundations of arithmetic by marking off the objects or events upon his fingers and toes. As already indicated, however, special practice makes a great difference in the accuracy with which such instantaneous computation can be made. Several years ago Prof. Preyer, of Jena, tried some experiments upon this subject, and found, if I remember correctly, that after a course of special training one might acquire the power of instantaneously distinguishing between twenty and twenty-one dots promiscuously scattered over a piece of paper.

Now, it is clearly only in this way that animals can be supposed to count at all; and, therefore, the only question is as to how far they are able to take immediate cognisance of the precise numerical content of a perception—or, in the case of a series of events, how far they are able to take similar cognisance of their past perceptions. But, as Sir John Lubbock observes, there is no record of any experiments having been made in this direction. Houzeau (tom. ii. p. 207) says that the mules used in the tramways at New Orleans are able to count five; for they have to make five journeys from one end of the tramway to the other before they are released, and they make four of these journeys without showing that they expect to be released, but bray at the end of the fifth. If this is really a case of "counting," in the incorrect sense of the term (and not due to observing some signs of their approaching release), it is probably due to their perception of a known amount of fatigue, a known duration of time, or some other such measure.

Several years ago my sister tried to teach an intelligent terrier to fetch a stated number of similar little woollen balls placed in a box at a distance from herself—the number stated, or ordered, being purposely varied from one to six. But although she is good at teaching animals, and here went to work judiciously in ways which I need not wait to describe, the result, as in the case given by Sir John Lubbock, was a total failure.

My object in making these remarks is to point out that in experiments of this kind the game seems scarcely worth the candle. Even if it were proved that a dog could "count" up to any particular number, all that we should have proved would be that the dog is able to distinguish between the degrees of two or more perceptions of a given kind; we could not thus prove any abstract conception of number on the part of the animal, such as is implied on the part of the "Damara floundering hopelessly in a calculation." Howsoever hopeless such floundering may be if the man is really *calculating*—*i.e.* employing some system of numerical signs—his operations are being conducted on a totally different psychological level from those of the bitch who, in surveying her litter of puppies, perceives that there is not so great a mass of them as she remembers to have perceived before. Psychologically considered, the artifice of numerical notation is as far above any such faculty of simple perception, as the artifice

of alphabetical writing is above that of simple association. I cannot doubt that a moment's thought would have shown Sir John Lubbock how needless was his precaution—while establishing certain associations of ideas in a dog's mind between written words on a card and the things signified—of spelling the words phonetically, "so as not to trouble him by our intricate spelling."

It is a most interesting fact that a dog's attention can be so far fixed upon written signs that a special association of ideas admits of being established between them and the things signified; but the psychological distance between establishing such a special association and spelling a word is so enormous as not to admit of computation. And similarly, even if my sister had succeeded in teaching her terrier to fetch a stated number of balls at word of command, no one could have supposed that she had thus taught the animal to count, in the sense of employing any system of numerical notation: she would only have proved the degree in which this animal was able to perceive, *without counting*, the different appearances presented by this, that, or the other volume of balls in a box.

GEORGE J. ROMANES

Lodge's "Mechanics"

PERMIT me to thank Prof. Tait for his kind and amusing criticism of my little book. I am struck with comic horror at the thought that anything in the preface can be construed into a comparison between works like Thomson and Tait, Clerk-Maxwell and W. K. Clifford, with such elementary picture-books as Deschanel and Ganot. I do not indeed share Prof. Tait's contempt for these "foreign" books; a student will find in them details, about (say) barometers or air-pumps, for which he may search the other works mentioned in vain. I did not urge students to read Thomson and Tait, because to those who can the advice is superfluous; to those who cannot it is disheartening. I did, and do, recommend such junior students as we get at provincial colleges to read easy works on Physics—not always because they contain a profound and satisfactory statement of principles, for how few of them do, but because they explain a multitude of details and experimental developments with which it was unwise to encumber a little book dealing mainly with vital principles, and aiming at being, in its humble way, an introduction to the classics of the science.

My book is primarily intended as milk for babes; and while it would be cruel to tell a baby to look at the sun, it is possible to direct his attention to a gas-light with some pleasure and satisfaction.

OLIVER LODGE

University College, Liverpool, November 13

The Resting Position of the Oyster—A Correction

IN a late number of *NATURE* (vol. xxxii. p. 597) Mr. J. T. Cunningham makes the extraordinary announcement that Woodward, Jeffrey, and Huxley were wrong in asserting that the oyster rests on the left side. I am in a position to state with positive certainty that it is invariably the left valve of the fry of the oyster which becomes affixed to a foreign object. I have examined thousands of very young adherent spat ranging in size from 1-90th of an inch to 2 inches in diameter, and have never found an exception to this rule. Besides the positive statements to the same effect made by Huxley and others, I would refer the reader to a brief paper by myself entitled "On the Mode of Fixation of the Fry of the Oyster" (*Bull. U.S. Fish Commission*, vol. ii., 1882, pp. 383-387); but I would caution the reader that Figs. 3 to 8 were reversed through an unfortunate oversight, as the apices of the umbos of all the larval shells figured on p. 387 should be directed to the left instead of to the right side. This blunder of the artist is pointed out in the explanation to plate 75, where the figures from the above-cited notice are reproduced in my paper entitled "A Sketch of the Life-History of the Oyster," forming Appendix II. to "A Review of the Fossil Ostreidae of North America," by Charles A. White, M.D., and Prof. Angelo Heilprin. In another paper of mine, "The Metamorphosis and Post-Larval Development of the Oyster," Rep. U.S. Fish Commissioner, Part 10, for 1882, p. 784, Fig. 2 shows the larval shell, L, of the young spat in normal position, with the umbo directed to the left. This figure may be compared with advan-

tage in respect to the points raised here with the figure of the external anatomy of the adult on plate 73 in my "Sketch of the Life-History of the Oyster," already cited.

Mr. Cunningham's inference that the left valve, usually regarded as the lower one, is really the upper, because he finds worm-tubes and hydroids most abundant on the convex or left valve, is founded upon an imperfect acquaintance with the habits of the oyster. For, if living oysters are thrown into the water they will invariably fall upon the bottom with the left valve downward. If dead oyster-shells—free valves—be similarly thrown into the water, they will invariably fall with the hollow side up, and the convex one down. And furthermore, both living and dead shells remain in just the position in which they fall. Dead shells sown as cultch, or collectors, fall in such a position, and most of the spat is "caught" on the exposed parts of the under surface of such shells, whereas little is found on the upper surface. The reason for this is that the sediment which is deposited on the upper surfaces, asphyxiates the young oyster-fry and the other larvæ which affix themselves before they can become established and strong enough to resist its effects. The affixed organisms on the exposed inclined under surfaces of the shells, are, on the other hand, protected from the accumulation of sediment.

It is also well known that the right valve of the oyster is always the most deeply pigmented, while the lower or left one is paler. This is always the case when oysters lie almost flat on the bottom. When crowded together on the natural banks in a vertical position, there is less difference between the colours of the valves. This difference is obviously due to some influence exerted by the position of the aspects of the body of the animal in respect to light, the same as in land and aquatic animals generally. I would conclude, for this last reason alone, that the right valve of the oyster is normally always uppermost, were it not for the fact that I have observed all the stages of transition from the spat to the adult condition in confirmation of such a conclusion. It is true that many young oysters have the right valve looking down when allowed to grow upon cultch or shells which have been sown upon the bottom to favour the collection of the spat, but that circumstance by no means subverts the conclusions of such cautious and careful observers as Brooks, Woodward, Jeffrey, Huxley, Horst, and others.

JOHN A. RYDER

Smithsonian Institution, Washington, U.S.A.,
November 11

The Rotation Period of Mars

ONE or two points in Prof. Bakhuisen's investigation of the rotation period of Mars require correction:—

First, my determination of the period, as finally corrected, was 24h. 37m. 22.72s., correct within 0.02s. The correction arose from the detection of a small clerical error. My final result was deduced from a comparison of observations by myself in 1873, with observations by Kaiser in 1864, Madler in 1830–1837, Sir W. Herschel in the last quarter of the eighteenth century, Huyghens and Hooke in 1666. I have since used the best observations during the oppositions of 1881 and 1883, without finding any occasion for changing my result even by one-hundredth part of a second, though I place no reliance on the second decimal figure.

But, secondly, Prof. Bakhuisen, like Mr. Denning some time since, has taken Kaiser's result uncorrected for the clerical errors—very seriously affecting it—which I detected in 1873. Kaiser counted three days too many in comparing Hooke's observation with his own: one day through a mistake in correcting for change of style, and two days (apparently) from counting the years 1700 and 1800 as leap-years. His thus taking three days too many of terrestrial time had the effect—since three corresponding Martian rotations were taken in—of introducing a deficiency amounting to three times the excess of a Martian over a terrestrial day, that is, $3 \times 37m. 22.7s.$, or 67.281 tenths of a second. This, divided by the total number of Martian rotations, about 88,900, gives as a correction about 0.0077s. to be added to both Kaiser's estimates, making them respectively 24h. 37m. 22.697s. and 24h. 37m. 22.668s., the mean of which, 24h. 37m. 22.6825s., is practically the same as the value I assigned, viz. 24h. 37m. 22.7s.

I think it probable that Schmidt (and perhaps Prof. Bakhuisen, too) followed Kaiser so far as the error of three days was

concerned. It would naturally be taken for granted that this part of Kaiser's work was free from error. If I had not been determined to find out where and how Kaiser's calculations differed from my own, I should not have found out his mistake, for certainly one would not expect to find two large errors in a work perfectly free from small ones. But so it was. I may remark that Prof. Newcomb, of Washington, went through my calculation, finding it correct, and that Kaiser really had made the mistake I indicated.

As this correction re-established what Kaiser had doubted, the accuracy of Hooke's observations, and of my own interpretation of them, Prof. Bakhuisen's correction is scarcely admissible. For a difference of 0.06s., multiplied by 88,900 for the Martian rotations between Hooke and Kaiser in 1873, gives a total discrepancy of an hour and a half, nearly all of which must be assigned to Hooke's observations and Huyghens' (which I brought into agreement with Hooke's by correcting Kaiser's one-day error for change of style).

RICHD. A. PROCTOR

5, Montague Street, Russell Square, W.C., November 23

Beloit College Observatory

MY attention has been called to a paragraph in NATURE (vol. xxxii. p. 514), which, quoting from *Science*, speaks of "the Astronomical Observatory of Beloit College as closed for lack of funds." It is not strange that you express surprise at this announcement. Permit me to say that it is positively untrue. So far as I can learn, the only authority for it is a strange and entirely unwarranted statement from our late Director, Mr. Tatlock's connection with our Observatory closed on July 1 last. Within one week of that date Mr. Charles A. Bacon was appointed his successor, and a few weeks later he appeared and took charge. He has already proved himself competent, both as an observer and as an instructor. New arrangements are made for both meteorological and astronomical observations, and special attention will be given to solar and spectroscopic work. Though not richly endowed, our "Smith Observatory" is well equipped, and under its present direction its facilities will be made helpful to both the advancement and the diffusion of science. A. L. CHAPIN
Beloit College, Wisconsin, October 31

CONFERENCE OF DELEGATES OF CORRESPONDING SOCIETIES OF THE BRITISH ASSOCIATION, HELD AT ABERDEEN

A NEW branch of the British Association glided unobtrusively into existence at Aberdeen, under the new rules passed in the previous year: I mean the Conference of Delegates of Corresponding Societies. The Committee to whom the general arrangements connected with this new branch is intrusted are now issuing a circular to the Corresponding Societies, signed by myself as its Chairman, and by Prof. Meldola as its Secretary, in which they give an account of the proceedings at Aberdeen, with comments thereon. Much of this will be of general interest, as it helps to explain the functions of the Conference, which, as the proceedings showed, were imperfectly understood by many of the delegates themselves. The report is too long to ask you to print it in full, and on the other hand its general purport is more easily seized by leaving out details. I therefore limit myself to sending you extracts from it, with connecting-links of explanation to make them read continuously.

The Corresponding Societies Committee of the British Association beg to lay the following statement of work done at Aberdeen, with comments thereon, before the Presidents and Councils of the various Societies whose applications for enrolment as Corresponding Societies of the British Association had been granted.

The Conference of Delegates was held on Thursday, September 10, and on Tuesday, September 15, both meetings having been called at 3.15 p.m., and lasting in each case about one hour.