a very severe one—granular kidney with uræmic asthma, cardiac dilatation, retinitis, and sanguineous effusion into the pleura and pericardium. The fact that paracentesis pericardii gave unmistakable relief in so desperate a case is the best testimony to the efficacy of this mode of treatment. All who saw the case—and they were many—were convinced that but for the tapping the patient could not have lived many days. The first tapping, performed in the fourth left intercostal space close to the sternum, though it gave temporary relief only yielded five ounces of fluid. Accordingly on the next occasion the puncture was made in the fifth space two inches from the sternum. Dieulafoy recommends that the trocar be introduced at no great distance from the sternum, so as to avoid opening the pleura, but there was thought to be little danger of this occurrence in the present case as there were good reasons for believing that the pleura and pericardium were adherent. The greater success of the second paracentesis is largely to be attributed to the semi-erect position of the patient at the operation facilitating the displacement forward of the exudation, which, as so often happens, had gravitated to the posterior part of the pericardial sac behind the heart. The hæmorrhagic character of the fluid withdrawn from the pericardium as well as from the pleura is noteworthy, as pointing to the severity of the renal affection. The quantity of fluid removed on the second occasion, twenty-eight ounces, is rather large as compared with the average of cases recorded, though a much larger quantity has been removed in individual cases. It is, however, well to remember that the removal of a much smaller amount—e.g., four or five ounces—has been productive of excellent results in several cases that have been reported. The good effects of the tapping were manifested by the circulation, respiration, and secretion of urine. The rapid relief of the dyspnoea shows that this symptom was largely due to the mechanical difficulties under which the heart was labouring. Not less remarkable is the rise in the amount of urine that occurred. Thus, in the twenty-four hours preceding the first operation the quantity was twenty-seven ounces; during the next twenty-four hours fifty-five ounces were passed. After the second operation, on Aug. 10th, ninety-seven ounces were measured, and ten days later the quantity had risen to 137 ounces. No digitalis, strophanthus, or other diuretics were given after Aug. 9th, so that the diuresis can only be attributed to the improvement in the circulation brought about by paracentesis. The oedema of the skin in the præcordial region began to diminish as soon as the pericardial pressure was relieved, but it was some days before it had completely disappeared. The diagnosis of the pericardial effusion presented no difficulty. Although the pulsus paradoxus was not present in its typical form it was noted on the day of the second tapping that the intermissions of the pulse mostly coincided with inspiration. This condition could no longer be observed after the fluid had been removed, and it did not return. The repeated hæmoptysis that occurred early in the history of the case is worthy of note in connexion with the hæmorrhagic character of the effusions into the serous cavities, and with the fact that the necropsy revealed no disease of the lungs with the exception of slight marginal emphysema. I have met with several cases of pulmonary hæmorrhage in middle-aged and elderly people who were free from any sign of cardiac or pulmonary disease beyond slight emphysema, but in whom the existence of granular kidney in a latent form was highly probable. Referring to these cases in the discussion that took place at the Medical Society of London a few years ago on the late Sir Andrew Clark's paper on the Non-tuberculous and Non-cardiac Hæmoptysis of Elderly Persons, I ventured to attribute the hæmorrhage to a rise of pressure in the pulmonary circulation secondary to the high tension in the aortic system. When, as is usually the case in such circumstances, some degree of emphysema and general arterial degeneration coexist we must recognise the presence of conditions favourable to the production of pulmonary hæmorrhage. I still regard this explanation as simpler and more probable than Sir Andrew Clark's assumption of a specific lesion of the pulmonary arterioles and capillaries dependent on "the arthritic diathesis." In a valuable paper by Dr. Samuel West on a case of Purulent Pericarditis treated by Paracentesis and by Free Incision, with Recovery,² statistics of paracentesis pericardii

are included, and many interesting points are brought out. Since this paper was published other cases have been recorded; but as Dr. West's conclusions still seem to comprise all the essential points I will now quote them in conclusion: (1) Paracentesis pericardii is not only justifiable, but an operation which may be safely undertaken with ordinary precautions; (2) the most suitable place for puncture is, in ordinary cases, in the fifth left intercostal space one inch from the edge of the sternum; but if the pleura be adherent the puncture may be made safely much further out, and even in the sixth space; (3) the instrument employed should be a trocar and cannula, with or without aspiration; and (4) the operation may be performed with advantage, not only in the pericardial effusions of rheumatic or primary origin, but also in those which occur in the later stages of general dropsy, if it should appear that the fluid in the pericardium is adding to the difficulties under which the heart is placed.

THE DIFFICULTIES OF PROGNOSIS IN INSANITY.¹

BY HENRY SUTHERLAND, M.D. OXON.

EVERY practitioner of medicine is aware how difficult it often is to give a prognosis in ordinary cases of bodily disease. There is no point in our professional career which is so likely to endanger the opinion the public may have formed of our skill in diagnosis and treatment. It can, therefore, be easily understood that in all cases of mental disease this difficulty becomes much augmented, for the reasons that the patient will not assist us by describing his subjective symptoms, and will frequently complain that he is suffering from various bodily complaints of which there are no signs whatever. In forming an opinion about any given case of bodily disease we are usually asked: "Will the patient recover?" "How long will he be ill?" "Will the patient die?" "If so, how long has he to live?" And, of course, such questions as these refer just as much to the somatic troubles which may accompany mental disease; but in the prognosis of insanity we have to answer in addition the following style of questions: "Will the patient recover *mental* health?" "If so, how soon?" "Will he die of the *mental* disease?" "If he recovers, will he have another attack?" "If he remains permanently insane, will his life be a long or a short one?" The classifications which have been made of mental diseases do not throw much light on the subject of prognosis. Greissinger divides them under two heads—curable with functional disorder of the brain and incurable with organic disease. Maudsley drops an excellent practical hint in his subdivisions into emotional and intellectual insanity, emotional insanity being a condition in which horrible crimes are committed, while at the same time the public are unable to detect any abnormality in the intellectual faculties. But to my mind the rule of thumb employed at St. Luke's Hospital is by far the best foundation upon which we can rely for a really useful division of the subject. At this hospital curable cases are alone admitted. The medical officer is instructed to ask the friends who are applying for the admission of a patient: "What is his age? What is the duration of the attack? Is it the first attack or not?" and, lastly, "What is the form of mental disorder from which the patient is suffering?" One of the most important points in regard to prognosis is undoubtedly the age of the patient. Thurnam tells us in his statistics that "the probability of recovery is greatest in the young, and undergoes a regular diminution as age advances." Most writers look upon an attack occurring after the age of fifty as hopeless. Bearing this point in mind I gave a guarded but unfavourable prognosis in the case of a Member of Parliament aged fifty years who consulted me some years ago. His grandmother had committed suicide and his father had died insane. The patient had what he called "une idée fixe," a sort of modified delusion, with other well-marked mental symptoms. I certainly believed his case to be incurable. I prescribed rest, and I attended to his liver, which was much enlarged. Three months later he had completely recovered, and was able to make a long speech in the House of Commons. I must confess I was completely

² Transactions of the Royal Medical and Chirurgical Society, vol. xlvii, 1223.

¹ A paper read before the West London Medico-Chirurgical Society.

deceived on that occasion, as from the history of the case I had every reason to believe it to be incurable. As a contrast I may allude to the following case. A youth aged eighteen years was brought to me some time ago who was said to be a victim of self abuse. The late Dr. William Wood, who had a more profound knowledge of insanity than anyone I ever met with, pronounced the case to be hopeless. I did not agree with him; but I was wrong. In the course of time the patient so far recovered as to be able to start on a sea voyage. But the first night at sea he told his attendant that a horse had put his head in at the cabin window. After this he became violent, was placed in irons, sent home, and placed in an asylum. He is now under my care, having been insane for more than twenty years without any lucid interval. My opinion that his youth would save him from being a chronic lunatic turned out to be incorrect. The number of attacks a patient has had seriously affects the prognosis. A woman at the change of life consulted me. She had become depressed in consequence of having been left a large fortune unexpectedly, became suicidal, was sent to an asylum, and recovered. I prophesied that she would be a victim to recurrent melancholia. She went out and tried to keep house, but had not the strength to do so. She had a second attack and recovered. The patient went out again and got into the hands of two drunken women who swindled her. She had a third attack and returned to the asylum. She again left, and was taken in hand by another woman, and has had several attacks since. During one of them her brother and I endeavoured to persuade her to become a Chancery patient. She refused to take our advice, and she is now gradually drifting from a state of melancholia into one of hopeless dementia. On this occasion I was right. But in a case later I was not so fortunate. A woman had six times suffered from puerperal insanity and had been received into an asylum. When her seventh child was being thought of I advised that she should place herself under medical supervision as a boarder, and thus avoid the fatigues of the London season at this critical period. She passed through her seventh confinement without an attack. She had an attack on the birth of an eighth child. After that she was absent from her husband for two years, but after one year's absence from him a ninth child was born; a divorce was the consequence. Since then she has had children, and has remained perfectly sane on each occasion. My idea that she would end in chronic dementia has thus far proved incorrect.

The form of insanity is also of the greatest importance in giving an opinion. In Dr. Hack Tuke's Dictionary of Psychological Medicine there is an excellent article by Dr. Blandford giving the probability of recovery in every form of insanity. It can only be remarked here that acute cases as a rule are curable and that chronic cases are incurable. According to a very clever classification of mental diseases, drawn up by a committee of the Medico-Psychological Association, the following forms are incurable: general paralysis, epileptic insanity, senile insanity, paralytic insanity, and perhaps idiocy. In Dr. Mickle's work on General Paralysis of the Insane he mentions the names of not less than thirty-six writers who have published cases of recovery from general paralysis. In my own opinion a recovery from general paralysis, when the symptoms are well marked, never has and never will take place. In such so called recoveries either a wrong diagnosis has been made or the patient has left the asylum during a temporary convalescence and has been lost sight of. In the advanced stage the diagnosis of this disease is generally very easy. At least, so I and some others thought about a man whose case at the time attracted much attention in the psychological world. This patient's left pupil was always much contracted and insensible to the stimulus of light. The right pupil was much dilated. He had marked hesitation in his speech, more especially in his pronunciation of linguals and labials, which, as everybody knows, is very characteristic. I remember his delusions of grandeur very well. He said he had a horse which was going to win the Derby, that he was about to row for and win the Diamond Sculls, and that he had bought a four-in-hand. The case was carefully watched, epileptic attacks and other signs of advancing disease being daily anticipated. At the end of three months the patient made a good recovery, was discharged from the asylum, and I have reason to know never had a second attack. It was found out afterwards that his hesitation in speech was congenital, that the contraction of the left pupil was due to an old iritis, and that the

delusions of grandeur were only those which may be met with any day in certain cases of acute mania.

Any form of insanity associated with epilepsy is generally considered incurable. The exceptions to this rule are infantile convulsions, epileptic insanity at puberty in girls, and epilepsy after a prolonged and excessive use of alcohol. I was once sent down to Wales to rescue a lunatic who was being robbed of his property. He had been staying for some months in a low village inn, where, in order to conciliate him, he was supplied with unlimited beer. I found him reeling about with a glass of beer in his hand, which he sipped all day. I brought him up with great difficulty to London. On his arrival he had three distinct epileptic fits. We all pronounced the patient to be suffering from incurable epileptic insanity. In this we were wrong, as the patient for ten years never had another epileptic attack, the three on admission having been entirely due to drink. About the same time a woman aged twenty-eight years was sent by Dr. Buzzard suffering from epileptic mania. She commenced having these fits at about twenty-five years of age, and they were increasing in frequency every week. The diagnosis was clear and the outlook hopeless. After being in an asylum for some years she died from exhaustion, complicated with bronchitis. On this occasion we were all correct in our opinion.

A word as to idiocy. The prognosis in this form is most unfavourable, but more hopeful than might be supposed. Dr. Shuttleworth tells us that of patients discharged from idiot asylums after full training, 10 per cent. are self-supporting and more than 40 per cent. are capable of performing the ordinary transactions of life. Some years ago I was locum tenens at St. Luke's Hospital. As already stated, no incurable patients are admitted to this institution, and all patients are seen by the committee on their arrival. A male patient was brought into the board-room whose relatives had told us he was suffering from acute dementia. Two other medical men and myself were about to give directions as to the ward in which the case was to be placed, when a layman who had sat for many years on the committee exclaimed, "That is a congenital imbecile." I was sent out of the room to make further inquiries, and I found the layman was right, and that the three medical men, including myself, were wrong in their diagnosis. The patient was not admitted. An infant was brought to me by its parents about sixteen years ago. Forceps had been used at its birth, and the frontal bone had been compressed so that its form resembled the keel of a boat more than anything else. My prognosis was unfavourable. The girl is now eighteen years of age, can only articulate as a child of five years might do, and delights to play with dolls and to read childish books. She is quite incapable of doing any arithmetic or managing her own estate. It is impossible to describe cases illustrating points in prognosis in every variety of insanity, but we may take it for granted that in any patient in whom there is a regular periodicity in the attacks all hopes of cure are at an end. I have a woman under my care at the present time who suffers alternately from mania and dementia, each form of mental disorder lasting as nearly as possible nine weeks. I have also a man similarly affected. In most cases of *folie circulaire* which have come under my notice I have found dementia taking the place of mania more frequently than melancholia. Homicidal insanity is in my experience incurable. I was called upon some years ago to examine a man who had murdered his wife and two boys and also attempted to murder his two other children, but they refused to drink the poison he had prepared for them. The murderer then placed his own head on the line of a railway when an express train was expected, intending to commit suicide. He was rescued from this position by a pointsman. He was very much attached to his wife and the children he had killed, he had a good balance at the bank, and was in a good situation. He gave himself up and confessed all to the police. I had not the slightest doubt that the prisoner was suffering from the form of insanity known as "impulsive," which has been so ably described by Dr. Maudsley, and that he was an incurable lunatic. The judge, however, took a different view, informed the court that he would have no more murderers let loose on the plea of mental derangement, and was pleased to say that I had thrown dust in the eyes of the jury. After retiring for a few minutes the jury returned to the court, their verdict being "Not guilty, on the grounds of insanity." I visited Broadmoor Asylum some years later, where the prisoner had been sent, and found him to be one of the most maniacal homicidal patients in the institution, and also perfectly

incurable. The duration of the attack has naturally a great influence upon the chances of recovery. At the Retreat, York, 80 per cent. recovered who had been treated within the first three months of their attack. During the first year about 60 per cent. recovered. After the first year about 10 per cent. recovered. Dr. Thurnam says: "It may be stated fairly that six times as many cases recover within the first year as recover after the first year of the illness. After two years recovery is rare." Nevertheless, recoveries have been known to take place after six, ten, and even twenty years' duration. A woman was admitted to an asylum suffering from suicidal melancholia. Before she was placed under treatment she had gouged out her right eye with her thumb, in obedience to the command in the Bible, which says, "If thine eye offend thee pluck it out and cast it from thee." After some time the melancholic symptoms disappeared, and she settled down into what we considered to be a state of chronic incurable dementia. At the end of seven years she made a sudden recovery and was discharged perfectly able to manage her own affairs. In a long experience I have never known a case recover after so long a duration as seven years. I have had experience of patients, both male and female, who are still under my care after thirty, forty, and in one case fifty years. It is needless to remark that the prognosis is perfectly easy in such cases. With regard to sex it may be observed that a larger proportion of women than men recover. The numbers are 44 per cent. for females and 36 per cent. for males. One reason for this excess of recoveries on the women's side is that they are less afflicted with general paralysis than are the males.

I will now turn to somatic influences with regard to this subject. In my experience the prognosis in insanity from sunstroke is favourable in mild cases, but most unfavourable in severe cases. Sir Joseph Fayrer tells us that on post-mortem examination after sunstroke the brain and its membranes are found congested. My own idea is that when a person is attacked either slightly or seriously by *coup de soleil* the sympathetic circles of nerve fibres surrounding the arterioles become paralysed permanently, and in consequence the bloodvessels do not contract naturally. Hence a larger amount of blood is conveyed to the brain under even a slight stimulus, and excitement and mania result. I was once at a private ball at a seaside town. A young man entered the room, much flushed and talking loudly. He proceeded to pull the cornet out of the mouth of the player, lifted the hands of the pianist from his instrument, and pushed the violin from under the chin of the fiddler. He was expelled from the house, everyone saying he was drunk. He recovered next day, called on the hostess, and made his apologies in due form. This young man was an officer in the army and had had a slight sunstroke. He was a teetotaler, but feeling tired before the ball he took a glass of wine to revive himself, which went to his head and caused a transitory attack of mania. I had no reason to disbelieve his statement that one glass of wine produced these awkward consequences. But insanity from a severe sunstroke is a very formidable and incurable form of mental disorder. I had for many years an Indian general under my care who was one of the most dangerous patients it has been my lot to deal with. His condition was entirely due to a severe sunstroke he had sustained during an unusually hot summer in India. In such cases I have observed that there is a period of incubation. The symptoms most characteristic of the disease do not appear till some months after the exposure to the solar rays, although perhaps in the meantime a slight intellectual and emotional departure from the healthy standard may be observed. The catastrophe I am about to relate took place during the Indian Mutiny some forty years ago. The general had been told off to guard a certain number of Commissariat waggons. His story, which he presented to me most modestly, has neither a clinical nor a pathological interest, but well illustrates how the mind may become unhinged even some months after a sunstroke has visited the patient. He said: "I was in command of a party of forty horsemen. We came suddenly upon a force of 400. It was a case of ten to one. It was my duty to retire, but I felt that murder was in the air. I gave the order to my men to charge. My trumpeter was shot down on the right of me and my orderly on the left of me. I cut my way in and I cut my way out. All perished except myself, and I thought I was well out of it." In these few words were expressed what appeared to me to be a most romantic episode of military life. Unfortunately there is a seamy side to the story. The general was tried by court-martial, and his behaviour was so eccentric on this occasion

that a medical examination as to his state of mind was ordered. He was found insane and sent from India to an asylum, where he remained for twenty years and finally died an incurable lunatic.

I now pass to the prognosis in cases of alcoholic insanity. There is no class of cases which recover so rapidly as do those of acute alcoholic insanity due to one inordinate drinking bout, and there is no class of cases more incurable than that of patients who during a long course of years have soaked themselves day by day in stimulants. A captain in the Royal Artillery was admitted suddenly one afternoon to an asylum. He had indulged in an unusual amount of alcohol and had become mischievous and almost homicidal. He had taken his wife in his arms and had held her over the balcony of his house and threatened to dash her on the spikes of the area railings below. The police were called in, and he was certified and sent to a harbour of refuge. On admission he was in a state of violent mania and passed the night in the padded room. I visited him again next morning, and after a three hours' examination could find no indications whatever of mental aberration. He had completely recovered in one day, the recovery being simply due to abstinence from stimulants. By his own request he was detained as a boarder for a week, as he said, to keep him from the drink. On his discharge he took more freely than ever to alcohol, and died six weeks later from delirium tremens. The prognosis was, however, very different as from even a temporary recovery in the case of a man aged seventy-six years whom I was sent for to attend one Sunday morning. He had for not less than forty years been in the habit of getting drunk frequently, although able to go down to his office in the City and manage his affairs fairly well. At last he became so unrepresentable that his sons were obliged to protest against his taking any further part in the business. The unfortunate part of the case was that the man would neither die nor become insane. In France a *conseil de famille* would have been held to decide what was to be done with this worshipper of Bacchus. Unfortunately no such law obtains in England. I only saw him once, but I heard afterwards that his and his sons' business was going from bad to worse owing to his unfortunate propensities. In this case it was suggested to my mind that the law of China as to the treatment of lunatics might have been justifiably employed. In that enlightened country they have but one method. All lunatics there are treated by decapitation. I have before alluded to Dr. Maudsley's subdivision of mental diseases under the two heads emotional and intellectual insanity. Emotional insanity is a very important form to recognise. It is very rare. It leads to horrible crimes being committed by its victims. The intellect being nearly but not quite intact, it is impossible to persuade the relatives that there is anything wrong with the mind of the patient. The prognosis is usually most unfavourable in this form. I have a woman at present under my care residing in the house of a medical man. She has been insane for about five-and-twenty years. On one occasion a high authority on lunacy visited her and pronounced her to be perfectly sane. I asked him how long he had talked to her. He said ten minutes. I told him to go back and give her ten minutes more. In the course of the second ten minutes all the delusions and hallucinations came out, and he was obliged to confess that she was an incurable lunatic. This patient knows a great deal, but not all, about her investments and dividends, but ask her to describe a carriage accident or any place of amusement she has visited and she breaks down before the end of even a short narration. She indulges in the most filthy conversation, rendering it impossible for her to take her meals with the family. The case is incurable. I had another very curious case under my care a few months ago. The patient was a clerk in a Government office, engaged in the most complicated transactions during many hours of the day. Excess of alcohol was the root of the evil in his case. If he were in the public streets or an omnibus he fancied that people put up their hands to their mouths, stamped their feet, or made grimaces on purpose to annoy him. He had a brother and a wife, one of whom always accompanied him to and from the office. It was useless to tell him that they, who had met the same people in the street that he had, had never seen them do anything which might be construed into an insult. He became worse, and I obtained leave of absence for him for six months. My prognosis was that as long as he kept from drink he would be perfectly sane. He became a teetotaler and recovered perfectly. The extraordinary part of the case was that this man, affected with delusions of suspicion, was able

to carry on severe mental work all the time he was apparently insane. As to delusions generally, it may be said that if the patient has a variety of them, quickly changing and accompanied with noisy violence, the outlook is hopeful. Fixed delusions, existing for more than a year and continuing when the acute symptoms subside, mark the case as incurable. From among such patients we obtain our asylum kings and queens, who are content to work in the scullery or laundry during the day and assume Court dress or a regal crown when festivities take place in the evening; but of all delusions those that are progressive are the worst as regards prognosis. The general paralytic will commence by telling you he is going into Parliament, that he has been knighted, made a baronet, or created a peer, that he is the Prince of Wales, and finally that he is God Almighty Himself. This is an important point to be remembered in diagnosing a case of general paralysis from one of chronic mania. A great deal of light is thrown on prognosis by observing the postures and movements of the insane. I have a young woman at present under my care who sits all day with her arm over the back of a chair and her head leaning on it. This position has remained fixed during the day for three years. The outlook is hopeless. Another woman, young on admission, rubs her nose continually, making it very red. This she has done for many years. A man under my care walks round two flower-beds in the garden, his course taking the shape of the figure 8. He kicks anyone who gets in his way, but is otherwise harmless. One elderly man has for twenty years rolled a pocket handkerchief up in a ball and turned it round and round in his left hand. With his right he continually percusses his knee and thus wears out the cloth of his trousers in less than a week, necessitating endless patches. *Hæmatoma auris* almost always is found amongst chronic incurable cases. I have a woman with a marked swollen ear at present under my charge who will never recover. I have, however, seen *hæmatoma* in sane people, generally prize-fighters. I exhibited a professional boxer at the Clinical Society of London some years ago who had this peculiarity. Certain bodily complications have great influence on the outlook. For instance, the recoveries in puerperal insanity are very numerous, but if the case should be accompanied by albuminuria the patient almost invariably dies. If the patient should increase in weight, the mental condition remaining unaltered, the outlook is very unfavourable. A woman admitted in 1891 weighed 8st. 13lb., and on the third anniversary of her admission—that is, in 1894—she had increased in weight to 12st. 21lb. This was an increase of 3st., or 1st. a year. Her delusions are always the same—that she is suffering from various bodily complaints of which there are no tangible proofs. Syphilitic insanity in the early stage is fairly curable, but later ends in the death of the patient. I had an Indian colonel under my care for many years suffering from syphilitic mania. Not being satisfied with my treatment a very eminent physician was called into consultation. A third medical man, a friend of the patient, was present. The physician diagnosed the case as one of syphilophobia—that is, a delusion on the part of the patient that he has contracted syphilis. I made no remark and the physician departed. When he had left the room I turned up the patient's trousers and showed his friend, the third medical man, a large open brown syphilitic sore which the patient had had on his shin for many years. The patient died a month later. The condition of the blood in the insane is a great guide in giving an opinion as to how long a patient has to live. When I was at the Wakefield Asylum I examined under the microscope the blood of some hundreds of lunatics. The results were very satisfactory. It would take too long to enumerate them here, but I cannot resist mentioning one point which was to me very striking. I found that if, in general paralysis of the insane, a much larger proportion of white corpuscles as compared with the red was present, the patient was sure to die within four days, although he might apparently be in fairly strong health. This and my other observations have been confirmed by Dr. Lauder Lindsay and Dr. Rutherford.

Before concluding, I should like to say a word or two about the diagnosis of recovery in a patient. Recoveries are always calculated on the admissions. According to Dr. Thurnam they may vary from about 25 to 50 per cent.; but, as I have said, at St. Luke's Hospital, where only curable patients are admitted, the number of recoveries reaches 60 per cent. Generally speaking a rapid recovery is likely to be succeeded by a second attack. A slow recovery is much more hopeful.

Convalescence from insanity is indicated by quietness of mind in the patient: he speaks freely about his disease, he does not perpetually bother the physician as to when he is to leave the asylum, he asks to see his friends, and he returns to his usual employments. An outburst of weeping in females during an attack of acute mania is a good sign, as showing that the patient is aware of her condition. On the other hand, a contented state of mind points to increasing dementia. Esquirol remarks that many patients who are considered cured by their relatives are not so really. In this I quite agree. In proof of this I may mention that not less than three patients removed from our care by the wish of friends and contrary to advice committed suicide, one on the very night of his discharge. Patients are frequently sent back to me to be examined after they have left in order to ascertain whether or not they are capable of managing themselves and their affairs. A list of their former delusions is prepared from the case-book, and the patients are taxed with them one by one. The conclusions are drawn from their answers. Valuable information can also be got from the attendant who usually accompanies them as to their actions and habits being sane or insane since they left the asylum. In almost all cases three months after discharge there are still present traces of the old disorder. Frequently the patient looks lost and distressed, the angles of the mouth are drawn down, there is a general feebleness about the outlines of the features, and the bodily health is never quite re-established. They complain of great irritability and they are easily moved to tears. Their letters also eloquently express their undecided and vacillating state of mind. In females there is a weak smile on the face and frequent showing of the gums, the eyebrows are elevated as they speak, and there is a marked transverse corrugation of the forehead expressive of doubt and anxiety. But it is only by long experience and a careful examination of a case that we are able to determine from these rough rules what is to be done with the patient. The importance of giving a correct prognosis in cases of mental disease cannot be over-estimated. Frequently large pecuniary interests depend upon our opinion—as, for instance, when a large sum is offered to the patient if his life is only to be a short one, as in general paralysis, or if he is to receive a pension if his life is to be a long one, as in chronic melancholia. The matrimonial prospects of both sexes are also much affected by the form of insanity and other points which have to be gone into when an alliance of this kind is contemplated. Cases of hysterical mania in young girls are very hopeful, as well as to recovery as to the *non-probability* of a second attack. Any girl who has had two attacks of insanity ought not to marry in my opinion. The parents of both parties will come to consult us on these difficult and delicate matters. One rule I always lay down in such cases, and that is that the sane contracting party should be told the whole truth about any previous attack from which the other may have suffered, and then if he or she choose to proceed with the engagement they do so with their eyes open and take the responsibility on themselves. The wife of a medical man during her engagement told her future husband that she had some dreadful secret to reveal to him. He thought she referred to some previous engagement. She said to me afterwards, "He little thought I had two asylums at my back," meaning she had been twice certified as a lunatic. Six weeks after marriage she became insane and remained so for many years, the unfortunate husband having all this time to maintain a wife who was worse than dead to him. Finally, our method of arriving at a correct prognosis may be compared to weighing in a pair of scales those points which, on the one hand, are in favour of, and, on the other, are against the chances of recovery in any given case, such points, however, not being estimated by their number, but by their quality. Experience alone can teach us the value of each individual point. But when the importance of the subject is considered it is surely our duty to neglect no opportunity of improving our natural powers of observation, and thus to be enabled by the accuracy of our opinion to confer health and happiness upon our less fortunate brothers and sisters.

Richmond-terrace, Whitehall, S.W.

MEDICAL MAGISTRATE.—Mr. Thos. E. Jacobson, L.R.C.P. Edin., M.R.C.S., has been placed on the Commission of the Peace by virtue of the office of chairman of the Sleaford urban district council.