8th.—Ordered to get up for two hours.

From this date he gradually gained strength till he was discharged on the 1st of April perfectly recovered; and, notwithstanding the mutilated condition of the limb, and the consequent difficulty of procuring suitable flaps, there is a well-covered stump (*vide* lithograph, Plate VII.)

Considering the severity of disarticulating the shoulder-joint, the results are tolerably successful. In the Crimea the mortality was thirty-five per cent, but according to Dr. Gross, out of sixty-one cases occurring in American and European hospitals, there was a mortality of nearly fifty per cent. Whereas Ashhurst, quoting from Circular, No. 6, states the mortality to be thirty-nine and a-quarter out of two hundred and thirty-seven cases, and that when performed for other than traumatic causes, it is still more successful. In operating at the shoulder-joint for injury, the surgeon must select that method by which he can obtain the best flaps, and not be partial to any particular one. For this reason I selected the operation above described, in preference to Larrey's, or transfixion.

ART. XII.—*Cases of Orbital Disease.* By H. MACNAUGHTON JONES, M.D., M.Ch., Fellow of the Royal Colleges of Surgeons, Ireland and Edinburgh; Surgeon, Cork Ophthalmic and Aural Hospital; Senior Demonstrator of Anatomy, Queen's College, Cork, &c., &c.

THERE can be no doubt of the intractable nature of many cases of accumulation of pus in the frontal sinus. Mr. George Lawson has drawn special attention to this affection, and, with his usual skill, has proved how successfully it may be treated by a drainage-tube passed from the sinus into the nose and out through the nostril (*vide Practitioner*, July, 1870). This, undoubtedly, is the most correct and effectual method of curing the disease, for it offers at once a natural passage through which the pus can gravitate, at the same time that it affords a ready means of syringing out the cavity with any disinfectant fluid we choose to employ, and, also, the difficulty of keeping an opening in the soft parts for a long time patent, is obviated. The great displacement of the eye, the interference with normal vision, the swelling, are the prominent symptoms

which we recognize in old-standing cases of this affection—all these are completely combated by the drainage-tube and the operation proposed by Mr. Lawson. But occasionally we meet persons who will not submit to this plan of treatment. There is no doubt that the unpleasantness of wearing a tube is great, and shuts out the patient from society for the time during which it has to be worn.

In both the cases I am about to describe, the disease had been of old standing; neither would allow me to operate and insert the tube. In No. 1, I had the advantage of having my opinion strongly endorsed by Mr. Streatfield, who was in communication with me in reference to the case.

CASE I.—*Frontal Abscess and accumulation in the Frontal Sinus, resulting in Destruction of portion of the Roof of the Orbit, and extending to the Dura Mater.*—Mr. J., aged fifty-nine years, consulted me for epiphora, in January, 1871. *History*—Several years since operated on by Mr. Butcher, successfully, in the presence of Mr. Cusack, for a tumour over the eye, which he thinks was of a cystic nature; the eye has lately become more depressed, and the vision not so good as formerly. The patient suffers from diplopia, and has to close the sound eye to see objects well, as otherwise he gets a double image. He says he has then to look up to enable him to see an object distinctly. He never had the least pain, there was no fulness over the lachrymal sac, and, on examination with a probe, I found but a slight obstruction in one canaliculus; the upper lid drooped, and there was a slight adhesion and puffiness of the lid, which was marked by the remains of an old cicatrix where the original operation had been performed. His health had been otherwise good for years; he lived freely, and complained of nothing save an old chronic psoriasis of the scalp, which occasionally burst out, and from which he was suffering when I saw him. When he first consulted me I was inclined to attribute the changes in the position of the organ and the fulness in the lid, to the ulterior results of the previous operation, and considered that the watering of the eye might be due simply to obstruction in some portion of the lachrymal apparatus. I accordingly advised that I should operate, and slit up the canaliculus, and pass a probe into the canal, thinking that if any accumulation connected with the sac existed in the orbit, I would come on it in this manner. This, or any other operation at

Fig 1

Fig 4.

the time, he would not consent to, though I pressed him much to allow me. Not long after he got an accute attack of eczema of his face, scalp, and ears, which was very severe, and he also passed a gall-stone, from which he suffered intensely. I attended him, and was surprised to find that one day he got a profuse discharge of purulent matter from the nose, increased by pressure on the ball of the eye, and coming away in quantities when he blew the nose. The eye now began to water constantly. It was with great difficulty I persuaded him to permit me to perform any operation; however, on the 8th of April I opened up the sac, from which a quantity of foul, purulent matter came. I immediately enlarged the opening with a bistoury freely, and evacuated a quantity of the most foetid discharge, of a semi-transparent, slimy nature, and on examining with a probe, I found that the bone was extensively diseased. On pressing on the ball of the eye a large quantity of foul, purulent, and gelatinous discharge came, and I found the probe passed far upwards into the orbit and into the frontal sinus. For some few days I kept the wound well open, but on the third day the previously copious discharge ceased, the eyelid and brow began to swell, and he got an attack of erysipelas in the side of the face and head. On the 25th of April, under appropriate treatment, he had got much better of this complication, and as I was obliged to go to England for a few days, I left the case in Dr. Cummins' hands until my return. During my absence the wound was kept as well open as it was possible. Already there was a tendency to contract greatly, and accordingly I advised on my return, conjointly with Mr. Streatfield (to whom I had spoken of the case when in London), that he should have a drainage-tube passed, and a proper communication established. This he would not listen to, so I had no alternative but to insist on making a free incision into the orbit at its inner angle, and parallel with its upper margin, hoping that I would be able to devise some plan of keeping this open, and certain that an extensive cavity existed filled with pus, which might, if not evacuated, give rise, after so long a period of incarceration, to dangerous results; accordingly, by the aid of chloroform, on the 10th of June (with Drs. Cummins and Grattan) I did so, and evacuated a large quantity of the most foetid pus mingled with gelatinous discharge. I then examined the cavity carefully, having first syringed it through an Eustachean catheter, which was most convenient, as I was enabled to turn it in all directions inside. I found that the roof of the orbit was partly destroyed, a large

ragged communication existing between the orbit and the frontal sinus. The catheter easily passed in to the extent of three inches, and I could turn the curved end completely round. The following day I passed a self-retaining winged catheter into the cavity through which the fluid drained, and I had the cavity syringed out twice daily with permanganate solution. I found, on watching the flow of pus, that it welled up to the orifice of the wound with each cerebral pulsation, and with these pulsations there continued all through, while the discharge lasted, a constant synchronous rise and fall of the fluid at the opening. This I attributed to the cavity communicating with the dura mater. The winged catheter was now kept constantly in, and the wound enlarged daily with a probe. Weak solutions of carbolic acid were alternated with the permanganate, and the cavity well cleared out twice daily. He suffered on two occasions from rigours, followed by hectic and severe constitutional disturbance, with vomiting; and on the 18th of July, when everything appeared to be going on favourably, he got a severe febrile attack with an erythematous blush all over the scalp, which I was afraid would eventuate in general erysipelas. By the aid of salines, iced drinks, and diaphoretics, this subsided. After each of these attacks there appeared to be a fresh secretion of pus internally, judging from the character and quantity of the discharge. He was now taking iodide of potassium and liquid extract of cinchona, and this he continued for a length of time. The opening now began to contract gradually, notwithstanding that I slightly enlarged it on one or two occasions, and daily used the probe and catheter. I accordingly saw that before long the sides of the flexible and soft catheter would be encroached on, and the exit of pus prevented. On the 30th of July, the discharge of pus having greatly decreased, and there being no return of any bad symptom, I put in a tube, of the pattern figured, ordered by me from Messrs. Matthews, of London, a small silver canula (and subsequently a gilt one), one inch and a half long, through which passed two diverging blades of spring-steel, united below, and the distance of the wings at the orifice from each other, as also the extent of their protusion, being regulated by a screw handle. This acted as a holder, and kept the canula in position.

The tube was easily withdrawn by the patient himself, or the cavity syringed out through it, on the removal of the central

piece, by simply pressing in the tube and withdrawing the holder. He continued to do well, and on the 28th of August I made the following note in my case-book:—"Pus almost completely disappeared; a mere sign twice daily. Wearing tube, able to go about with protectors, and enjoy himself; using at present one grain nitrate of silver solution, and permanganate alternately." On the 1st of September, there being no apparent discharge of purulent matter, and the eye having perfectly returned to its normal position, I determined to remove the tube and let the opening heal.

I saw him again on the 4th; there was no pain; he said he felt quite well. On the 7th he went yachting, remaining out until late. On the 8th he came to me with a slight discharge of pus from the wound, which had now nearly completely closed. I immediately freely re-opened the wound (under nitrous oxide gas, given by Mr. Corbett, junior); a quantity of pus had evidently re-formed, and this was evacuated. From this out he got on very well. Though the discharge of matter had almost completely ceased on the 25th of September, save a slight flocculent sign, which always came after syringing, I did not again remove the tube, which now remained in without difficulty, and required no catch to support it. He alternately wore the gilt tube, or the end of a winged catheter. The vision was completely restored, and is now perfect. He enjoys better health than he has had for many years. The eye itself has returned to its normal position; the pupils are on the same level. (See Plate VIII., Fig. 1.)

The photograph was taken early in 1872. He still wears the tube, as at times there is a very slight appearance of a discharge, after which he syringes with a mild carbolic solution. He is in perfect health, and his vision is excellent. The following letter shows his present condition:—

"ALMA VILLA, BALLINTEMPLE,

"CORK, August 16th, 1873.

"MY DEAR DR. JONES,—I desire to inform you that my eye has now, and for the last fifteen or eighteen months, completely returned to its former position. The vision is excellent. I enjoy the best of health, and, but that I am nervous lest any of my former troubles should return, I could, with safety, remove the small tube. This, however, does not prevent me enjoying a good bout of my favourite game of whist, or to take my accustomed cruise in my yacht. When in town next I shall call to see you. Yours ever sincerely,

"GEORGE JOHNSON.

"H. M. Jones, Esq., M.D., Camden-place, Cork."

CASE II.—James T., aged forty-six, a carpenter, came to the Ophthalmic Hospital, on the 1st of April, 1871. He was able to count fingers and discern objects; he could read twenty Snellen at fifteen feet; his vision was diplopic. The eye was much displaced, being pushed upwards and outwards, and situated quite obliquely in the orbit, causing great disfigurement. There was a large swelling at the inner angle of the orbit, filling it up, above two inches in circumference. This swelling had lasted for about two years, and gradually increased to its present size, the eye being pushed entirely out of position as it increased. I had a drawing of the eye taken at the time. (See Plate IX., Fig. 2.)

He would not consent to any operation, and it was with difficulty I persuaded him to allow me to freely open the abscess, by a deep incision over the extent of the tumour. I evacuated a large quantity of pus, and found that the probe had extended a long way back into the orbit, and, as I anticipated, the bone was carious and necrosed. The fluid which came had a very foul odour, and was characteristic of diseased bone. I treated this case in the same way as the preceding. He attended the hospital for about three months. I kept the wound open by means of the constant syringing out of the cavity through an Eustachean catheter, and pieces of linen saturated with chloride of zinc solution, which I passed into it. By the repeated cleansing out of the cavity with solution of permanganate of potash or chloride of zinc, I succeeded in finally arresting all discharge; the cavity gradually narrowed, and I allowed the wound to heal in about three months from the time he applied. The photograph was then taken as represented in the drawing, and he had since continued at his employment, and his vision is perfectly restored. (See Plate IX., Fig. 3.)

CASE III.—*Melanosis of the Orbit*.—Mrs. R., widow, aged fifty, consulted me in August, 1869; then had a sightless ball, and only a vestige of cornea, greatly enlarged and prominent, with a peculiar elastic feel, giving the sensation of a large cystic tumour; lids movable, but partly adherent, especially the lower; the ball slightly movable, and discoloured from the action of the nitrate of silver; no pain generally, but at times—attendant on sudden changes in the atmosphere—pain of an excruciating character comes on, which stretches to the side of the head and brow; her general health at present good.

History.—Right eye affected for the past eight years; commenced after cold, and she was treated several times for attacks of inflam-

Fig. 2

Fig. 1

mation by leeches. Two years since the eye began to increase in size, and of late has grown more rapidly. Left eye perfectly healthy, naturally myopic, and dark pigment in iris. I advised immediate removal of the diseased ball, suspecting the disease to be of a malignant nature; she hesitated, and I did not again see her until the 17th of October (Sunday), when I was hastily summoned to her residence in the country, with a report that she was alarmingly ill. Frost had set in during the past few days, and the bad symptoms had supervened. I found her unable to speak, with a rapid and weak pulse, and for the preceding twenty-four hours she had been attacked with purging and vomiting; she begged for immediate operation. I returned home, and bringing with me Drs. Cummins and Grattan, I proceeded to remove the diseased mass, uncertain as I was of the nature and extent of the disease. She was thoroughly under chloroform. Immediately on freeing the adhesion which partly bound the upper eye-lid, and opening the conjunctiva, a quantity of black, gelatinous, melanotic matter burst out, of almost fluid consistency, and we at once recognized the terrible form of disease we had to deal with. I immediately enlarged the palpebral aperture, and freed the lids; then, with scissors and scalpel, I removed the ball and the contents of the orbit, until I cleared it out as far as possible, owing to the extremely severe hæmorrhage. The disease had evidently attacked chiefly the upper and back parts of the orbit, in which places it was of a firm consistence. The patient lost a considerable quantity of blood, and this continued for some time after I had ceased to operate; however, by pressure, and the use of perchloride of iron, it was arrested (I had no cautery with me at the time). The orbit was now plugged with pads soaked in strong perchloride solution, and we left the patient in very little pain, Dr. Grattan keeping her under the influence of chloroform for some time after the operation; all subsequent pain was relieved by opiates, and the application of iced bags to the temple and brow; suppuration set in, and the plugs were removed about fifty hours after the operation; the orbit was syringed out with strong chloride of zinc and carbolic acid injections. The sloughs gradually separated with poulticing, and there appeared a fair granulating surface underneath. She progressed rapidly in her general health and spirits; the lids fell in, and finally became adherent. About three months after the operation, I brought her case before the Cork Medical Society,

making these remarks:—"Her personal appearance is greatly improved; she is able to attend to her domestic and business affairs (which, as she has a large family, is of great importance to her); her general health and spirits she asserts, 'are better than they have been for a long time previous to the operation.'" I need not say that, knowing the nature of the case, I did not expect more than a merely temporary and palliative relief, and as the subsequent history of it will show, the anxiety with which I used to look forward to her future state of health was not without a cause. I ceased to visit her early in November, 1869. For the next six months I saw her but seldom, and there was no change of any importance in the condition of the affected part; she managed all her business affairs, the only thing being that on two or three occasions she exposed herself to cold, going about her farm, and got a return of the old neuralgic pain in the seat of the tumour. She went to the sea-side in the summer of 1870, and her general health improved greatly; she came to me, during this time, once or twice, and I was sorry to perceive that there was a decided tendency to re-growth, and prominence of the lids, with slight pain on pressure; however, she continued fairly until the 20th December, when she got sudden pain from exposure to cold. The part was greatly swollen on my visiting her; leeches, ice, chloral, and opiates, gave her relief, and the swelling partly subsided, but from this out there was a slow but steady growth from the orbit, accompanied at times with great pain, which nothing relieved but chloral in small and repeated doses. The tumour gradually increased, until, in April, 1871, it assumed the size and appearance depicted in the drawing. (See Plate VIII., Fig. 4.)

It was then soft and fluctuating to the feel, especially round the superior border of the orbit. It had in parts a dark livid appearance, and large veins covered its surface. It encroached on the nose and temple, and the pain was occasionally relieved by profuse hæmorrhage from the adjacent nostril. She now became anxious for re-operation, which I was, at first, totally against, and explained to her and her relatives, fully, the hopeless nature of any such step, and that at best it could give her only temporary relief. However, the pain became unbearable, and, unless when under the influence of chloral, her screams and agony were most distressing to all around her. I was most unwilling to assume the entire responsibility of the case, and as I was going to London, for my holiday, about this time, I took a drawing of the tumour with me, and

consulted a few friends there on the desirability of an operation. They, as far as they could judge (without actual observation of the patient), were in favour of giving her a chance, *if she desired it*, of prolonging life and obtaining relief from pain. She pressed urgently for operation on my return, the pain becoming, if anything, more agonizing, and the tumour enlarging. Just then I received (in reply to a letter of mine on the case) a communication from my friend, Mr. Streatfield, in which, after stating his opinion as to the certainty of return, he also advised operation—"seeing that her health was otherwise good, and to give her another remission from pain." On the 28th of May, over one year and seven months after the first operation, Drs. Cummins, Curtis, and Grattan assisting me, I, by two oval incisions, removed the tumour, about the size of an orange, which completely filled the orbit, and was of a distinctly cystic nature, being enclosed by a firm envelope, the entire tumour shelling out. The bleeding was arrested by the actual cautery, which was freely applied to the raw surfaces, the orbit then dressed with pieces of lint soaked in chloride of zinc paste. On removal of the tumour a fungoid mass was found protruding from the ethmoid bone, the orbital plate of which, and a portion of the frontal, were completely gone. Pain entirely subsided after the operation, from which she recovered well. She was supported freely by iced drinks, beef-tea and wine, and for two days she took one grain of opium every four hours. She would not permit the chloride of zinc paste to be applied to the raw surfaces, as I desired, and I had to be satisfied with strips of linen, soaked in chloride of zinc and carbolic solution, placed in the orbit.

There was now little or no pain; but the fungoid growth, from the inner wall of the orbit, increased rapidly, and was attended by an occasional sanious discharge from the right nostril, and also from its surface. Matters continued thus until the 30th of June, when there was sudden profuse hæmorrhage, and also a discharge of purulent matter from the growth. She had now for some time cessation from severe pain, until the 25th of August, when I again saw her. The tumour had in the meantime increased enormously, projecting from the orbit, and pushing the nose aside, a huge "*fungus hæmatodes*," which had become intensely painful, and which at times bled profusely. The hæmorrhage partially relieved the pain, assisted by subcutaneous injections of morphia, administration of chloral, and iced bags applied to the forehead and

temple. She finally sank in September, 1871, about one year and eleven months after the date of the first operation.

This case is instructive in some particulars. It adds one more to the list of those which prove the absolute necessity of early, *very early* operation in cases of suspected malignant disease; its slow growth; the comparative freedom from pain for a long period; the mobility of the ball; the freedom of the lids; and the absence of external signs of the disease, were uncommon in a case of such extensive malignancy. The disease did not at any time appear to attack any other organ, as her general health remained fair, until at last she succumbed to the continued pain. The advantage of operation (irrespective of all personal considerations), even at the eleventh hour, was manifest, inasmuch as it was after the last operation that she was enabled finally to arrange her affairs, which were much confused, and to leave her orphan children beyond the power of litigation. To relieve suffering, to prolong life, are two great objects to be gained by any operation, and certainly in this instance both were attained.

CASE IV.—*Epithelioma of Orbit.—Removal of Contents.*—Jeremiah S., aged sixty, admitted to the Cork Ophthalmic and Aural Hospital on the 9th of September, 1870. *History.*—From the time he was three years of age he had a small red pimple at the outer margin of the orbit; about three years since he noticed it enlarging, and then it rapidly spread. He was treated for the growth in various ways by several medical men in America. It quickly attacked the eye, and it is now over twelve months since he had the least vision in the affected ball. He presented himself to me in a state of horrible deformity; the greater part of both lids was completely destroyed, especially the lower; there was a great chasm between the ball and the inferior margin of the orbit; it had lost all power of motion, and I was barely able to discern the situation of the cornea; the skin round the orbit was partly involved, especially at the outer margin; in short, the whole orbit presented to view an unsightly red mass, the disease evidently extending deeply, involving all the muscles, some of which were completely destroyed. Lately he has suffered some pain, and his general health is declining. On the 9th of September, Dr. Grattan having placed him well under chloroform, I removed the whole contents of the orbit, completely clearing it out, and also removing the diseased portion of integument and tissues at its edges. I then

lightly ran the actual cautery over the whole, and filled the orbit with pledgets of lint, soaked in chloride of zinc paste. He then got a full dose of chloral, and from his admission until his departure from hospital he suffered no pain. The sloughs separated with carbolic poultices in some days. He was up the following day in the ward, and was permitted to walk about outside on the third day after the operation. Healthy granulations sprung up, and he left the hospital on the 28th of October, the interior of the orbit, the cavity of which had a good deal contracted, presenting a healthy smooth surface, the only part which had anything of a suspicious appearance being the outer margin; here the integument looked vascular and thickened. While in hospital he was syringed out daily with chloride of zinc and carbolic acid solution, and the orbit dressed with carbolate of zinc ointment.

Since writing the above, I have managed to trace this patient, who lives a long distance from the city. I learn that the disease has again made its appearance, extending not so much in the orbital as in the temporal direction and on the face. His health, however, is good, and he suffers no pain.

ART XII.—*Three Thousand Cases in Practical Medicine. An Analysis, with Remarks.* By RICHARD J. HALTON, Licentiate of the King and Queen's College of Physicians; Licentiate of the Royal College of Physicians, Edinburgh; Licentiate of the Royal College of Surgeons, Ireland; Licentiate in Midwifery and Diseases of Women, King and Queen's College of Physicians, Coombe Hospital, &c.; Medical Officer to the Kells Dispensary, and Medical Officer of Health to Kells.

IN compiling the present paper I have had it principally in view to give a list of all the diseases met with during two years in the practice of the Kells Dispensary District, distinguishing those which occurred with most frequency; and as some diseases had come under observation in private practice which did not happen to occur in those years in the dispensary, I thought a list of them might not be without interest. I have selected reports of a few cases out of many hundreds carefully written, which may serve as specimens of some of the common forms of disease of every-day occurrence in country practice. I have detailed shortly the treatment pursued, and accurately recorded the results.