TRANSACTIONS

OF THE

PHILADELPHIA ACADEMY OF SURGERY.

Stated Meeting held October 7, 1912.

DR. GWILYM G. DAVIS, President, in the Chair.

SPRAIN-FRACTURES.

DR. PENN G. SKILLERN, JR., presented skiagraphs of cases of sprain-fracture as follows:

CASE I.—*Sprain-fracture of coracoid process of scapula.* A football player, aged 20, fell upon his right shoulder, causing luxation at acromioclavicular joint. Skiagram (Fig. 1) showed a scale of bone torn off from the coracoid, probably from traction upon the coracoclavicular ligaments. Fractures of the coracoid process are rare, their line usually involving the base. The frequency of combination of this sprain-fracture with luxation at this point has not been established.

CASE II.—*Sprain-fracture of wrist.* This skiagram (Fig. 2) of an ordinary "sprained wrist" showed avulsion of a scale of bone from the dorsum of the carpus, probably from the os magnum. This scale was not palpable on account of the swelling, but there was distinct localized tenderness over it. A skiagram should be made of every "sprained wrist" and the treatment should be immobilization.

CASE III.—*Sprain-fracture of anterior superior spine of ilium.* Male, aged 16, during a foot race felt something snap in upper part of left thigh, but finished race (five yards). Pain aggravated by flexion of thigh. Skiagram (Fig. 3) showed avulsion of a shell of bone from the anterior superior spine and its immediate vicinity, evidently from action of the sartorius muscle.

CASE IV.—*Sprain-fracture of cuboid.* S. K., male, aged 23. Twisted left foot inward and heard something crack, immediately after which swelling appeared at external tarsometatarsal joint. No previous injury here. Examination showed swelling and ecchymosis between external malleolus and this joint, and defi-

nitely localized tenderness at antero-external corner of cuboid, suggesting sprain-fracture of same. Skiagram (Fig. 4) showed chipping off of a sliver of bone from antero-external corner of cuboid, evidently from overstretching of the dorsal tarsometatarsal ligament at this site. Foot strapped in over-abduction with relief of pain.

CASE V.—*Fracture of adductor tubercle of femur.* Boiler-maker, aged 20, received a blow upon the lower part of the left thigh. There was localized tenderness just above internal condyle. Skiagram (Fig. 5) showed separation of adductor tubercle, and the tendon of the adductor magnus leads to it as a shadow.

CASE VI.—*Fracture of sustentaculum tali.* Male, aged 35, fell from a height of 10 feet, landing on feet. Skiagram (Fig. 6) showed an impaction of the sustentaculum into the body of the os calcis.

CASE VII.—*Fracture of clavicle, sternal end.* Male, aged 40, was struck by a heavy object upon the right clavicle. Examination revealed a dense and tender swelling over the clavicle near the sternum, which to inspection resembled a neoplasm and a luxation at the sternoclavicular joint. Skiagram (Fig. 7) revealed a line of fracture within an inch of the sternoclavicular joint. In the literature this fracture is very infrequently met with.

CASE VIII.—*Syphilis hereditaria tarda of femur.* Male, aged 22, farmer. Except for lesion in left thigh is robust and healthy. Three years previous to admission had what was diagnosed and treated as a fracture of the femur. For several years before that had had trouble with left femur, giving rise to a perceptible limp. Examination revealed marked bowing of left thigh, the point of greatest convexity being 13 cm. below the anterior superior iliac spine. Left thigh 6.5 cm. shorter than right. On palpation the upper part of the femur was of great uniform diameter, markedly roughened, but not tender. No inflammatory manifestations, no areas of softening, no sinuses. There was no history of tuberculosis, malignancy, or syphilis in the family. Diagnosis of late hereditary syphilis of femur made. Skiagram (Fig. 8) revealed marked increase in diameter of upper half of femur; obliteration of medullary cavity; alternating areas of osteoporosis and osteosclerosis; and the line of an incomplete fracture at the point of greatest convexity of the femur. Wassermann reaction positive (Ivy). Mercury and iodide treatment

FIG. 1.

Fracture of coracoid process scapula.

FIG. 2.

Sprain. Fracture of wrist.

FIG. 3

Sprain. Fracture of anterior superior spine of ilium.

FIG. 4.

Sprain fracture of tarsal cuboid.

FIG. 5.

Separation of adductor tubercle of the femur.

FIG. 6.

Fracture of sustentaculum tali.

FIG. 7.

Fracture of internal end of clavicle.

FIG. 8.

Hereditary syphilis of femur.

FIG. 9.

Mesocolic fossa of extraordinary depth.

was instituted and an orthopædic splint adjusted by Dr. Willard so as to transfer weight supported by left lower limb from ground to pelvis as a base of support. After 8 months, while the patient had better use of and less inconvenience with the limb, yet skiagrams indicated but little change in the condition of the bone, and quite recently a dose of neo-salvarsan was administered.

CASE IX.—*Mesocolic fossa of extraordinary depth.* Found in a female body in the angle between the duodenojejunal flexure and mesocolon, this fossa measured 3.5 cm. in width, and 4 cm. in depth (Fig. 9). The mouth of the fossa was bounded on the left by the left colic branch of the inferior mesenteric artery and vein; in front by an arterial arch which connects the middle colic artery with the left colic. Leaving this arch at its middle and behind it is the inferior mesenteric vein, while attached to the arch and enclosing it is the posterior lamella of the mesocolon. On the right the mouth is bounded by the middle colic artery and the trunk of the superior mesenteric. Posteriorly is the third portion of the duodenum and the renal artery.

The anterior wall of the pouch is bounded by the pancreas; the posterior wall by the duodenojejunal flexure; the right by the superior mesenteric vessels; and the left by the descending colon and anterior half of the left kidney.

This fossa offers an excellent site for lodgment of an internal abdominal hernia.

DR. W. G. ELMER remarked that one should be careful not to be led into error by examination of a late X-ray picture. If the ligament tears away the osteogenetic layer from the bone, it is open to supposition that a small island of bone may be formed in the vicinity of the bone from which it was supposed to have been torn away.

DR. ASTLEY P. C. ASHHURST said that it seemed to him that there was no need to exaggerate the frequency of sprain-fractures, when it was possible to explain the lesions in other ways. Drs. Ross and Stewart (*ANNALS OF SURGERY*, 1912, ii, 599, Fig. 4) included in their series of sprain fractures an experimental fracture of the coronoid process of the ulna which was quite clearly caused by direct violence and not by ligamentous distraction. So in the case reported by Dr. Skillern, it seemed to Dr. Ashhurst that the fracture might very well have been caused by direct violence.

REDUCTION OF OLD DISLOCATIONS OF THE SHOULDER.

DR. T. TURNER THOMAS presented a paper with the above title, for which see page 217.

DR. OSCAR H. ALLIS said that he possibly stood alone in one respect, he does not think any one gets a perfect result in dislocation of the shoulder. The mechanism of the shoulder is very peculiar in some respects; as all know, muscle has the function of stretching within prescribed limits. When the head of the humerus is thrown suddenly into the axilla the short muscles of the scapula, its normal retainers, are apt to be torn in two or stretched with interstitial tearing of the muscle-fibres, so that after restoration there will be a later atrophy and tendency to a second dislocation. A tendency to a second dislocation is not necessarily because the capsule does not unite, but because it has no longer these little muscles to hold it in position. He was inclined to think that no one ever gets the full amount of motion after one of these dislocations. In a paper read before the College of Physicians on one occasion he stated the theory that the dislocation takes place through muscular action when the arm is thrown away from the body. He had seen a number of such cases. A person going downstairs makes a misstep and throws the arm out, producing a dislocation, although the individual does not fall. When Dr. Thomas said that he does not believe that the opening or laceration in the capsule is ever so small that the head cannot be replaced, he was sure experimental work bore that out. The old theory of a slit in the capsule cannot be held by any one that has done any experimentation.

DR. D. L. DESPARD said that traction prolonged over a number of days possesses some advantages over the method of applying a great force only at the time when reduction is being attempted. The former method of traction was successfully used recently in the dislocation of the shoulder of an exceedingly muscular man. The injury had existed for over six weeks and he had anticipated a great deal of difficulty in effecting a reduction. By means of Buck's extension apparatus, traction was started with 6 pounds, which was gradually increased to 10 pounds and continued over a period of 8 or 9 days. The patient was then anæsthetized and reduction effected by the Kocher method upon the first attempt.

DR. JOHN H. JOPSON said that he had always found Dupuytren's modification of Mothe's method easier than the Kocher method. It consists in drawing the arm directly upward fully extended in a broad sweep while counter-pressure is made by an assistant's fist in the axilla.

DR. A. P. C. ASHHURST remarked that all methods of reducing dislocations might be classified, either as *direct* or *indirect*. Dr. Thomas advocates one of the direct methods; it is the same as Stimson's or Sir Astley Cooper's, only it is applied differently. Stimson puts the patient in a sort of sling and lets the arm hang through a hole in this sling, attaching a weight to the hand. Cooper pulled the arm away from the chest, using the foot for counter-pressure. When the head of the humerus is brought away from the side of the chest and opposite the glenoid process of the scapula, then it is pushed or pulled into the socket, either by manual pressure as in Stimson's and Thomas's methods, or by leverage over the foot, as in Astley Cooper's method. All of these are direct methods, similar in principle to Allis's method for reduction of dislocations of the hip. In this sense there was nothing new in Dr. Thomas's application of this principle to the shoulder. Henry H. Smith's, Kocher's, and other methods of the kind are so-called indirect methods, like Bigelow's.

It was interesting, Dr. Ashhurst thought, to recall that anæsthesia not only made reduction much easier, but that it demonstrated that it was not only the muscles which interfered with reduction. Though Kocher was the first to recognize that the ligaments were the main obstacle to reduction in dislocations of the shoulder, as Moses Gunn had been the first to recognize their action at the hip, Kocher's claim that the coracohumeral ligament acted at the shoulder as did the iliofemoral at the hip was demonstrated to be false by Farabeuf, who showed that the essential structure was the posterior part of the capsule.

The end results of dislocations of the shoulder formed an interesting subject. Though the reports of consecutive series of cases are meagre, it appeared that two out of three patients have been found to present persistent debility. Out of more than 20 patients under the speaker's own care, it had been possible so far to ascertain the end results only in five cases. In only two of these was perfect function regained; the three other patients were quite satisfied with their condition, though two of these had had the same shoulder dislocated twice or more subsequently,

and the third had distinct limitation of motion, though subjectively he claimed his shoulder was "all right."

DR. GWILYM G. DAVIS remarked that this discussion had broadened out into recent as well as old dislocations. Etherization eliminates the consideration of muscle resistance, but as regards the difficulty of replacing luxations if no anæsthetic is given, then the muscles play an important part as a hindrance to replacement; in shoulder luxations it is often thought unnecessary to give an anæsthetic. Then the elimination of the muscles must be undertaken by other means, hence it is that use is made of the methods of Stimson, Astley Cooper, and others, by direct traction, etc., all with the idea of overcoming the muscle resistance. When this is overcome by an anæsthetic then the bones and ligaments only are to be dealt with, and the reason that the luxation cannot at times be reduced under these circumstances is because, as Dr. Allis has shown in the hip, the arm is not placed in the correct position. Thus, when Stimson puts the arm through the sling he lets the weight swing and the arm swings round until the capsule is opened and the head slips in. The same thing occurs in the Astley Cooper method,—when the arm is rotated it opens the rent and in goes the head, and it is the same with all the direct methods. In the old method of hanging the patient over the top of a door, the arm rotates until the capsule is open to its greatest size, when the head will slip in. If traction is made in abduction and backward, absolutely eliminating the muscles by anæsthesia, then the only thing to do is to rotate until the capsule is open, and push the head in. If it does not go in, then there is an irregularity in the condition; it is not a true simple luxation but one complicated with a fracture, etc. When it comes to the method, the speaker agreed with Dr. Thomas that this is the proper method of reducing luxations of the shoulder. He had had several cases in which he had put the foot against the bedstead and pulled outward and backward and rotated; the longest time taken for reduction was fifteen minutes, while in others it was effected in three minutes. Kocher's method is unnecessary although effective if it is desired to reduce without an anæsthetic. Reduction does not depend solely upon the coracohumeral ligament. This ligament goes from the coracoid process, which is to the inner side, outward and in front. It has near it the long tendon of the biceps and the glenohumeral ligament. Above is that part of the cap-

sule which frequently remains untorn in the cases in which the Kocher method is effective. One can determine on the cadaver that when the upper and posterior parts are intact one can rotate the humerus, and the head will move out from the chest toward the glenoid cavity. But if that portion of the capsule is torn, the head will bore between the glenoid process and the side of the chest, and in order to reduce such a luxation all that is necessary is to proceed with the direct method as the capsule is all torn. In such cases the Kocher method is ineffective. In the old cases in which that part of the capsule is intact then the Kocher method is effective because one can wind the remaining portion of the capsule around the upper portion of the head of the bone and push it outward, and it pushes in the old cases the supraspinatus, infraspinatus, and teres minor off of the glenoid process and cavity. The Kocher is the most dangerous method, because if the external rotation is made when the arm is down by the side, as is so often taught, then the coracobrachial is stretched so firmly over the head of the humerus that the lesser tuberosity catches on it and prevents it being rotated outward, and not infrequently the head of the humerus is jammed so tightly between the glenoid process of the scapula and the side of the chest that if one persists in rotating outward one will fracture the bone. It is far safer to do a wide abduction and traction.

DR. T. TURNER THOMAS (in closing) said that he did not mean to infer that the abduction method was new. It is much older than the Kocher method, but what he had tried to show was that one of the older methods is more effective than the new, or Kocher method, which has been the prevailing method almost from its introduction. In regard to the Stimson method for recent dislocations, the underlying principle is the same as that which he had been applying in old dislocations. With regard to the disabling limitation of movement after the reduction of old dislocations, if the dislocation has existed for many months before reduction, the chances for a complete return of function are small. He would not say that it is impossible to get it. The return of function will be more rapid and will more nearly approach the normal after a non-operative reduction than after an operative reduction, as a rule. The underlying cause of the difficulty is essentially the same as for the corresponding condition found after the reduction of recent dislocations, *i.e.*, the stiff and painful shoulder of which so much has been written in re-

cent years. The cicatricial contraction at the site of the capsule tear in the axilla will be more unyielding after the reduction of the old dislocations than after the reduction of the recent. He had, in a considerable number of cases of the latter variety, broken up this resistance by forcible manipulations under ether, without trouble and with very satisfactory results.

He had had very little experience with recent dislocations, since these are usually reduced by the family physician or the hospital interne, so that in discussing the abduction method he had confined himself to the old dislocations.

The superiority of the abduction method should be more evident in the reduction of dislocations of the shoulder associated with fracture of the surgical neck of the humerus, just as the Allis method is superior to the Bigelow method in the corresponding condition at the hip. The Kocher method may have the advantage in the reduction of recent dislocations, without an anæsthetic, in very powerful individuals. Such patients can resist more effectively the simple, direct pull in abduction than a series of more or less complicated movements as in the Kocher method, although the general spasm may effectively resist all these.

Stated Meeting held November 4, 1912.

DR. GWILYM G. DAVIS, President, in the Chair.

STAB-WOUND OF THE HEART; RECOVERY AFTER SUTURE.

DR. CHARLES F. MITCHELL presented a colored man, 59 years of age, who was brought to the Pennsylvania Hospital by the patrol at 5 P.M., July 30, 1912, having received a stab-wound of the left chest a short while before. He had been drinking heavily, and there was a marked odor of alcohol on his breath. His previous history was negative, except that he always used alcohol to excess. He was admitted to the service of Dr. Richard H. Harte, in whose absence Dr. Mitchell was called upon.

On admission temperature was normal, pulse 90 to the minute, while breathing was rapid and rather labored. There was no sweating, lips and conjunctiva blanched, heart sounds regular, but rather faint. Arteries atheromatous, marked arcus senilis. Area of cardiac dulness not increased. Right chest normal, but signs of pneumothorax over whole left chest. There was some

dulness of left chest posteriorly. In sixth left interspace in anterior axillary line was a stab-wound, about $1\frac{1}{2}$ inches in length, from which bright red blood was flowing.

From above symptoms and physical examination a penetrating wound of the chest was diagnosed with probable injury to the heart. At 6.24 P.M., 1 hour and 24 minutes after admission, the patient was given ether preceded by ethyl chloride. The field of operation was sterilized with 3 per cent. iodine solution and wound in interspace enlarged. Left lung was found collapsed. The sixth rib was then divided and retracted, and immediately a large opening was found in the pericardium. The edges were rough and the wound appeared to be more like a tear than a clean cut. There were a number of clots found in the pericardium, which when removed showed a transverse cut in the heart, from which at each systole there flowed bright red blood. The cut in the heart was about one inch in length, apparently in the left ventricle about an inch above the apex. A curved intestinal needle, threaded with fine Pagenstecher thread, was then passed through the cardiac muscle and tied, the ends being left long and used as a tractor in the introduction of the second stitch. When the second stitch was tied, it was found that the wound was completely closed and the hemorrhage from the heart stopped. The pericardium was then partially closed, after its cavity had been washed out with normal hot solution. A small gauze drain was left in the wound and the retractor holding the rib withdrawn, and the wound closed with silk-worm-gut sutures. There was no hemorrhage from the chest or pericardium, and no ligatures were used during the operation.

Previous to the operation the patient was so well stimulated by the alcohol already imbibed, that he did not require any stimulation, either before, during, or after the operation. He apparently left the operating table without any symptoms of shock, his temperature being 96.4° , pulse 88, respiration 32. Time of operation 22 minutes. During the night he was given a sixth of morphine hypodermically, but this is all the medication he received. The following morning his temperature rose to 101° , pulse remained about the same (92), respiration 36.

On August 2, three days after admission, he developed delirium tremens, and was irrational for a couple of days. On August 5 a to-and-fro friction rub synchronous with a heart-

beat was noted to the right of the sternum at level of the third rib. This disappeared in three days. There apparently was no increase of cardiac dullness or other signs of cardial effusion. Temperature at this time was 100°, pulse 100, respiration 28. Drain was removed on August 17. Patient sat up in bed at this time. On August 21 dullness in the left chest posteriorly, with distant breath sounds, was noted over this area. Temperature 102.3°, pulse 124, respiration 28. On August 26 chest was aspirated and about eight ounces of a dark reddish, clear fluid evacuated. Upon culture this was found to be sterile. On September 14 chest was again aspirated, but only a small quantity of the same sort of fluid obtained. From this time on, patient rapidly improved. Signs of fluid in left chest diminished and when patient left the hospital on October 19 there was but slight dullness over left chest posteriorly, probably due to a thickening of the pleura. The heart at this time seemed to be slightly pulled to the left, apex beat being in sixth interspace one inch to the left of the nipple line. There were no murmurs present, sounds regular but a trifle rapid. Dr. Mitchell added that so much has been written of late as to the treatment of heart wounds that it does not seem necessary at this time to go very deeply into this subject.

König¹ in his article on "Technic for Access to Suture of the Heart," gives a full discussion on this subject, and Poole,² gives a most exhaustive study of the technic, as well as the bibliography of recorded cases up to the year 1912. He has succeeded in tabulating 77 cases of heart suture, which added to those already tabulated in 1909 by Peck totals 236.

Ranzi³ gives Rehn the credit of publishing the first successful case of heart suture in 1896, and has collected 223 operative cases with a mortality of 53.3 per cent. He adds to this number three cases of stab-wound and also three of gun-shot wound of the heart, who were operated upon in Von Eiselsberg's Clinic at Vienna, but only one of which recovered. He mentions in the successful case, that five hours intervened between the time of injury and operation, and states in naming the time of the operation that the anæmia was not very marked.

¹ Deutsche Zeitschrift für Chirurgie, vol. cxii, Nos. 4 and 6.

² ANNALS OF SURGERY, April, 1912.

³ Wiener klinische Wochenschrift, Vienna, Dec. 14, vol. xxiv, No. 50.

Bircher⁴ reports a case of gun-shot wound of the heart which healed under conservative measures alone, and also reports a case of multiple stab-wounds of the heart requiring operative treatment, which case recovered. He goes on to say that only one stab-wound was found at the first operation and the second wound required suture twelve hours later. At the time of the second operation the wound first sutured showed that firm repair had already begun to take place. He states that conservative measures seemed more promising for gun-shot wounds, and intimates that operative procedures are necessary in all cases of stab-wound.

It is needless to say that all stab-wounds of the heart require surgical intervention, and that all wounds in the neighborhood of the heart should be explored, as this is the only positive method of determining the extent of the injury done.

When the symptoms of extreme shock accompany a wound of the chest in the cardiac region, the diagnosis is fairly sure of an injury to either the pleura, pericardium, or heart, but this cannot fully be determined without exploratory procedure; as in the case above cited, the symptoms were those of an injury to the pleura, whereas at operation we found not only the pleura injured, but the pericardium and heart as well.

As to the method of operating in these cases, I do not believe that any fixed rules can be laid down other than those of expediency. Usually the enlarging of the original wound (Peck) and the division of the costal cartilages, the retraction of which will allow a good exposure of the heart, is all that is required. Kocher and others suggest various flap methods. Kocher divides the fifth, fourth, and, if necessary, the third costal cartilages, while Wilms recommends the intercostal incision, as it can be much more quickly performed than the various flap methods. In the majority of heart wounds the pleura is injured. Sauerbruch says that 80 per cent. of the cases are so complicated.

As to the suture material, either well-vaselined silk, chromicized catgut, or Pagenstecher may be used. In our case Pagenstecher thread was the one selected. No doubt the use of the differential pressure apparatus is of great advantage in the administering of the anæsthetic, but when this apparatus is not available, ether, by the drop method, is the most efficient method.

⁴ *Archive für klinische Chirurgie*, Berlin, vol. xcvi, No. 4, pages 831-1075; last indexed, April 27, page 1318.

The pericardium may or may not be completely closed. If there is much injury it is better to partially close it by interrupted sutures and carry a small drain down to the opening that is left so as to drain the excessive serous discharge which is apt to occur as a result of the traumatism. Drainage of the pleural cavity may or may not be done at the primary operation; it depends on the likelihood of infection. In doubtful cases it should always be performed. Poole says it is better to delay drainage until infection has occurred and then to perform a secondary thoracotomy.

DR. FRANCIS T. STEWART remarked that it was an error to give credit to Rehn, of Frankfort, for the first suture of the heart. Farina and Cappelen each operated in 1896, but the patients died. In 1897 Rehn published the first successful cardiorrhaphy. So far as he was aware there had been 11 cases of suture of the heart in Philadelphia, one by Dr. Mitchell, two by Dr. Gibbon, one by Dr. Bradbury, one by Dr. Billings, one by Dr. Harte and five by himself, nine of these having been cared for at the Pennsylvania Hospital. As to the diagnosis, in the beginning it is often a matter of doubt. He had explored a number of thoraces for wounds and had found only five cases in which the heart was wounded, although in many a wound of the heart was suspected. Simply from the degree of shock no conclusion can be drawn. He remembered one case of stab-wound over the heart which appeared as if there must be a wound in this organ, but it was found on examination that the knife had not penetrated the thorax, the patient suffering only from emotional shock. In some of the cases in which the thorax is penetrated the heart is seriously disturbed because of so-called concussion of the heart, the heart being merely bruised. This is more frequent in gun-shot wounds. In several cases that he had explored the pericardium had been wounded but not the heart, but the symptoms were indicative of a wound of the heart. The most reliable symptoms, when they exist, are those of compression of the heart (cyanosis, distention of the veins from pressure on the auricles, etc.). These symptoms are not conclusive, however, because effusion of blood into the pericardium may result from wounds of the vessels of the pericardium or the great vessels at the base of the heart. The site of the wound is usually over the heart, although in some cases it is in the axilla or even in the abdomen. No conclusions can

be drawn from external bleeding, because serious bleeding may proceed from a wound of the internal mammary artery or from the intercostals. The diagnosis can be assured only by exploration. His own custom in these cases has been to incise the skin; if the wound penetrates the muscles, to incise the muscles; if it penetrates the thorax, to enlarge the wound throughout its entire depth and to expose the pericardium; if a wound is found in the pericardium, to enlarge that wound, perhaps by resection of a rib above or below, or both. But if, on exploration, the symptoms of a wound of the heart being present, it is found that the pericardium has not been wounded, or no wound is discovered, then the pericardium should be punctured with a needle, because, although one who has had no experience in this class of surgery may think it easy to determine whether or not blood is in the pericardium simply by inspection, it is not always an easy matter, as has been proved by several reported cases. If there is doubt after a needle has been put in the pericardium, this membrane should be incised in order to allow full exploration. As to the method of exposure, it is a matter of expediency. The size and shape of the incision or flap must be determined by the situation of the external wound and the situation of the stab in the heart. If a flap, consisting of one or more ribs, is turned inward, the pleural cavity will always be opened. If a flap is turned up or down double section of the intercostals is necessary. Whenever possible the flap should be turned outward, toward the arm. In this way, if need be, the pleura can be separated without injury, and the exposure made extrapleurally, as in one of his cases, a wound of the auricle, where he was able to make a large flap, to push off the pleura, which was not wounded by the knife, to expose the pericardium over a wide area, and to suture the wound in the heart without injury to the pleura; the patient made a rapid recovery. If the pleura is wounded infection usually follows. About one-half of the cases die of empyema of the pleural cavity or pericarditis or infective myocarditis, etc., so that if infection can be avoided the number of recoveries will be vastly increased. Of those that recover, about one-third do so in spite of infection. Of the five cases operated on by himself three recovered, two in spite of infection. As surgeons now recognize that wounds of the heart should be sutured and it has been demonstrated

that the hemorrhage can be controlled, the greatest problem is to learn how to prevent the infection. He mentioned three things as being of some value in this direction. First, the rapid disinfection of the skin with iodine. In most of the cases he had operated on he used soap and water, alcohol, and bichloride of mercury, which takes a little too long, if done properly, and is not very sure if done hurriedly. With a 10 per cent. tincture of iodine solution the disinfection can be done rapidly and certainly. In the second place, if possible, drainage should be avoided; careful hæmostasis should be made and clots removed, as those which remain either form adhesions or encourage infection. If empyema arises later it can be drained. Third, that the presence of air in the pleural cavity must if possible be avoided, at least after operation. He did not know of any case having been operated on with the positive or negative pressure apparatus. In his last case he intended to use a home-made Auer-Meltzer apparatus, but the different parts could not be assembled quickly enough. The question arises, however, as to whether suction upon or distention of the lung may not increase the bleeding. If a positive or negative pressure apparatus is not at hand, the air in the pleural cavity should be removed by aspiration, after the wound in the chest is closed. A pleura full of air contains a large number of bacteria, which, after they have settled on the pleura give rise to infection. If the lung can be expanded the chances of infection will be less. In one case in which he opened the thorax for exploration, finding a wound of the lung and not of the heart, the patient was treated in this way and recovered without difficulty or infection.

DR. W. JOSEPH HEARN said that a few years ago a colored woman was brought to the Jefferson Hospital one morning with evidence of puncture of the heart (stab-wound), and one of his medical delegates examined her carefully with the stethoscope and was satisfied there was leakage from the heart and suggested immediate operation, which he attempted. He made a U-shaped incision, turning the two bows of this flap toward the sternum, and avoided cutting the mammary vessels. He made an incision through which he could almost put his hand, and after washing out with salt solution, found an opening one inch long in the pericardium; the pericardial wound was enlarged sufficiently to see there was no wound of the heart,

merely a scratch, which had been made by the point of the knife. It was simply a wound of the pericardium into which he put 2 or 3 sutures, closing the wound without drainage. The woman 2 years later died of phthisis. His method of exposure gave a good view of the heart without much hemorrhage. The only difficulty encountered in sewing up the pericardium was the heart movement.

PERIRENAL HÆMATOMA.

DR. JOHN SPEESE read a paper with this title.

DR. FRANCIS T. STEWART remarked that he had never seen a spontaneous perirenal hæmatoma, but when Dr. Speese read his explanation of the tympanites it reminded him of the tympanites of other renal lesions, particularly of renal colic, which must be purely nervous in origin. On two occasions he had been asked to operate on a patient for intestinal obstruction who was found to be suffering from renal colic, and recently he had seen another case of the same character.

DR. CHARLES H. FRAZIER was reminded of a case presented by him to the American Medical Association five or six years ago. This was a young man 25 years of age, who was brought to the hospital 36 hours after the onset of his illness, believed to be suffering from an acute abdominal lesion. Upon examination after admission there was found board-like rigidity of the right side, exquisite tenderness on pressure midway between the appendix and gall-bladder, and also tenderness, but not to the same degree over the right kidney. There was marked acceleration of the pulse, leucocytes were 25,000, and an elevated temperature was present. An exploratory incision was made in the right rectus, nothing was found in the peritoneal cavity and the wound was closed. An incision was then made over the right kidney. A very large perirenal hæmatoma was exposed. Drainage was introduced and the patient made an uneventful recovery. The hemorrhage was attributed to a cortical tubercular lesion. The patient had pulmonary tuberculosis.

AN APPROACH TO THE HYPOPHYSIS THROUGH THE ANTERIOR CRANIAL FOSSA.

DR. CHARLES H. FRAZIER read a paper with the above title, for which see page 145.

DR. JOHN H. JORSON mentioned a case of fracture of the skull in which the line of fracture extended to the roof of the

orbit and the patient developed a panophthalmitis. Dr. Shoemaker, who performed the enucleation, stated at the time that this condition was a not infrequent complication of fractures involving the orbit. It had occurred to him that in the operation described by Dr. Frazier, some such pathological condition in the eyeball might result.

DR. CHARLES H. FRAZIER remarked, in response to what Dr. Jopson had said of involvement of the orbit, that great care should be exercised in separating the periosteum from the roof of the orbit before any attempt is made to remove the bone. The periosteum is quite thin there and may be readily torn unless one proceeds cautiously. It is rather presumptuous at this juncture to say that the transfrontal method of approach will be preferred to others, particularly the transphenoidal route. He could not, however, help but feel that surgeons will never become accustomed to working through such a long and contracted avenue as is necessary when approaching the sella through the sphenoidal sinuses. For the nasal specialist who is accustomed to open and drain the sphenoidal sinus, it may be a simple matter to go a step further and remove the thin shell of bone which forms the floor of the sella turcica. Or if nothing more than the removal of the floor of the sella, a sella decompression, is contemplated, the transphenoidal route may be given preference; but if one wants an exposure of the sella turcica sufficient to enable one to see the character of the lesion to be dealt with, some method other than the intranasal method of approach will be found to be absolutely essential.

THE FORMATION OF AN ARTIFICIAL VAGINA BY INTESTINAL TRANSPLANTATION.

DR. FRANCIS T. STEWART read a paper with the above title, for which see page 210.

TO CONTRIBUTORS AND SUBSCRIBERS:

All contributions for Publication, Books for Review, and Exchanges should be sent to the Editorial Office, 145 Gates Ave., Brooklyn, N. Y.

Remittances for Subscriptions and Advertising and all business communications should be addressed to the

ANNALS OF SURGERY,
227-231 South Sixth Street,
Philadelphia.