of alkaloidal matter when tested with the solution of iodine in potassium iodide. The alkaloid was found to be soluble in water, and was extracted from the aqueous solution by chloroform, just as in the case of the contents of the stomach described above. The residue, on evaporating off the chloroform, had precisely the same properties as that from the stomach contents, but the quantity of the oily liquid obtained was far greater—viz., 0.188 gramme (=2.9 grains). This residue was converted into oxalate by adding excess of oxalic acid to the aqueous solution and evaporating to dryness. The residue was then boiled with absolute alcohol, which dissolved the oxalate of nicotine; filtered alcoholic solution was evaporated to dryness, the oxalate dissolved in water, the solution rendered alkaline with potash and shaken with chloroform. The chloroform solution yielded on evaporation the pure nicotine, paler in colour than the original oil, but otherwise quite similar in properties. Of this purified alkaloid, 2 centigrammes (= $\frac{3}{10}$ ths of a grain) were injected in the state of aqueous solution into the pleural cavity of a large rabbit by my friend and colleague Mr. C. J. Martin. Almost instantly the animal was thrown into violent convulsions, during which it turned a complete somersault, and was observed to be in a state of extreme opisthotonos as soon as it reached the ground. This condition was rapidly followed by general muscular paralysis and relaxation, and the breathing stopped within three minutes after the exhibition of the poison. The thorax being opened at once, only fibrillar contractions of the heart were observed. At the adjourned inquest before the deputy coroner, Mr. H. E. Barnes, a relative of the deceased having heard my evidence, and my suggestion founded therefrom, that the deceased might have made an infusion of tobacco in some alcoholic liquor, and taken the mixture, informed the coroner that "in conversation the deceased had told him that that was how he meant to do it," from which it would appear that the man was cognisant of the highly toxic nature of tobacco infusions. As the bottle which the deceased threw away was broken, no clue whatever was given to me as to the nature of the poison. As regards the absence of convulsive movements in this case, such movements appear not to be invariably produced by tobacco intoxication in the human subject. Tetanic convulsions were present in one case recorded by Taylor, but absent in another. Possibly the quantity of the poison taken may account for these different effects, for no convulsion took place in the case observed by Taylor himself, where a man swallowed pure nicotine from a bottle, and died within five minutes.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF SPORADIC CRETINISM.

By ARNALLT JONES, L.R.C.P. LOND., M.R.C.S. ENG.

THE following notes refer to the case of R. G—, who suffered from sporadic cretinism, and died at the age of two years and a half, having been under my observation for nearly two years. I was first consulted when the child was ten months old, because the parents were alarmed about the condition of his tongue, which had grown large and unwieldy, protruding constantly through the lips and threatening to interfere with respiration. The child certainly had macroglossia, but he also had a cretinoid appearance. When seven months old he was able to stand upon his legs and also to articulate "dadda" and similar nursery calls, and he took an interest in things that went on around him. When nine months old, however, he commenced to have violent fits of crying, and a blight seemed to come over the child. The tongue now started growing till it protruded through the lips, keeping the mouth constantly wide open. After he reached the age of nine months he ceased to grow, and his appearance became very striking. The hair was coarse and scanty; the skin thickened, rough, and scaly in parts, the forehead shallow and face wide, the nose

broad and flat, the lips thick, with a wide open mouth constantly dribbling saliva, while the features were utterly devoid of expression; the hands were broad and rough-skinned, and the fingers stumpy. No trace of the thyroid gland could be made out, but there was a mass of fat in each supra-clavicular triangle. Some stress was laid by the late Dr. Hilton Fagge upon the presence of these collections of adipose tissue in this situation, and in this particular case these fat collections were at one time nearly the size of a hen's egg, but during the last few weeks of the patient's existence they dwindled down to the size of a pigeon's egg. The teeth were late in appearing; they were friable and ill-formed and crumbled away early. The child was subject to violent fits of crying; these came on at varying intervals, and if he once started he would go on perhaps for hours at a stretch. Otherwise, he accepted his fate in a placid, indifferent manner. Three days before his death violent coughing came on with hæmoptysis. The child died suddenly in convulsions, and, unfortunately, this event occurred during my absence on a holiday; consequently a necropsy was not obtained. The actual cause of the disease in this case is difficult to ascertain. The water drunk by the family was not impregnated with the salts of lime or magnesia; their house is built on pennant rock (sandstone), which here is some hundreds of feet in thickness; there was no consanguinity on the part of the parents, and no specific history on either side. The father was addicted to intemperance at times, a fact which would carry some weight with those who accept a theory as to the etiology of sporadic cretinism suggested by Dr. Langdon Down. This disease has been noticed more than once in several children of the same parents; and here that is likely to be the case, as a younger child of this family is at the present time developing some of the symptoms of incipient cretinism.

Aberavon, Glamorgan.

XANTHOMA MULTIPLEX.

By G. C. STEPHEN, L.R.C.P. LOND.

ON consulting Dr. Crocker's book on Diseases of the Skin, I find that only forty cases of xanthoma multiplex occurring after puberty have been recorded. I therefore trust the following notes of the case of a patient who recently consulted me for brow neuralgia and jaundice, associated with that interesting affection of the skin xanthoma multiplex, may prove of interest.

Miss —, aged sixty, states that she had always enjoyed fairly good health until from two to three years ago, when she suffered a good deal from bronchial and nasal catarrh and diarrhoea; but from these she appeared to have quite recovered until she was taken ill in November last with supra-orbital neuralgia and erysipelas of the right side of the face. In January, patches of xanthoma appeared on the hands, feet, wrists, and face, followed in a short time by very intense jaundice. As the patient was at this time residing in a malarious district of America, I think the jaundice was attributable to the malarial cachexia there acquired. When I first saw her, May 26th, 1890, there were numerous buff-coloured patches on the palms and backs of the hands and wrists, and in the flexures of the elbows and knees; also on the face there were several patches not raised above the level of the skin. The lesions on the hands and feet were from about the size of a grain of wheat to that of a three-penny piece. Some were perfectly flat, and on passing the fingers over them no protuberance was noticeable (xanthoma planum). Others, however, were slightly raised above the level of the skin (xanthoma tuberosum). The subjective symptoms were severe itching, particularly at the time of the formation of new lesions. I have repeatedly examined the urine without discovering any trace of sugar. Under anti-malarial treatment, and drachm doses of phosphate of soda three times a day, the jaundice has improved very much, and the irritation of the skin and neuralgia have almost entirely disappeared. The patient feels confident that some of the patches of xanthoma have disappeared, but I cannot say that I have noticed more improvement than that they have become less raised above the level of the skin. The patient's statement is no doubt important, however, in view of the fact that Mr. Jonathan Hutchinson, in a recent number of the *Archives of Surgery*, reports a case with symptoms very similar to those of my patient, in which xanthoma diabeticorum was suspected from the irritating character of the

lesions &c., although no glycosuria was discoverable, and in which he thinks there was sufficient improvement in the lesions to warrant a more hopeful prognosis being taken of the disease than the available statistics would lead one to suppose. Sutherland-avenue, W.

SIMULTANEOUS DISLOCATION OF BOTH SHOULDER JOINTS.

By T. W. J. ALLEN,
HONORARY SURGEON TO THE GRIMSBY HOSPITAL.

IN connexion with the accounts of simultaneous fracture of both clavicles, recently published in THE LANCET, I think the following case worth recording.

A young man, aged twenty-four, works on the Fish Dock, driving a cart, and packing fish. Within the last four years he has from different causes dislocated the right shoulder four times and the left shoulder three times, and on each occasion came under my observation. Ten days ago I was called in to see him, and found him suffering from a subcoracoid dislocation of both joints. He said that on getting hurriedly out of bed to assist an invalid brother he turned faint and fell, and upon recovering his senses found both his shoulders "out." I had no difficulty in reducing the deformity by the heel in each axilla in the usual way.

Great Grimsby.

CASE OF TRISMUS NEONATORUM.

By J. C. O'SULLIVAN, L.R.C.P.E., &c.

ON the night of July 3rd I was called to see the child of Mrs. H—, who gave it birth four days previously. The jaws were rigidly fixed and the finger could not be introduced into the mouth. There was a deep ridge on the forehead, giving the face an expression of frowning. The angles of the mouth were drawn outwards. The temperature was normal, and the umbilical cord had healed healthily at the place where it was divided. I told the patient's friends that tetanus was probably impending. The routine modes of treatment, warm baths, &c., were ordered to be tried. Convulsions came on the following morning at 2 A.M. These were strong, rigid, and continuous. The fingers were clenched on the palm, the back arched, abdomen upwards, and the head thrown back. There was much lividity, and the convulsive movements occurred five or six times an hour. The child died this morning (July 5th). There was no defective sanitation or other recognisable cause to explain the disease. The mother has been, and continues to be, healthy.

East-street, Walworth, S.E.

FIBROID TUMOUR OF THE CERVIX UTERI.

By PERCY BOULTON, M.D., M.R.C.P. LOND.,
PHYSICIAN TO THE SAMARITAN FREE HOSPITAL.

THE rarest position for a mural fibroid is in the cervix below the os internum, and the following is the report of a typical case which has recently been under my care.

F. C—, married, aged thirty-six, had been suffering from severe hæmorrhage almost continuously for six months, and when not bleeding had a copious discharge, with debility and backache. She had been seen by several practitioners who had told her she had a tumour, but a distinct diagnosis had not been arrived at. On exploration the vagina was found to be occupied by a tumour, more or less oval in shape and as large as a goose's egg. It clearly was not a polypus, as no encircling cervix could be felt, and the tumour first suggested to my mind a complete inversion of the uterus. High up in front above the pubes, a slit-like depression like an ordinary button-hole from $\frac{1}{2}$ in. to $\frac{3}{4}$ in. in length could be felt. I thought this might be the os, but to clear up the idea of inversion, I passed my finger into the rectum when the diagnosis was at once made apparent, for the uterus could be distinctly felt above the tumour, which was found to be entirely in the cervix. By the bimanual process this was confirmed, though not so unmistakably as by rectal examination, as the patient was fat. I then dilated with my finger what I had supposed might be the os. The anterior lip was exceedingly thin and the posterior lip effaced

by the tumour, so the finger passed backwards over its convex anterior surface. I made an incision with the cautery knife down the centre of the tumour, beginning at the os, and enucleated the growth, and the patient has made a good recovery.

The case is interesting because it is uncommon, but especially from a diagnostic point of view, for if the growth had been mistaken for a polypus, and it had been removed by the écraseur, Douglas' pouch would certainly have been opened; while if it had been treated as an inversion, the attempts at reposition might have led to pelvic inflammation or other trouble.

Seymour-street, W.

A Mirror

OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

CHARING-CROSS HOSPITAL.

CALCULOUS PYELITIS; RIGHT NEPHRO-LITHOTOMY; PELVIC ABSCESS; RECOVERY.

(Under the care of Mr. BELLAMY.)

REPORTS of all cases requiring operation on the kidney are of the greatest clinical value, and an accurate record should be taken of every condition and symptom, for we are but on the borderland of perfection in kidney surgery. Hardly any two cases are alike, and we learn from each, either from error or success. We think the case here reported of peculiar interest. The extreme difficulty of the operation, a peritoneal wound (absolutely unavoidable), the immense protrusion of bowel, and, as Mr. Bellamy suggested, a possible abnormal position of the organ, made the case a very unfavourable one. The subsequent pelvic abscess was possibly the result of the wound of the peritoneum, but, as will be seen from the notes, owing to the length of time after operation at which it appeared, this is doubtful. For the notes of the case we are indebted to Mr. Da Costa, surgical registrar.

T. L—, aged forty-one, a messenger, was admitted on March 18th, 1890. Had always been well. Enjoyed good health as a boy until sixteen years of age, when he was struck over the right loin, after which he was laid up in bed for ten days, for five days of which he suffered from hæmaturia. He recovered from this completely as far as he remembers, but had a slight return of hæmaturia eighteen months afterwards following unusual exertion. He was quite well until 1879, when he noticed a mucous deposit in his urine (in 1870 he had an attack of gonorrhœa, followed by swollen testicles; this attack lasted only ten days), which was ascribed by a surgeon to cystitis. He then went to Guy's Hospital, where he was treated in the wards for cystitis. He was under treatment for four months. The deposit has gradually been increasing and getting thicker. Four months ago he had bronchitis, and was then advised to wash out his bladder with boro-glyceride, a drachm to half a pint of water, twice daily. He did not derive any benefit from this. Four weeks ago he saw a surgeon, who sounded him for stone, but did not find any. Mother died of asthma when fifty. Father died of old age. All his brothers and sisters were healthy, no relative dying either of cancer or phthisis. The patient is well nourished. Heart and lungs are healthy; pulse 68; temperature normal. Urine: (a) Frequency. Passes urine four or five times daily, not at all during the night. (b) Character of stream: Passes a fair-sized stream, with good impulse. (c) Pain: Has slight pain on passing urine, especially at the end of micturition. Stream does not stop suddenly; with the first and last part of the stream urine comes away with a white turbidity. Patient has not passed any blood for the past twenty-four years. Examination of urine: Cloudy, and on standing yields considerable deposit; the odour is not offensive; sp. gr. 1012; slightly acid; contains about one-sixth albumen. Under the microscope it contains a large quantity of pus. The patient has