This article was downloaded by: [University of Florida]
On: 16 February 2015, At: 16:36
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Annals and Magazine of Natural History: Series 8

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah14>

XXXIII.—Description of a new cichlid fish of the genus *Heterogramma* from the La Plata

C. Tate Regan M.A.

Published online: 08 Sep 2009.

To cite this article: C. Tate Regan M.A. (1909) XXXIII.—Description of a new cichlid fish of the genus *Heterogramma* from the La Plata , *Annals and Magazine of Natural History: Series 8*, 3:15, 270-270, DOI: [10.1080/00222930908692574](https://doi.org/10.1080/00222930908692574)

To link to this article: <http://dx.doi.org/10.1080/00222930908692574>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

two perfectly distinct species, one western (*anceps*, Bougainville) and one eastern (*atrata*, Guadalcanar). The faunistic areas of the Solomon Archipelago indicated by the distribution of the five species of the *Pteropus rayneri* group are very nearly the same as those recognized by ornithologists (see W. Rothschild and E. Hartert, Nov. Zool. xii. pp. 243-244, 1905).

XXXIII.—*Description of a new Cichlid Fish of the Genus Heterogramma from the La Plata.* By C. TATE REGAN, M.A.

Heterogramma pleurotænia.

Depth of body $2\frac{2}{5}$ in the length, length of head $2\frac{1}{6}$. Snout shorter than eye, the diameter of which is 3 in the length of head; interorbital width $3\frac{3}{4}$ in the length of head. Depth of præorbital $\frac{1}{3}$ the diameter of eye. Maxillary extending a little beyond the vertical from anterior edge of eye; jaws equal anteriorly; fold of the lower lip continuous; cheek with 4 series of scales; no distinct gill-rakers on the lower part of the anterior arch. Scales $23\frac{2}{8}$; upper lateral line well developed on 8 or 9 scales only; lower lateral line vestigial or absent. Dorsal XVI 6; spines subequal from the fourth, the last $\frac{2}{3}$ the length of head. Anal IV 5. Pectoral nearly as long as the head. Caudal rounded. Caudal peduncle deeper than long. A dark lateral stripe from eye to base of caudal; a broad oblique blackish bar from eye to interoperculum; caudal fin with a few transverse bars or series of spots; outer edge of pelvic fin dusky.

A single specimen, 40 mm. in total length, from the La Plata, received from Herr J. Paul Arnold.

The different coloration, somewhat deeper body, lower spines, narrower interorbital region, &c. distinguish this species from the allied *H. corumbæ* and *H. trifasciatum*. A specimen of *H. corumbæ* in the British Museum has IV 5 anal rays, and it is probable that in the species here described the normal number is III 6-7.