DIPHTHERIA.*

By H. RENNEY, M.D., D.P.H.,
Medical Officer of Health, Sunderland.

WHEN I received an invitation from our worthy Secretary to read a paper at this meeting of the Northern Branch, I felt some difficulty in choosing a subject. Just about that time, however, we were having what was, for Sunderland, a great increase in the number of notifications of diphtheria, and I thought I could not do better than introduce a discussion on that disease, and the measures which might be taken in order to prevent its spread. I approach my subject with considerable diffidence, for I feel that I have not much that is new to tell you, and I would much rather be listening to one of you than standing here as a reader of a paper. However, I hope to learn a good deal from the discussion of this important subject.

Diphtheria was, at one time, comparatively speaking, a rare disease in Sunderland, and I believe also in the county of Durham; but of late years it has been considerably on the increase. Not only in the County of Durham has this been so, but in the whole of England and Wales, for according to the Annual Reports of the Registrar-General, the mortality has been much increased during the last decade.

In Sunderland, the average number of notifications of diphtheria (in which are included membranous croup), has been 48·6 for the last ten years. During the whole of last year we had only 74 cases notified, but in the first quarter of this year we have had 55 notifications, so that it is time something was done to endeavour to limit the spread of the disease.

On investigating these 55 cases, I found that most of the children affected attended the Council Schools. Nine cases occurred in one class-room of the infants' department of one of the schools. On March 21st, the class-rooms and cloak-rooms of this department were disinfected by means of formalin spraying and formalin vapour. Since that date there has only been one case in the department, viz., on March 24th, and it had probably become infected before the disinfection was carried out.

I do not wish to assume that the source of infection was to be found only on the floor, walls, and desks of this class-room, but I think it is a very significant fact that since March 21st there has not been another case there. I had not an opportunity of taking swabs from

* Read at a Meeting of the Northern Branch of the Incorporated Society of Medical Officers of Health, May 12th, 1905.

the throats of the unaffected children, and I cannot therefore say if any of them acted as "carriers." I had intended taking swabs from the dust on the floor and from the top of the door previous to disinfection, but my disinfector had done his work before I was aware of it, and thus probably prevented me acquiring some interesting information. I took swabs from these places after the disinfection had been completed, and the result was negative. Of the nine children who were attacked, the first case occurred on February 17th, and the others in the following order, viz. March 4th, 9th, 13th, 14th, two on the 15th, 16th, and 24th. How the first case originated I am unable to say. I have not been able to discover any infected area in close proximity to the school. Slates were used for writing on, but there was no provision made whereby each child got its own slate every day. There had been no cases of diphtheria for a considerable time in this school previously.

In an epidemic of diphtheria which occurred in Bradford during the earlier part of 1903, according to information gleaned from a paper read by Dr. Crowley at the Glasgow Congress last year, swabs were taken from 93 throats where no clinical evidence of diphtheria existed, nor even sore throat. Out of the 93, there were diphtheria bacilli found in 42, that is, in a little over 45 per cent. If then, in every epidemic in schools, there are bacilli in anything like this proportion in healthy throats, or at all events in throats where no evidence of diphtheria exists, we have a very potent factor which makes for the spread of the disease; especially in the infants' departments, where children of susceptible age are crowded together under conditions which are by no means as sanitary as they might be. There is now no reason to doubt that Klebs-Löffler bacilli are found in healthy throats, and that these act as carriers of diphtheria to others, whose vitality, from one cause or another, is lessened. I believe that the use of pens, pencils, slates, and drinking-cups in common is a very frequent means of infection. The dust of class-rooms in the joints between the floor boards, on the top of doors, and in other out-of-the-way places, no doubt forms a nidus for bacilli.

Let me here relate what took place recently at our borough sanatorium for infectious diseases. On November 25th, 1904, a boy was admitted suffering from a typical attack of scarlet fever, the rash being well out on admission. He had, however, a very suspicious throat, and I at once injected 2000 units of anti-toxin, and had him removed to a pavilion by himself. Swabs from his throat were positive. He did exceedingly well, and a month later was returned to the scarlet fever pavilion, as he was still peeling. Previous to his re-admission this pavilion had been thoroughly disinfected with formalin. On January 22nd, 1905, a nurse who had been

working in this pavilion, in the male ward, developed diphtheria. She was removed to the isolation block, and disinfection was again carried out. On January 31st, a boy, who had been admitted into the scarlet fever pavilion on December 22nd (after the pavilion had been disinfected owing to the other cases), developed diphtheria. He also was removed, and disinfection was again carried out. On February 6th, Klebs-Löffler bacilli were found in the healthy throat of a child in the same ward, but nothing came of this. On March 24th, yet another case of diphtheria appeared in another scarlet fever patient who had been admitted on March 9th. He was removed, and the pavilion was emptied of the rest of the patients. Swabs were then taken from the dust on the top of the door, and also from crevices in the floor. Diphtheria bacilli were found in both instances. Thorough disinfection with formalin was again resorted to, and swabs were taken as before. The result was negative. I have no doubt that a certain percentage of scarlet fever patients have Klebs-Löffler bacilli present in the throat on admission, and although the majority of these do not develop diphtheria, yet they act as carriers to others. I think it is a matter for serious consideration whether all scarlet fever patients should not first be admitted into an observation pavilion, where swabs could be taken, and the patient detained until the swabs showed a negative result; they could then be sent to the scarlet fever pavilion.

We know but little of the etiology of diphtheria. One bacteriologist has said that probably the bacillus is ubiquitous; if that be the case, there is no difficulty in understanding how it is that the disease occurs in so many different localities, and in different sections of the large towns. The curious thing about its distribution is, that it is found so frequently to be much more prevalent in the newer houses where sanitary defects do not exist, and this is not necessarily on "made" ground, or indeed, on any particular kind of soil, although a clay soil has been said to be more favourable to it. We need not consider the question of a milk-borne epidemic, or the occasional outbreaks which have been thought to be due to the disease in the cow itself, as when such outbreaks occur they can generally be traced to their source. As far back as 1877, Dr. Thorne, in his Milroy Lectures, gave it as his opinion that school influence operated in a number of ways, by bringing together children at the most susceptible age, and placing them in elementary schools, where, for long periods, they inhale the exhalations of their fellows. He thought that such aggregation induced a form of sore throat. One has only to attend the practice of a large throat hospital for some time to see what a large number of children have pharyngitis and tonsillitis, and these must form only a minor proportion of the total number who are suffering in a similar manner, so that

there must be in every community a great number of children who are, more or less, susceptible to diphtheria.

The infection may be carried, as we know, in milk, by fomites, and some have thought, probably by domestic animals and even birds. Aerial convection I very much doubt, if by that is meant convection by winds. There is an aerial dissemination, however, that I think is possible. It has been proved that in coughing, sneezing, and, I believe, in speaking also, tubercle bacilli are projected from the mouth some few feet, and I do not see why the same thing should not take place when a person suffering from diphtheria coughs or sneezes. In this way it may be spread in schools, not only by the so-called "carriers," but by children suffering from an undetected mild attack. It has been thought that the bacillus may find a habitat in the surface soil, and that a rise in the ground water may, by expelling the ground air, lead also to the expulsion of the bacillus; but this is going back to the old theory that micro-organisms float in the air independently of any vehicle. It is not likely that the ground air will contain the little rafts of dust on which the bacilli could float. Besides, diphtheria has been known to have increased during dry years when there was no rise of ground water.

We shall never be able to successfully combat and destroy this disease until its etiology is fully worked out, but we can take action against it in a way which, I think, will materially lessen its incidence. I mean with regard to prohibiting children attending school at the most susceptible age period. Statistics show that the incidence and mortality increase up to the fourth year of life, and thereafter steadily decline. The Education Department certainly makes provision for children attending school from the age of three to five years, but does not *compel* them to attend until they reach the latter age. If then, we could prevent them attending school until this age were attained, I think there can be no doubt that there would be a very considerable reduction in the number of cases of diphtheria. I presented a report to my Health Committee recently, in which I pointed out the greater susceptibility at this age period, and I recommended that the Education Committee should be asked to endeavour to prevent children under the age of five years being admitted to school. I am glad to say that the Health Committee thereupon passed a unanimous resolution that my report be sent to the Education Committee, and that the Chairman and I should go as a deputation to further support their resolution*. On my suggestion, the Health Committee have also

* Unanimous resolution passed by the Education Committee to ask the Education Department to prohibit children attending school under the age of five.

requested the Education Committee to take steps to prevent any child attending school for at least one month after having had diphtheria, a similar regulation to apply to children living in the same house with an infected case*.

As further precautionary measures I would suggest the abolition of slates, and the substitution of paper for writing on. Children should be taught that it is dangerous to suck pencils, pens, etc., and the interchange of such articles should be forbidden. Drinking cups should be abolished, and the children forced to drink only from the running stream of water from the tap. Every Education Authority should have a specially appointed Medical Officer who would examine the children at regular intervals, and who would thus be in a position to detect the first signs of the disease ; in addition to which he would take swabs from such throats as he thought necessary and have them examined. There is no doubt there are, from time to time, children attending school who have a mild attack of diphtheria ; these would be excluded by such inspection, and isolated either in an infectious diseases hospital or at home.

If, in addition to these precautions, every child suffering from diphtheria were refused admission to school until the throat were free from bacilli, outbreaks of the disease would be very much diminished.

DISCUSSION.

The paper was very fully discussed, and the discussion elicited the fact that during the past five or six years diphtheria had increased in frequency alarmingly in the County of Durham. The increase of new houses built on previously cultivated soil on unpaved sites, was suggested as one reason for the increase of this disease. School life, especially the younger lives, was the means of disseminating diphtheria markedly, and that the use of school furniture, such as slates, etc., was probably one of the chief agencies in the school at work in spreading diphtheria.

There was no proof that drinking water entered into the causation of diphtheria.

As means of prevention there was unanimity respecting the suggestions made in the paper.

The increased incidence during the colder months of the first and fourth quarters of the year compared with that in the warmer second and third quarters, was also remarked.

PROFESSOR KOCH ON TSETSE FEVER.—The *Cologne Gazette* publishes a letter from Professor Koch, who went to German East Africa at the end of last year to examine the causes of tsetse fever, recurrent fever, and plague, on behalf of the German Government. In the letter, which is dated Iringa, Uhehe, April 26th, the Professor announces that he has succeeded in ascertaining the existence in the tsetse fly of trypanosoma, confirming the observations of British workers. His investigations show that this parasite undergoes special development in the tsetse fly.

* This also was agreed to by the Education Committee.