been the subject of intestinal obstruction from extreme atony of the bowel, was quite convalescent, and has ever since remained well.

These cases, amongst others which I have met with, fully illustrate the importance of a line of treatment so ably advocated by Dr. Ogle in his monograph on the treatment of abdominal distension by acupuncture. Having had such satisfactory experience of its utility, and never yet having witnessed any bad effects from it, I place these results before the profession in advocacy of its employment.

Newcastle-on-Tyne.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE IN WHICH A TRANSIENT ATTACK OF LOCAL ASPHYXIA (RAYNAUD) WAS DETERMINED BY A DOG-BITE.

BY WILLIAM PASTEUR, M.D.,

ASSISTANT PHYSICIAN TO THE MIDDLESEX HOSPITAL.

HERBERT B—, aged nine, a very anæmic but well-nourished boy, of lymphatic temperament, was bitten on the back of the left hand by a puppy-dog on April 23rd. His previous health had been good on the whole. He had snuffles in early childhood, and is now suffering from chronic ulceration of the back of the pharynx, which is probably syphilitic. He has never been subject to cold hands and feet, and is certain they never turned blue. There is no history of any urinary trouble.

The wound, which was quite superficial, was cauterised and poulticed. On one or two occasions during the first two days his mother noticed that the left hand was slightly blue. On April 28th the wound began to suppurate, and on the same day the whole hand became icy cold and very painful, and turned a dark-blue colour. On the application of a warm poultice it soon regained its normal state. The boy was admitted under my care at the North-Eastern Children's Hospital on April 30th. When seen by me at 2 o'clock the left hand was slightly œdematous, and of a deep blue-black, inky colour, especially well seen in the matrix of the nails. The hand had been wrapped in cotton-wool, and was no more cold. The discolouration reached half-way up the forearm, and was of a uniform colour. There was no venous mottling above the level of the discolouration. The radial pulse on the affected side was decidedly less full than in the corresponding limb. The patient complained of considerable pain in the hand. There was also very slight lividity of both ears. The right hand and arm were perfectly healthy. Some urine passed shortly after admission was quite normal. It was tested every day for a fortnight for albumen and blood, with negative results. There was a deposit of urates on the second day after admission. Two hours after admission the attack had completely passed off. On the morning of May 6th the left hand became slightly blue and cold, and somewhat painful. The attack lasted only ten minutes. Later in the day a mild attack of local asphyxia was induced by immersing the hand in cold water for some minutes. Sensation was never blunted to any appreciable extent during the attacks. There were two very mild attacks on May 8th and 9th. On the 12th both hands were immersed in cold water for fifteen minutes, with negative result. No difference was observed on the two sides. On the 13th, after being washed in warm water, the left hand became blue and cold for about an hour. During his stay in the hospital the temperature varied between 97.6° and 99.8°. The wound healed rapidly under suitable treatment, and the patient was discharged cured on May 14th. Three days later he had a well-marked attack affecting the whole hand for about an hour. The left hand is now often colder than the right. On May 26th, after a warm bath, the left foot became cold and the great toe turned blue. Since that date there have been no attacks. The boy is again in the hospital on account of the pharyngeal ulceration. I am indebted for the notes of the case to Mr. E. Brock, M.B.

Remarks.—The anæmia and syphilis may be held to have been predisposing causes in this case, but there seems no reason to doubt that traumatism was the immediate determining cause of the attack. Although it would be profitless to speculate on the bearings of this observation, one may legitimately draw attention to the limited distribution of the asphyxia, to its asymmetry, and to its primary location in the wounded extremity. It is also of interest to note, in connexion with the reflex vaso-motor neurosis theory of Raynaud's disease, that in the present instance there was a well-defined source of irritation in that part of the periphery which subsequently became the seat of the asphyxia.

Queen-street, Mayfair.

ON THE TREATMENT OF HYDROCELE.

BY A. E. HIND, F.R.C.S.

THE treatment of hydrocele of the testis by injection of solution of perchloride of mercury was discussed, I believe, in the medical journals some years ago, but has not yet received the attention it deserves. In my hands it has always given complete satisfaction, but my cases have only been a limited number. I use a solution three times the strength of liquor hydrarg. perchlor., and first draw off the hydrocele fluid and then wash out the cavity with the perchloride solution and draw off any remaining fluid. The last case on which I operated was an old double hydrocele with thick walls. The result was quite satisfactory. The injection does not cause pain, and is not followed by inflammation. In these respects it is superior to iodine. Its action depends on its power of forming an insoluble albuminate of mercury, by which the walls of the cavity are glued together. As far as my experience goes, there is no danger of mercurial poisoning. The albuminate is fairly insoluble, and a large amount of mercury is not required. I have tried this treatment with success in a case of infantile hydrocele after tapping had failed. Iodine causes pain when used, and often much pain afterwards by setting up more inflammation than is required to cure. Iodine also is not always efficacious, and may be followed by abscess. I believe perchloride is sure in its action, and its use does not necessitate rest in bed afterwards. I should like to know if the experience of others who may have used it coincides with mine. Have any ill results been known to follow? Has it ever been known to fail? What is the weakest solution that is effectual?

Jersey.

A CASE OF PUERPERAL CONVULSIONS TREATED BY ADMINISTRATION OF OPIUM; RECOVERY.

BY R. E. JOHNSON, L.S.A.

I WAS called on Christmas morning, 1887, at 1 A.M., to see a young woman, aged sixteen, reported to be suffering from convulsions. Her confinement was expected, so I administered twenty minims of tincture of opium. This had the effect of soothing the patient, who was in a highly excited state after the first attack. In half an hour another fit, decidedly epileptiform, came on. I gave the patient fifteen minims of the same drug, and made a vaginal examination, when I found the foetal head well down, nearly on the perineum, the os being widely dilated and the membranes unruptured. She had another fit and then the child was born. Immediately after its birth she was seized with a severe rigor. This was at 3 A.M. Being a long way from home, I did not dare to leave her to go for any other remedy, so persevered with the opium, at last getting her well under the influence of the drug. At 6 A.M. the fits, though still occurring about every twenty minutes, were not so severe in character. At 10 A.M. I was informed that she had had seven fits since 6 o'clock. There was profuse diaphoresis. About four ounces of urine, containing a little albumen, were drawn off by the catheter. 12 noon: Two more fits. Still administering opiates. 7 P.M.: Has had eight fits since noon. 10 P.M.: Had one fit since last seen. Had one while I was present. Catheterisation: about eight ounces of urine.

Dec. 26th.—10 A.M.: No fits since last night. Adminis-