Enteric fever stands on a different basis, for the principal measures for the prevention of this disease are those tending to the general improvement of the sanitary condition of the county. Its presence, unless imported, is an indication of some insanitary condition, and we should not rest satisfied until this disease has been exterminated. The cause of the actual decrease of typhoid fever may, I suppose, be put down to improved sanitary conditions and to the increased cleanliness and carefulness in living of the people.

The greatly decreased mortality from phthisis is a matter for congratulation, although it is probable that some of the earlier deaths entered as phthisis were not due to that disease.

The rapidly increasing number of deaths from cancer is the one discouraging feature of this table.

SCARLET FEVER AND MILK SUPPLIES.*

By EDMUND M. SMITH, M.D. (Ed.), D.P.H. (Camb.),
Medical Officer of Health for the City of York.

At the end of October, a sudden outbreak of nine cases occurred in the Clifton District, entirely amongst *adults, and in better class homes*, such as are rarely invaded by this disease.

Every line of enquiry was followed up, but no other common origin of such an unique outbreak could be discovered except one, viz. :—That all the patients were accustomed to drink raw milk, and that all obtained their supply from the same milk-dealer, whose reputation had been excellent.

Full investigation failed to discover any other source of infection through the milk than that of a labourer, two of whose children were convalescent from attacks of scarlet fever, and who had not cleansed his house properly after the infection, according to our printed instructions. This man had been employed on the said milk-farm for about a week prior to the outbreak, his employer being ignorant of the man's home circumstances. He was not supposed to have anything to do with the milk or milking, and he and others on the farm denied that he had, but his home conditions, and connection with the farm, taken together with the very uncommon features of the outbreak (its simultaneousness, the class of patients, and the common milk-supply), made the milk supply an extremely suspicious circumstance. Although the milk dealer's custom was an extensive one, and the number of persons known to be attacked very few, yet, under all the circumstances, I cannot think that the relationship was a mere coincidence; but proof seemed impossible.

Upon realizing what the possible source of infection was, the labourer's employment on the farm was at once stopped, measures were taken to enforce the cleansing of his house, and raw-milk consumers were advised to boil the milk before use.

The outbreak then ceased.

Presuming that the milk supply was infected, the reason why so few persons were attacked is somewhat difficult to explain, but many of the customers were known to boil or sterilize their milk before use, and many adults do not drink raw milk. It is probable, also, that only some small portion of the milk supply was infected.

* Annual Report, 1904.