therefore, the approximation to the security assured by the artificial member, the greater will be the improvement in locomotion. This is a fact not generally recognized in the construction of the ordinary brace.

It is not my purpose to urge operative treatment for all cases of calcaneus, but simply to suggest an effective procedure for those cases in which, for one reason or another, mechanical treatment is inefficient—a class which, in hospital practice at least, is a very large one.

THE DIFFICULTIES IN MAKING A DIAGNOSIS IN THE BONE LESIONS OF NURSINGS.¹

BY R. TUNSTALL TAYLOR, B.A., M.D.,

SURGEON IN CHARGE OF THE HOSPITAL FOR CRIPPLED CHILDREN; FELLOW OF THE AMERICAN ORTHOPEDIC ASSOCIATION; ASSOCIATE PROFESSOR OF ORTHOPEDIC SURGERY IN THE UNIVERSITY OF MARYLAND, ETC.

The title of this paper suggested itself to me on account of the paucity of symptoms presented by several cases in babies in my wards during the past year which rendered an early and definite diagnosis most difficult and unsatisfactory. I present it in the hope that the discussion which it may bring forth from my fellow-members, together with observations I myself may make, will add somewhat to our means of making a more definite and an earlier diagnosis.

By nurslings I have meant to convey those babies under two years of age who are either breast-fed or bottle-fed, and in whom dentition has not begun or is only partially completed. However, the limitation need not be made, as many of the symptoms and blood changes to be spoken of occur in young children as well as infants.

On being called to see a baby with one or more of the cardinal symptoms of bone inflammation certain of the following formidable array of diseases may present themselves to the clinician for differentiation and exclusion, viz.: tuberculous osteomyelitis or epiphysitis, syphilitic periostitis and osteochondritis, achondroplasia, rickets proliferation, softening or clubnation, scorbutic subperiosteal hemorrhage, or, perhaps, epiphyseal separation, osteosarcoma, epithelioma (?), rheumatism, "growing pains," rheumatoid arthritis, acute anterior poliomyelitis, acute osteomyelitis, acute arthritis, and also influenza, typhoid infection, colon infection, and pneumococcus infections; osteomata, whether exostoses or hyperostoses; osteopsathyrosis, spontaneous fractures, and cysts.

¹ Read before the American Orthopedic Association, 1901.

The following is a case in point, involving a differential diagnosis between tuberculous osteomyelitis, specific osteochondritis, the subperiosteal hemorrhage of scurvy, and sarcoma:

CASE I.—C. C., a boy, aged five months, was put under my care on January 1st of this year. The child seemed in great pain in the right leg, which was held flexed at about 45 degrees, rotated outward, and abducted the same amount. The mother stated that the child had had pain apparently for three months—i. e., from the time he was two months old—and she noticed he had held his leg constantly still and moved the left leg freely, so that she thought the right leg must be paralyzed.

Family History. The parents were refined and well-to-do, apparently healthy people. They had been married ten years before any children were born to them, but any miscarriages were denied. The father had led a rough life on the plains, and had been more or less dissipated. He acknowledged drinking to excess, had had venereal warts, but denied syphilis. He had had no rash, sore-throat, or headache, etc., at the time of the appearance of the venereal warts, and had received no medication for a prolonged time. (Further data on this point was unobtainable.) He is subject to "rheumatic gout." The patient has a brother and sister who are said to have always been strong and well. A paternal uncle died of phthisis, and a maternal uncle had a sinus in the back. The first wife of the patient's father died of consumption eleven months after marriage. Any history of malignant tumor in the family is lacking.

Previous History. The labor was easy and normal, and the child at birth was large and apparently healthy. No definite history of "snuffles" or rash is obtainable; but the baby had, when one month old, a large red, ecchymotic spot on the left cheek, which shortly disappeared. When two months old a swelling appeared at the left wrist, with malposition of the hand, which the mother thinks involved the joint. This was treated by the family physician with pasteboard splints and mercurial ointment, and followed by recovery; but some thickening is still easily detected at the lower left radial epiphysis. The feeding had been combined bottle-feeding and nursing. At the time of admission the child nursed twice daily, and was bottle-fed on equal parts of milk and water, *sterilized*, every three hours.

Pain and night-cries had been present for two months before I saw the patient, and insomnia has been a marked symptom. The maximum temperature, reached a few days before admission, was 103° F., but did not exceed 100° F. after that. As to any history of trauma, the father thought a former nurse was of such a nature that she would have unhesitatingly struck the child or wrenched an arm or a leg in anger, and attributed the child's condition to this; but he had no definite knowledge to this effect.

Condition of child on admission to the hospital, January 8th: On inspection a plump, well-nourished and well-muscled baby, and large for its age, was seen, weighing eighteen pounds two ounces. The lips, mucous membranes, cheeks, and anils were of a good color. The skin everywhere seemed normal, with no scars or discolorations suggestive of ecchymoses or subcutaneous hemorrhages. The gums were normal in appearance, and no teeth showed signs of erupting. The head was of good shape, with fontanelles open, but not protuberant. Chest and

abdomen were normal in shape, and there was no heading of ribs or enlargement of epiphyses except as above stated (*viz.*, left radius). The eye at once noticed the malposition of the right thigh and a distinct, fusiform, diffuse, spindle-shaped swelling at the upper third of the thigh, measuring $10\frac{1}{2}$ inches (28 cm.) in circumference against 10 inches (25.5 cm.) on the left. This swelling was said to have increased rapidly of late.

On palpation the lymphatic glands were not appreciably indurated anywhere except in the right inguinal region and right iliac fossa. The spleen was palpable, and extended 2 cm. below the costal margin. Percussion also disclosed this. Auscultation was negative except for a slight bronchitis involving the larger tubes.

The swelling in the right femur seemed to extend from about the middle third of the thigh to a point near the coxo-femoral articulation, but not into it. It was larger above than below, hard, did not fluctuate, and involved apparently more the anterior and external aspect than the posterior and internal. Muscular spasm was marked, and any movement seemed to cause pain and brought on a crying spell. The other joints did not show this tenderness. One point on the external aspect seemed to be more sensitive than the rest. A radiograph was made, which showed an hour-glass shape in the bone, with straight, regular edges. The blood count gave: Reds, 4,100,000; whites, 18,400; hemoglobin, 80. The differential count showed lymphocytes increased; no myelocytes or eosinophiles. Beginning poikilocytosis was noted. The diagnosis, so far as the blood was concerned, showed a slight secondary anemia.

With this data before me I at first thought of two possibilities—scurvy and tuberculous osteomyelitis. For the former I gave modified Pasteurized milk, lemon-juice, and beef-juice; and for the latter employed traction in the line of the deformity of $1\frac{1}{2}$ to $2\frac{1}{2}$ pounds, which seemed to afford almost immediate relief. On the third or fourth day after the traction was instituted the baby seemed perfectly comfortable, slept all night, appetite became vigorous, and increased weight was shown. I reluctantly gave the child for three successive nights tuberculin from Trudeau's laboratory, the most reliable I could get, with absolutely negative results. I say "reluctantly," because I felt if the result had been positive a bronchial lymph node might be responsible, and not the thigh trouble; and, beside, the usual site for tuberculous invasion—the epiphysis—did not seem the location of the trouble. If negative, we all know tubercula is not infallible, and a wrong inference might be obtained. The picture was not that of a true tuberculous coxalgia, and the child's age and the rapidly increasing induration was against such a diagnosis. The blood-findings, according to Dana's forty-one published observations,¹ led me to think the high leucocyte count would indicate tuberculous abscess formation; but Brown,² after an

¹ Boston Med. and Surg. Journ., May 23, 1896.

² Transactions of California Medical Society, 1897.

examination of seventy-two cases, dissents from this conclusion unless there is a secondary infection; "simply increased activity of the tubercular process itself," he claims, "would not give an increased leucocyte count over 2000 to 3000 above normal." Thus my diagnosis of tuberculous osteomyelitis was most unsatisfactory and by no means conclusive.

My diagnosis of scorbutus also lacked many usual symptoms to sustain it. The gums were normal, no subcutaneous hemorrhages had appeared, and there was no anemia to speak of. The child was well nourished, and the other limbs showed no tenderness. The only possible straws for the scurvy theory to hold to were the possible subperiosteal hemorrhages and the fact that many of the best authorities, such as Starr, Rotch, Lewis Smith, H. L. Taylor, and others, observe that scurvy shows no bleeding gums prior to dentition. Aspiration brought out only a few drops of blood, which was normal in appearance and not black or disintegrated. Thus following the old whist rule, "when in doubt, lead trumps," I ordered mercurial inunctions. After one week of this treatment the leucocytes had fallen to 16,800, but after two weeks the swelling had increased one-half an inch (1.5 cm.), and was appreciably harder, and my diagnosis of congenital syphilis was shaken; however, as the child seemed so much improved in every other way, I persisted in the inunctions. I then called in an eminent consultant, who suggested sarcoma, and advised an exploratory incision, with hip-joint amputation if sarcoma was found; but he felt possibly metastases had already taken place; but if not, and sarcoma should be found, permission should be obtained, in advance of any puncture, to go ahead with the amputation, for fear of metastases from the puncture. The child looked so rosy, bright, and merry that I determined to trust to "the trump" a little longer, with the result that the patient was discharged from the hospital on February 21st, with no induration in the right thigh and a freely movable joint. His general condition was excellent. To my mind it was a clear case of syphilitic osteochondritis. Carl Beck's paper on "Cartilaginous Cysts Near the Epiphysis of Long Bones in Children" had not then appeared, else I should doubtless have added this disease to my difficulties; but he tells us the skiagraph differentiates this condition from osteosarcoma, in that the former gives a translucent appearance at its situation, with straight, sharply marked edges of bone, while sarcoma gives no such translucency and an irregular, external demarcation. Wiegel has called my attention to the mottled or spongy look in his excellent skiagraphs of sarcoma in bone.

Ridlon, in vol. iv. of the *Transactions of the American Orthopedic Association*, speaks of a number of syphilitic bone and joint lesions seen by him, and quotes from Burnstead and Taylor's *Venercal Diseases*

(fourth revised edition, p. 735) in support of his diagnosis, which had been questioned by eminent authorities and pronounced tuberculous, although his results with antisyphilitic medication were quite suggestive.

To what extent blood examinations will help us remains to be seen, and with the vast material at our command it would seem to me well to make blood examinations a part of our ward records in bone diseases, just as urine has been heretofore. This seems especially so, in our special work, when one thinks of the important part played by the bone-marrow in the genesis of blood. Such records should bring forth fruit in due season and increase our means of differential diagnosis. In attempting any inferences from the blood-finding in infants we must bear in mind that a normal leucocytosis exists, according to Schiff, Gundobin, Bayer, Hnyem, and others, for we may have at birth from 17,000 to 21,000 white cells, and after the first feeding a digestion leucocytosis as high as 36,000. On the tenth day 10,000 to 14,000, and at the sixth month 12,000. After a meal 30,000 leucocytes is never a very high count in infants under two years. Cold baths likewise increase the leucocyte count. It is also to be observed that a backward child's blood is more of the infantile than of the adolescent type.

Cabot gives us an excellent foundation on which to build in *Clinical Examination of the Blood* (third revised edition, 1898), and I will refer to it for certain points on the blood in infancy and childhood:

"1. All signs by which sickness is shown in the blood of adults are exaggerated in children.

"2. All forms of infantile anemia are apt to be associated with enlarged spleen.

"3. In infantile blood the small mononuclear elements are more numerous than in the adult ('lymphocytosis of infancy').

"4. The red cells show variations in shape and size, and nucleated reds may be found.

"5. The hæmoglobin is relatively high at birth and relatively low during childhood."

It is therefore important to make our observations before menle or a cold bath, to state the child's age, and, if possible, obtain a blood record before the sickness began, to determine how much is physiological and how much pathological.

To apply this practically for diagnostic purposes in tubercular osteitis or periostitis without abscess formation (according to Dnna's observations), and without secondary infection (according to Brown's), we have no leucocytosis, while in syphilitic bone disease we have the leucocytosis of a secondary anemia. The same is true in scurvy and rickets. Acute osteomyelitis can be distinguished from so-called "rheumatic paine," "growing pains," and neuralgia by the leucocytosis present.

The average leucocyte count for osteosarcoma in adults (?) is high—about 17,000. Injections of tuberculin produce leucocytosis; therefore blood counts should not be made immediately after such injections. Acute rheumatism gives about 16,000, subacute 9000, and chronic 7000 leucocytes, on an average.

Hereditary syphilis is perhaps the best-known cause of a relative lymphocytosis in children, but scurvy may produce the same result.

In infancy the percentage of eosinophiles is very often higher than in adults, so that in them eosinophilia may be considered physiological; but it is also very characteristic of malignant bone disease and osteomalacia, as found by Neusser and his pupils. Cahot, from whom I have obtained most of the foregoing, mentions having seen eosinophilia in osteomyelitis also. In some cases of syphilis and syphilitic disease of the spinal cord we find eosinophilia.

Myelocytes are found especially in malignant bone disease, hereditary syphilis, and acquired syphilis and other conditions where there is a grave anemia with leucocytosis.

Normoblasts show more a severe secondary anemia, and megakaryoblasts show a very severe secondary anemia in bone disease as elsewhere. Applying this data to the case cited, we find the infant had 18,400 leucocytes and relatively increased lymphocytes, which would suggest exclusion of tuberculous bone disease in favor of syphilis or scurvy. Absence of nucleated reds, myelocytes, and eosinophilia, together with no marked diminution in the red count, would tend to exclude sarcoma from the preponderance of authority, especially as this was not what might be called a small tumor.

Thus, if this reasoning is logical, we would narrow down to syphilis and scurvy, and as the supposed subperiosteal hemorrhage was the only likely symptom of scurvy, hereditary syphilis alone remained.

Certainly, it seems to me, the blood examination was of value as an additional aid in the diagnosis of this case.

I will cite briefly another perplexing case.

CASE II.—H. R., aged eighteen months, son of a laborer, was admitted to the Hospital for Crippled Children, April, 1901, with a marked swelling extending from the upper border of the right hypochondriac region to the lower border of the right iliac region, and from the mid-axillary line around the right side of the loin nearly to the spinal column. The duration of this abscess was said to be four weeks; that is, the parents had noticed it for that length of time. In the lumbar region fluctuation was detected, and the pus seemed near the surface. The child was anæmic, emaciated, tubercular looking, and evidently in pain. Large furuncles were present on the occiput and right eyelid. The temperature was 101° F. The spine was rigid and double. Psoas-contraction of some 90 degrees was present, but no kyphosis or other deformity was apparent. Family history was good. The patient had been healthy up to September, 1900, when he had measles and whooping-cough, and had

a traumatic history of having fallen down five or six steps and off a chair previous to the onset of the present trouble, a month ago. His father stated that in the morning the child was apparently well, playing and talking, and in the evening could not use the right leg. Pain had been constant since, with fretful nights, anorexia, thirst, fever, sweats, constipation, but no vomiting. Palpation in right iliac fossa gave no induration in the region of the appendix.

The symptoms and picture were those of Pott's disease, with double psoas-contraction and lumbar abscess, but no kyphosis.

The blood count briefly showed 40,000 leucocytes, a low red count, and low hæmoglobin, increased polymorphonuclear neutrophils and small mononuclear leucocytes. Quite a number of cells resembling myelocytes and a few normoblasts were also found. This was against tubercular abscess in the extremely high leucocyte count and the presence of marrow cells found, as an incision demonstrated, evacuating 500 c.c. of thick greenish pus, sometimes mixed with blood, from which cultures showed the presence of the staphylococcus pyogenes aureus. The abscess cavity was curetted and drained, and the child showed rapid improvement. The psoas-contraction gradually yielded to traction in the line of the deformity.

THREE NOTEWORTHY CASES OF BRAIN INJURY.¹

By GEORGE TULLY VAUGHAN, M.D.,

OF WASHINGTON, D. C.,

SURGEON UNITED STATES MARINE HOSPITAL SERVICE; PROFESSOR OF SURGERY, GEORGETOWN UNIVERSITY; SURGEON TO GEORGETOWN UNIVERSITY AND EMERGENCY HOSPITALS; MAJOR AND DEBIADE SURGEON, UNITED STATES VOLUNTEERS, DURING THE WAR WITH SPAIN.

From the earliest dawn of medical science injuries of the brain—the seat of life and of intellect, if not the soul—have furnished an interesting field for study. The correct explanation of concussion—the mechanism by which laceration and hemorrhage take place without fracture, the laws which govern the direction of the lines of fracture, the prognosis as to life and health and intellect, and the best methods of treatment, are problems which have attracted the attention of surgeons and physicians from the days of Hippocrates, and are still burning questions for consideration by the surgeons of our day.

The dangers in cerebral injuries are shock, hemorrhage causing compression or anæmia, and infection.

The symptoms of shock occur immediately; those of hemorrhage at the same time or a few hours later, occasionally as much as a week later, and those of infection from two days to a week, or even several months or years after an injury, possibly as a result of a second injury which arouses a focus of infection which has long remained dormant.

¹ Read at the meeting of the Association of Military Surgeons of the United States, 1901.